

# Non-Hermitian (NH) Topology – What Changes?

Siddhant Midha

Department of Electrical Engineering  
Indian Institute of Technology Bombay

EE 787 Course Project  
November 28, 2022



# Introduction

- Hermiticity – Crucial?

# Introduction

- Hermiticity – Crucial? Guarantees,

# Introduction

- Hermiticity – Crucial? Guarantees,
  - 1 unitarity of the time evolution

# Introduction

- Hermiticity – Crucial? Guarantees,
  - 1 unitarity of the time evolution
  - 2 reality of the energy eigenvalues

# Introduction

- Hermiticity – Crucial? Guarantees,
  - 1 unitarity of the time evolution
  - 2 reality of the energy eigenvalues
- [Ben05] showed that hermiticity not necessary!

# Introduction

- Hermiticity – Crucial? Guarantees,
  - 1 unitarity of the time evolution
  - 2 reality of the energy eigenvalues
- [Ben05] showed that hermiticity not necessary!  $\mathcal{PT}$  Symmetry!

# Introduction

- Hermiticity – Crucial? Guarantees,
  - 1 unitarity of the time evolution
  - 2 reality of the energy eigenvalues
- [Ben05] showed that hermiticity not necessary!  $\mathcal{PT}$  Symmetry!
- Use?



# Introduction

- Hermiticity – Crucial? Guarantees,
  - 1 unitarity of the time evolution
  - 2 reality of the energy eigenvalues
- [Ben05] showed that hermiticity not necessary!  $\mathcal{PT}$  Symmetry!
- Use? Optic systems with gain and loss [LJS14]

# Introduction

- Hermiticity – Crucial? Guarantees,
  - 1 unitarity of the time evolution
  - 2 reality of the energy eigenvalues
- [Ben05] showed that hermiticity not necessary!  $\mathcal{PT}$  Symmetry!
- Use? Optic systems with gain and loss [LJS14], in open systems [Rot09]

# Introduction

- Hermiticity – Crucial? Guarantees,
  - 1 unitarity of the time evolution
  - 2 reality of the energy eigenvalues
- [Ben05] showed that hermiticity not necessary!  $\mathcal{PT}$  Symmetry!
- Use? Optic systems with gain and loss [LJS14], in open systems [Rot09][CW15]

# Introduction

- Hermiticity – Crucial? Guarantees,
  - 1 unitarity of the time evolution
  - 2 reality of the energy eigenvalues
- [Ben05] showed that hermiticity not necessary!  $\mathcal{PT}$  Symmetry!
- Use? Optic systems with gain and loss [LJS14], in open systems [Rot09][CW15] and in solid state systems [SF18].

# Introduction

- Hermiticity – Crucial? Guarantees,
  - 1 unitarity of the time evolution
  - 2 reality of the energy eigenvalues
- [Ben05] showed that hermiticity not necessary!  $\mathcal{PT}$  Symmetry!
- Use? Optic systems with gain and loss [LJS14], in open systems [Rot09][CW15] and in solid state systems [SF18].
- Surprising effects in store!

# Introduction

- Hermiticity – Crucial? Guarantees,
  - 1 unitarity of the time evolution
  - 2 reality of the energy eigenvalues
- [Ben05] showed that hermiticity not necessary!  $\mathcal{PT}$  Symmetry!
- Use? Optic systems with gain and loss [LJS14], in open systems [Rot09][CW15] and in solid state systems [SF18].
- Surprising effects in store!
  - 1 The Bulk Edge Correspondence does **not work** for NH systems!

# Introduction

- Hermiticity – Crucial? Guarantees,
  - 1 unitarity of the time evolution
  - 2 reality of the energy eigenvalues
- [Ben05] showed that hermiticity not necessary!  $\mathcal{PT}$  Symmetry!
- Use? Optic systems with gain and loss [LJS14], in open systems [Rot09][CW15] and in solid state systems [SF18].
- Surprising effects in store!
  - 1 The Bulk Edge Correspondence does **not work** for NH systems!
  - 2 NHSE: Non Hermitian Skin Effect

# Introduction

- Hermiticity – Crucial? Guarantees,
  - 1 unitarity of the time evolution
  - 2 reality of the energy eigenvalues
- [Ben05] showed that hermiticity not necessary!  $\mathcal{PT}$  Symmetry!
- Use? Optic systems with gain and loss [LJS14], in open systems [Rot09][CW15] and in solid state systems [SF18].
- Surprising effects in store!
  - 1 The Bulk Edge Correspondence does **not work** for NH systems!
  - 2 NHSE: Non Hermitian Skin Effect – *many* bulk states decay exponentially into the bulk!



# Introduction

- Hermiticity – Crucial? Guarantees,
  - ① unitarity of the time evolution
  - ② reality of the energy eigenvalues
- [Ben05] showed that hermiticity not necessary!  $\mathcal{PT}$  Symmetry!
- Use? Optic systems with gain and loss [LJS14], in open systems [Rot09][CW15] and in solid state systems [SF18].
- Surprising effects in store!
  - ① The Bulk Edge Correspondence does **not work** for NH systems!
  - ② NHSE: Non Hermitian Skin Effect – *many* bulk states decay exponentially into the bulk!
- Why? Topological Origins? Let's see!

# Why study NH systems?

# Why study NH systems?

- Physical Requirement: *Something in the way?*

# Why study NH systems?

- Physical Requirement: *Something in the way?*
- Need some forbidden points in order to have non trivial deformations.

# Why study NH systems?

- Physical Requirement: *Something in the way?*
- Need some forbidden points in order to have non trivial deformations.
- Simplest assumption: There exists a point gap.

# Why study NH systems?

- Physical Requirement: *Something in the way?*
- Need some forbidden points in order to have non trivial deformations.
- Simplest assumption: There exists a point gap. That is,

# Why study NH systems?

- Physical Requirement: *Something in the way?*
- Need some forbidden points in order to have non trivial deformations.
- Simplest assumption: There exists a point gap. That is,

$$\det\{H(k)\} \neq 0 \forall k \quad (1)$$

# Why study NH systems?

- Physical Requirement: *Something in the way?*
- Need some forbidden points in order to have non trivial deformations.
- Simplest assumption: There exists a point gap. That is,

$$\det\{H(k)\} \neq 0 \forall k \quad (1)$$

- Now?



# Why study NH systems?

- Physical Requirement: *Something in the way?*
- Need some forbidden points in order to have non trivial deformations.
- Simplest assumption: There exists a point gap. That is,

$$\det\{H(k)\} \neq 0 \forall k \quad (1)$$

- Now? Winding number,

# Why study NH systems?

- Physical Requirement: *Something in the way?*
- Need some forbidden points in order to have non trivial deformations.
- Simplest assumption: There exists a point gap. That is,

$$\det\{H(k)\} \neq 0 \forall k \quad (1)$$

- Now? Winding number,

$$w \equiv \int_{-\pi}^{\pi} \frac{dk}{2\pi i} \partial_k \ln \det H(k) \quad (2)$$

# Why study NH systems?

- Physical Requirement: *Something in the way?*
- Need some forbidden points in order to have non trivial deformations.
- Simplest assumption: There exists a point gap. That is,

$$\det\{H(k)\} \neq 0 \forall k \quad (1)$$

- Now? Winding number,

$$w \equiv \int_{-\pi}^{\pi} \frac{dk}{2\pi i} \partial_k \ln \det H(k) \quad (2)$$

$$w = \sum_{n=1}^N \int_{-\pi}^{\pi} \frac{dk}{2\pi} \partial_k \arg E_n(k) \quad (3)$$

# Hatano Nelson Model: A Surprise [Gon+18]

- Minimal one band model

# Hatano Nelson Model: A Surprise [Gon+18]

- Minimal one band model

$$H = \sum_j \left( J_R c_{j+1}^\dagger c_j + J_L c_j^\dagger c_{j+1} \right) \quad (4)$$

# Hatano Nelson Model: A Surprise [Gon+18]

- Minimal one band model

$$H = \sum_j \left( J_R c_{j+1}^\dagger c_j + J_L c_j^\dagger c_{j+1} \right) \quad (4)$$

- Take FT,

# Hatano Nelson Model: A Surprise [Gon+18]

- Minimal one band model

$$H = \sum_j \left( J_R c_{j+1}^\dagger c_j + J_L c_j^\dagger c_{j+1} \right) \quad (4)$$

- Take FT,

$$H(k) = J_R e^{-ik} + J_L e^{ik} \quad (5)$$

# Hatano Nelson Model: A Surprise [Gon+18]

- Minimal one band model

$$H = \sum_j \left( J_R c_{j+1}^\dagger c_j + J_L c_j^\dagger c_{j+1} \right) \quad (4)$$

- Take FT,

$$H(k) = J_R e^{-ik} + J_L e^{ik} \quad (5)$$

Computing Winding number



# Hatano Nelson Model: A Surprise [Gon+18]

- Minimal one band model

$$H = \sum_j \left( J_R c_{j+1}^\dagger c_j + J_L c_j^\dagger c_{j+1} \right) \quad (4)$$

- Take FT,

$$H(k) = J_R e^{-ik} + J_L e^{ik} \quad (5)$$

Computing Winding number

$$w = \begin{cases} 1 & |J_R| < |J_L|; \\ -1 & |J_R| > |J_L|. \end{cases} \quad (6)$$

# Hatano Nelson Model: A Surprise [Gon+18]




Figure 1: Hatano Nelson Model: Single Band Topology!

# Biorthogonal Setup [Bro14], [SAC06]

Given a NH system, we have a hamiltonian  $H$  such that  $H \neq H^\dagger$ .

# Biorthogonal Setup [Bro14], [SAC06]

Given a NH system, we have a hamiltonian  $H$  such that  $H \neq H^\dagger$ . There exist left eigenkets  $\{|\psi_R^n\rangle : n\}$  (not orthogonal) and right eigenkets  $\{|\psi_L^n\rangle : n\}$

# Biorthogonal Setup [Bro14], [SAC06]

Given a NH system, we have a hamiltonian  $H$  such that  $H \neq H^\dagger$ . There exist left eigenkets  $\{|\psi_R^n\rangle : n\}$  (not orthogonal) and right eigenkets  $\{|\psi_L^n\rangle : n\}$  such that,

# Biorthogonal Setup [Bro14], [SAC06]

Given a NH system, we have a hamiltonian  $H$  such that  $H \neq H^\dagger$ . There exist left eigenkets  $\{|\psi_R^n\rangle : n\}$  (not orthogonal) and right eigenkets  $\{|\psi_L^n\rangle : n\}$  such that,

- $H|\psi_R^n\rangle = E_n|\psi_R^n\rangle$

# Biorthogonal Setup [Bro14], [SAC06]

Given a NH system, we have a hamiltonian  $H$  such that  $H \neq H^\dagger$ . There exist left eigenkets  $\{|\psi_R^n\rangle : n\}$  (not orthogonal) and right eigenkets  $\{|\psi_L^n\rangle : n\}$  such that,

- $H |\psi_R^n\rangle = E_n |\psi_R^n\rangle$
- $H^\dagger |\psi_L^n\rangle = E_n^* |\psi_L^n\rangle$

# Biorthogonal Setup [Bro14], [SAC06]

Given a NH system, we have a hamiltonian  $H$  such that  $H \neq H^\dagger$ . There exist left eigenkets  $\{|\psi_R^n\rangle : n\}$  (not orthogonal) and right eigenkets  $\{|\psi_L^n\rangle : n\}$  such that,

- $H |\psi_R^n\rangle = E_n |\psi_R^n\rangle$
- $H^\dagger |\psi_L^n\rangle = E_n^* |\psi_L^n\rangle$
- $\langle \psi_L^n | \psi_R^m \rangle = \delta_{mn}$



## Biorthogonal Setup [Bro14], [SAC06]

Given a NH system, we have a hamiltonian  $H$  such that  $H \neq H^\dagger$ . There exist left eigenkets  $\{|\psi_R^n\rangle : n\}$  (not orthogonal) and right eigenkets  $\{|\psi_L^n\rangle : n\}$  such that,

- $H |\psi_R^n\rangle = E_n |\psi_R^n\rangle$
- $H^\dagger |\psi_L^n\rangle = E_n^* |\psi_L^n\rangle$
- $\langle \psi_L^n | \psi_R^m \rangle = \delta_{mn}$
- $\sum_n |\psi_L^n\rangle \langle \psi_R^n| = 1$

(Assuming no degeneracies for simplicity)

# Biorthogonal Setup [Bro14]

- Let  $|\psi\rangle \in \mathcal{H}$ ,

# Biorthogonal Setup [Bro14]

- Let  $|\psi\rangle \in \mathcal{H}$ ,

$$|\psi\rangle = \sum_n c_n |\psi_R^n\rangle \quad (7)$$

$$|\psi\rangle = \sum_n d_n |\psi_L^n\rangle \quad (8)$$

# Biorthogonal Setup [Bro14]

- Let  $|\psi\rangle \in \mathcal{H}$ ,

$$|\psi\rangle = \sum_n c_n |\psi_R^n\rangle \quad (7)$$

$$|\psi\rangle = \sum_n d_n |\psi_L^n\rangle \quad (8)$$

- The work in [ZW19] guaranteed the existence of a hermitian matrix  $X$  such that

# Biorthogonal Setup [Bro14]

- Let  $|\psi\rangle \in \mathcal{H}$ ,

$$|\psi\rangle = \sum_n c_n |\psi_R^n\rangle \quad (7)$$

$$|\psi\rangle = \sum_n d_n |\psi_L^n\rangle \quad (8)$$

- The work in [ZW19] guaranteed the existence of a hermitian matrix  $X$  such that

$$|\psi_L^n\rangle = X |\psi_R^n\rangle \text{ for all } n \quad (9)$$

# Biorthogonal Setup [Bro14]

- Let  $|\psi\rangle \in \mathcal{H}$ ,

$$|\psi\rangle = \sum_n c_n |\psi_R^n\rangle \quad (7)$$

$$|\psi\rangle = \sum_n d_n |\psi_L^n\rangle \quad (8)$$

- The work in [ZW19] guaranteed the existence of a hermitian matrix  $X$  such that

$$|\psi_L^n\rangle = X |\psi_R^n\rangle \text{ for all } n \quad (9)$$

- $\langle\psi, \psi\rangle_X \equiv \langle\psi | X | \psi\rangle = \sum_n c_n d_n^*$

# Biorthogonal Setup [Bro14]

- Let  $|\psi\rangle \in \mathcal{H}$ ,

$$|\psi\rangle = \sum_n c_n |\psi_R^n\rangle \quad (7)$$

$$|\psi\rangle = \sum_n d_n |\psi_L^n\rangle \quad (8)$$

- The work in [ZW19] guaranteed the existence of a hermitian matrix  $X$  such that

$$|\psi_L^n\rangle = X |\psi_R^n\rangle \text{ for all } n \quad (9)$$

- $\langle\psi, \psi\rangle_X \equiv \langle\psi | X | \psi\rangle = \sum_n c_n d_n^*$

# Exceptional Points (EPs) [BBK21]

- Interesting property of NH systems: *Exceptional Points*



# Exceptional Points (EPs) [BBK21]

- Interesting property of NH systems: *Exceptional Points*
- Hamiltonian matrix is defective  $\implies GM < AM$

# Exceptional Points (EPs) [BBK21]

- Interesting property of NH systems: *Exceptional Points*
- Hamiltonian matrix is defective  $\implies GM < AM$
- Illustrative example

# Exceptional Points (EPs) [BBK21]

- Interesting property of NH systems: *Exceptional Points*
- Hamiltonian matrix is defective  $\implies GM < AM$
- Illustrative example

$$H(\alpha) = \begin{pmatrix} 0 & \alpha \\ 1 & 0 \end{pmatrix} \quad \alpha \in \mathbb{C} \quad (10)$$

## Exceptional Points (EPs) [BBK21]

- Interesting property of NH systems: *Exceptional Points*
- Hamiltonian matrix is defective  $\implies GM < AM$
- Illustrative example

$$H(\alpha) = \begin{pmatrix} 0 & \alpha \\ 1 & 0 \end{pmatrix} \quad \alpha \in \mathbb{C} \quad (10)$$

$$E_{\pm} = \pm\sqrt{\alpha} \quad |\psi_R^{\pm}\rangle = \begin{pmatrix} \pm\sqrt{\alpha} \\ 1 \end{pmatrix} \quad |\psi_L^{\pm}\rangle = \begin{pmatrix} 1 \\ \pm\sqrt{\alpha} \end{pmatrix} \quad (11)$$

## Exceptional Points (EPs) [BBK21]

- Interesting property of NH systems: *Exceptional Points*
- Hamiltonian matrix is defective  $\implies GM < AM$
- Illustrative example

$$H(\alpha) = \begin{pmatrix} 0 & \alpha \\ 1 & 0 \end{pmatrix} \quad \alpha \in \mathbb{C} \quad (10)$$

$$E_{\pm} = \pm\sqrt{\alpha} \quad |\psi_R^{\pm}\rangle = \begin{pmatrix} \pm\sqrt{\alpha} \\ 1 \end{pmatrix} \quad |\psi_L^{\pm}\rangle = \begin{pmatrix} 1 \\ \pm\sqrt{\alpha} \end{pmatrix} \quad (11)$$

- Eigenvectors *coalesce* at these points.

# Exceptional Points (EPs) [MA19]




Figure 2: Exceptional Points Visualized

# EPs $\rightarrow$ Monopoles and Mobius Strips! [ZW20]

Hermitian Mobius Strips – need to ‘contact’ degeneracies.

# EPs $\rightarrow$ Monopoles and Mobius Strips! [ZW20]

Hermitian Mobius Strips – need to ‘contact’ degeneracies.




Figure 3: Hermitian Mobius Strip



# EPs $\rightarrow$ Monopoles and Mobius Strips! [ZW20]

Non Hermitian Mobius Strips – need EPs? Or do we?

# EPs $\rightarrow$ Monopoles and Mobius Strips! [ZW20]

Non Hermitian Mobius Strips – need EPs? Or do we?




Figure 4: NH Mobius Strip

# EPs $\rightarrow$ Monopoles and Mobius Strips! [ZW20]

Non Hermitian Mobius Strips – need EPs? Or do we?




Figure 4: NH Mobius Strip

Branch Cuts!  $f(z) := \sqrt{z}$  is not holomorphic on  $\mathbb{C}$ .

# Geometric Phase in NH Systems [ZW19]

- Usual setting  $H \equiv H(\mathbf{R}(t))$

# Geometric Phase in NH Systems [ZW19]

- Usual setting  $H \equiv H(\mathbf{R}(t))$
- Ansatz

# Geometric Phase in NH Systems [ZW19]

- Usual setting  $H \equiv H(\mathbf{R}(t))$
- Ansatz

$$|\psi(t)\rangle = \sum_n c_n(t) \exp\left\{\frac{-i}{\hbar} \int_0^t E_n(t') dt'\right\} |\psi_R^n(t)\rangle \quad (12)$$

# Geometric Phase in NH Systems [ZW19]

- Usual setting  $H \equiv H(\mathbf{R}(t))$
- Ansatz

$$|\psi(t)\rangle = \sum_n c_n(t) \exp\left\{\frac{-i}{\hbar} \int_0^t E_n(t') dt'\right\} |\psi_R^n(t)\rangle \quad (12)$$

- Using the TDSE,  $i\hbar\partial_t |\psi\rangle = H|\psi\rangle$ ,

# Geometric Phase in NH Systems [ZW19]

- Usual setting  $H \equiv H(\mathbf{R}(t))$
- Ansatz

$$|\psi(t)\rangle = \sum_n c_n(t) \exp\left\{\frac{-\iota}{\hbar} \int_0^t E_n(t') dt'\right\} |\psi_R^n(t)\rangle \quad (12)$$

- Using the TDSE,  $\iota\hbar\partial_t|\psi\rangle = H|\psi\rangle$ ,

$$\iota\hbar \sum_n \dot{c}_n(t) \exp\left\{\frac{-\iota}{\hbar} \int_0^t E_n(t') dt'\right\} |\psi_R^n(t)\rangle + \iota\hbar \sum_n c_n(t) \exp\left\{\frac{-\iota}{\hbar} \int_0^t E_n(t') dt'\right\} |\dot{\psi}_R^n(t)\rangle = 0 \quad (13)$$



# Geometric Phase in NH Systems [ZW19]

- Usual setting  $H \equiv H(\mathbf{R}(t))$
- Ansatz

$$|\psi(t)\rangle = \sum_n c_n(t) \exp\left\{\frac{-\iota}{\hbar} \int_0^t E_n(t') dt'\right\} |\psi_R^n(t)\rangle \quad (12)$$

- Using the TDSE,  $\iota\hbar\partial_t|\psi\rangle = H|\psi\rangle$ ,

$$\iota\hbar \sum_n \dot{c}_n(t) \exp\left\{\frac{-\iota}{\hbar} \int_0^t E_n(t') dt'\right\} |\psi_R^n(t)\rangle + \iota\hbar \sum_n c_n(t) \exp\left\{\frac{-\iota}{\hbar} \int_0^t E_n(t') dt'\right\} |\dot{\psi}_R^n(t)\rangle = 0 \quad (13)$$

- Further,

# Geometric Phase in NH Systems [ZW19]

- Usual setting  $H \equiv H(\mathbf{R}(t))$
- Ansatz

$$|\psi(t)\rangle = \sum_n c_n(t) \exp\left\{\frac{-\iota}{\hbar} \int_0^t E_n(t') dt'\right\} |\psi_R^n(t)\rangle \quad (12)$$

- Using the TDSE,  $\iota\hbar\partial_t|\psi\rangle = H|\psi\rangle$ ,

$$\iota\hbar \sum_n \dot{c}_n(t) \exp\left\{\frac{-\iota}{\hbar} \int_0^t E_n(t') dt'\right\} |\psi_R^n(t)\rangle + \iota\hbar \sum_n c_n(t) \exp\left\{\frac{-\iota}{\hbar} \int_0^t E_n(t') dt'\right\} |\dot{\psi}_R^n(t)\rangle = 0 \quad (13)$$

- Further,

$$\dot{c}_m(t) = -c_m(t) \langle \psi_L^m | \dot{\psi}_R^m \rangle - \sum_{n \neq m} c_n(t) \langle \psi_L^m | \dot{\psi}_R^n \rangle \exp\left\{\iota \int_0^t \omega_{mn}(t') dt'\right\} \quad (14)$$

# Geometric Phase in NH Systems [ZW19]

- Usual setting  $H \equiv H(\mathbf{R}(t))$
- Ansatz

$$|\psi(t)\rangle = \sum_n c_n(t) \exp\left\{\frac{-\iota}{\hbar} \int_0^t E_n(t') dt'\right\} |\psi_R^n(t)\rangle \quad (12)$$

- Using the TDSE,  $\iota\hbar\partial_t|\psi\rangle = H|\psi\rangle$ ,

$$\iota\hbar \sum_n \dot{c}_n(t) \exp\left\{\frac{-\iota}{\hbar} \int_0^t E_n(t') dt'\right\} |\psi_R^n(t)\rangle + \iota\hbar \sum_n c_n(t) \exp\left\{\frac{-\iota}{\hbar} \int_0^t E_n(t') dt'\right\} |\dot{\psi}_R^n(t)\rangle = 0 \quad (13)$$

- Further,

$$\dot{c}_m(t) = -c_m(t) \langle \psi_L^m | \dot{\psi}_R^m \rangle - \sum_{n \neq m} c_n(t) \langle \psi_L^m | \dot{\psi}_R^n \rangle \exp\left\{\iota \int_0^t \omega_{mn}(t') dt'\right\} \quad (14)$$

Where,  $\omega_{mn}(t) = (E_m(t) - E_n(t))/\hbar$ .

# Geometric Phase in NH Systems

- Need real eigenvalues!

# Geometric Phase in NH Systems

- Need real eigenvalues!
- 

$$\dot{c}_m = -c_m \langle \psi_L^m | \dot{\psi}_R^m \rangle \quad (15)$$

# Geometric Phase in NH Systems

- Need real eigenvalues!
- 

$$\dot{c}_m = -c_m \langle \psi_L^m | \dot{\psi}_R^m \rangle \quad (15)$$

- This gives us,

# Geometric Phase in NH Systems

- Need real eigenvalues!
- 

$$\dot{c}_m = -c_m \langle \psi_L^m | \dot{\psi}_R^m \rangle \quad (15)$$

- This gives us,

$$c_m(t) = \exp \left\{ i \left( \int i \langle \psi_L^m(\mathbf{R}) | \nabla_{\mathbf{R}} \psi_R^m(\mathbf{R}) \rangle \cdot d\mathbf{R} \right) \right\} \quad (16)$$

# Geometric Phase in NH Systems

- Need real eigenvalues!

- 

$$\dot{c}_m = -c_m \langle \psi_L^m | \dot{\psi}_R^m \rangle \quad (15)$$

- This gives us,

$$c_m(t) = \exp \left\{ \imath \left( \int \imath \langle \psi_L^m(\mathbf{R}) | \nabla_{\mathbf{R}} \psi_R^m(\mathbf{R}) \rangle \cdot d\mathbf{R} \right) \right\} \quad (16)$$

- 

$$\boxed{\mathbf{A}_m = \imath \langle \psi_L^m(\mathbf{R}) | \nabla_{\mathbf{R}} \psi_R^m(\mathbf{R}) \rangle} \quad (17)$$



# Geometric Phase in NH Systems

- Need real eigenvalues!



$$\dot{c}_m = -c_m \langle \psi_L^m | \dot{\psi}_R^m \rangle \quad (15)$$

- This gives us,

$$c_m(t) = \exp \left\{ \iota \left( \int \iota \langle \psi_L^m(\mathbf{R}) | \nabla_{\mathbf{R}} \psi_R^m(\mathbf{R}) \rangle \cdot d\mathbf{R} \right) \right\} \quad (16)$$



$$\mathbf{A}_m = \iota \langle \psi_L^m(\mathbf{R}) | \nabla_{\mathbf{R}} \psi_R^m(\mathbf{R}) \rangle \quad (17)$$



$$\mathbf{B}_m = \iota \langle \nabla \psi_L^m | \times | \nabla \psi_R^m \rangle \quad (18)$$

# Geometric Phase in NH Systems

- Need real eigenvalues!

- 

$$\dot{c}_m = -c_m \langle \psi_L^m | \dot{\psi}_R^m \rangle \quad (15)$$

- This gives us,

$$c_m(t) = \exp \left\{ \iota \left( \int \iota \langle \psi_L^m(\mathbf{R}) | \nabla_{\mathbf{R}} \psi_R^m(\mathbf{R}) \rangle \cdot d\mathbf{R} \right) \right\} \quad (16)$$

- 

$$\mathbf{A}_m = \iota \langle \psi_L^m(\mathbf{R}) | \nabla_{\mathbf{R}} \psi_R^m(\mathbf{R}) \rangle \quad (17)$$

- 

$$\mathbf{B}_m = \iota \langle \nabla \psi_L^m | \times | \nabla \psi_R^m \rangle \quad (18)$$

- Berry phase is real only if ([ZW19])

# Geometric Phase in NH Systems

- Need real eigenvalues!

- 

$$\dot{c}_m = -c_m \langle \psi_L^m | \dot{\psi}_R^m \rangle \quad (15)$$

- This gives us,

$$c_m(t) = \exp \left\{ \iota \left( \int \iota \langle \psi_L^m(\mathbf{R}) | \nabla_{\mathbf{R}} \psi_R^m(\mathbf{R}) \rangle \cdot d\mathbf{R} \right) \right\} \quad (16)$$

- 

$$\mathbf{A}_m = \iota \langle \psi_L^m(\mathbf{R}) | \nabla_{\mathbf{R}} \psi_R^m(\mathbf{R}) \rangle \quad (17)$$

- 

$$\mathbf{B}_m = \iota \langle \nabla \psi_L^m | \times | \nabla \psi_R^m \rangle \quad (18)$$

- Berry phase is real only if ([ZW19])

$$\langle \psi_R^m | (\nabla_{\mathbf{R}} X) | \psi_R^m \rangle = 0 \quad (19)$$

# Breakdown of the BBC




Figure 5: SSH Chain: Standard Energy Plot

# Breakdown of the BBC




Figure 6: SSH Chain: Non-reciprocity with  $\Delta = 0.2$

# NHSE!




Figure 7: SSH Chain: Standard Density Plot

# NHSE!




Figure 8: SSH Chain: Non-reciprocity with  $\Delta = 0.2$

# NH-BBC [ZYF20] [Gon+18]

- Perform the generalization



## NH-BBC [ZYF20] [Gon+18]

- Perform the generalization

$$\exp\{\iota k\} \rightarrow \beta \in \mathbb{C} \quad (20)$$

## NH-BBC [ZYF20] [Gon+18]

- Perform the generalization

$$\exp\{\imath k\} \rightarrow \beta \in \mathbb{C} \quad (20)$$

- Consider the *one* band model,

## NH-BBC [ZYF20] [Gon+18]

- Perform the generalization

$$\exp\{\imath k\} \rightarrow \beta \in \mathbb{C} \quad (20)$$

- Consider the *one* band model,

$$H = \sum_{i,j} t_{i,j} |i\rangle \langle j| \rightarrow \sum_{k \in BZ} H(k) |k\rangle \langle k| \quad (21)$$

## NH-BBC [ZYF20] [Gon+18]

- Perform the generalization

$$\exp\{\imath k\} \rightarrow \beta \in \mathbb{C} \quad (20)$$

- Consider the *one* band model,

$$H = \sum_{i,j} t_{i,j} |i\rangle \langle j| \rightarrow \sum_{k \in BZ} H(k) |k\rangle \langle k| \quad (21)$$

- Let  $t_{i,j} \equiv t_{i-j}$  be non-zero only for  $-m \leq i-j \leq n$ .

# NH-BBC [ZYF20] [Gon+18]

- Perform the generalization

$$\exp\{\imath k\} \rightarrow \beta \in \mathbb{C} \quad (20)$$

- Consider the *one* band model,

$$H = \sum_{i,j} t_{i,j} |i\rangle \langle j| \rightarrow \sum_{k \in BZ} H(k) |k\rangle \langle k| \quad (21)$$

- Let  $t_{i,j} \equiv t_{i-j}$  be non-zero only for  $-m \leq i-j \leq n$ .

$$H(k) = \sum_r t_r (\exp\{\imath k\})^r \implies H(\beta) = t_{-m} \beta^{-m} + \cdots + t_n \beta^n = \frac{P(\beta)}{\beta^m} \quad (22)$$

# NH-BBC [ZYF20] [Gon+18]

- Perform the generalization

$$\exp\{\imath k\} \rightarrow \beta \in \mathbb{C} \quad (20)$$

- Consider the *one* band model,

$$H = \sum_{i,j} t_{i,j} |i\rangle \langle j| \rightarrow \sum_{k \in BZ} H(k) |k\rangle \langle k| \quad (21)$$

- Let  $t_{i,j} \equiv t_{i-j}$  be non-zero only for  $-m \leq i-j \leq n$ .

$$H(k) = \sum_r t_r (\exp\{\imath k\})^r \implies H(\beta) = t_{-m} \beta^{-m} + \cdots + t_n \beta^n = \frac{P(\beta)}{\beta^m} \quad (22)$$

- Winding number

# NH-BBC [ZYF20] [Gon+18]

- Perform the generalization

$$\exp\{\imath k\} \rightarrow \beta \in \mathbb{C} \quad (20)$$

- Consider the *one* band model,

$$H = \sum_{i,j} t_{i,j} |i\rangle \langle j| \rightarrow \sum_{k \in BZ} H(k) |k\rangle \langle k| \quad (21)$$

- Let  $t_{i,j} \equiv t_{i-j}$  be non-zero only for  $-m \leq i-j \leq n$ .

$$H(k) = \sum_r t_r (\exp\{\imath k\})^r \implies H(\beta) = t_{-m} \beta^{-m} + \cdots + t_n \beta^n = \frac{P(\beta)}{\beta^m} \quad (22)$$

- Winding number

$$w(E_b) = \frac{1}{2\pi} \oint_C \frac{d}{dz} \arg[H(z) - E_b] dz \quad (23)$$

# NH-BBC [ZYF20] [Gon+18]

- Perform the generalization

$$\exp\{\imath k\} \rightarrow \beta \in \mathbb{C} \quad (20)$$

- Consider the *one* band model,

$$H = \sum_{i,j} t_{i,j} |i\rangle \langle j| \rightarrow \sum_{k \in BZ} H(k) |k\rangle \langle k| \quad (21)$$

- Let  $t_{i,j} \equiv t_{i-j}$  be non-zero only for  $-m \leq i-j \leq n$ .

$$H(k) = \sum_r t_r (\exp\{\imath k\})^r \implies H(\beta) = t_{-m} \beta^{-m} + \cdots + t_n \beta^n = \frac{P(\beta)}{\beta^m} \quad (22)$$

- Winding number

$$w(E_b) = \frac{1}{2\pi} \oint_C \frac{d}{dz} \arg[H(z) - E_b] dz \quad (23)$$

Now what?



# Generalized Brillouin Zone [ZYF20] [Gon+18] [YM19]

Two Steps

# Generalized Brillouin Zone [ZYF20] [Gon+18] [YM19]

Two Steps

- 1 Solve the algebraic equation

# Generalized Brillouin Zone [ZYF20] [Gon+18] [YM19]

Two Steps

- 1 Solve the algebraic equation

$$\det(H(\beta) - E) = 0$$

# Generalized Brillouin Zone [ZYF20] [Gon+18] [YM19]

## Two Steps

- 1 Solve the algebraic equation

$$\det(H(\beta) - E) = 0$$

- 2 Order the solutions, such that  $|\beta_i| \leq |\beta_{i+1}|$ .

# Generalized Brillouin Zone [ZYF20] [Gon+18] [YM19]

## Two Steps

- 1 Solve the algebraic equation

$$\det(H(\beta) - E) = 0$$

- 2 Order the solutions, such that  $|\beta_i| \leq |\beta_{i+1}|$ .

### Definition 1 (GBZ)

For the hamiltonian  $H(\beta)$  with solutions to the determinant equation  $|\beta_1| \leq |\beta_2| \leq \dots$  define the GBZ as the following set in the complex plane,

# Generalized Brillouin Zone [ZYF20] [Gon+18] [YM19]

## Two Steps

- 1 Solve the algebraic equation

$$\det(H(\beta) - E) = 0$$

- 2 Order the solutions, such that  $|\beta_i| \leq |\beta_{i+1}|$ .

### Definition 1 (GBZ)

For the hamiltonian  $H(\beta)$  with solutions to the determinant equation  $|\beta_1| \leq |\beta_2| \leq \dots$  define the GBZ as the following set in the complex plane,

$$GBZ := \{\beta \mid |\beta_m(H(\beta))| = |\beta_{m+1}(H(\beta))|\} \quad (24)$$

# Generalized Brillouin Zone [ZYF20] [Gon+18] [YM19]

## Two Steps

- 1 Solve the algebraic equation

$$\det(H(\beta) - E) = 0$$

- 2 Order the solutions, such that  $|\beta_i| \leq |\beta_{i+1}|$ .

## Definition 1 (GBZ)

For the hamiltonian  $H(\beta)$  with solutions to the determinant equation  $|\beta_1| \leq |\beta_2| \leq \dots$  define the GBZ as the following set in the complex plane,

$$GBZ := \{\beta \mid |\beta_m(H(\beta))| = |\beta_{m+1}(H(\beta))|\} \quad (24)$$

## Theorem 2 (GBZ Theorem)

*The GBZ is that closed contour in the complex plane which encircles the pole of order  $m$  at the origin and exactly  $m$  zeros of the polynomial  $P(z) - Ez^m$  for any  $E \in \mathbb{C}$ .*

# GBZ: Consequences [ZYF20] [Gon+18] [YM19]




Figure 9: [ZYF20] GBZ, BZ and spectra for different hamiltonians  $H(z) = z^{-2}/5 + 3z^{-1} + 2z$ ,  $H(z) = z^{-2}/5 + z^{-1} + 2z$  and  $H(z) = 2z^{-2}/5 + z^{-1} + z$



# GBZ: Consequences [ZYF20] [Gon+18] [YM19]

Three Consequences.

# GBZ: Consequences [ZYF20] [Gon+18] [YM19]

Three Consequences.

- 1 The OBC of an infinite chain is reflected upon by evaluation on the GBZ, **not** the BZ!

# GBZ: Consequences [ZYF20] [Gon+18] [YM19]

Three Consequences.

- 1 The OBC of an infinite chain is reflected upon by evaluation on the GBZ, **not** the BZ!
- 2 No winding of any complex energy in the mapped contour on the energy plane (by using Cauchy's Residue Principle).

# GBZ: Consequences [ZYF20] [Gon+18] [YM19]

Three Consequences.

- 1 The OBC of an infinite chain is reflected upon by evaluation on the GBZ, **not** the BZ!
- 2 No winding of any complex energy in the mapped contour on the energy plane (by using Cauchy's Residue Principle). The OBC spectrum collapses into a line, as seen in the figure.

# GBZ: Consequences [ZYF20] [Gon+18] [YM19]

Three Consequences.

- 1 The OBC of an infinite chain is reflected upon by evaluation on the GBZ, **not** the BZ!
- 2 No winding of any complex energy in the mapped contour on the energy plane (by using Cauchy's Residue Principle). The OBC spectrum collapses into a line, as seen in the figure.
- 3 Skin Modes: Claim in

# GBZ: Consequences [ZYF20] [Gon+18] [YM19]

Three Consequences.

- 1 The OBC of an infinite chain is reflected upon by evaluation on the GBZ, **not** the BZ!
- 2 No winding of any complex energy in the mapped contour on the energy plane (by using Cauchy's Residue Principle). The OBC spectrum collapses into a line, as seen in the figure.
- 3 Skin Modes: Claim in [ZYF20]

# GBZ: Consequences [ZYF20] [Gon+18] [YM19]

Three Consequences.

- 1 The OBC of an infinite chain is reflected upon by evaluation on the GBZ, **not** the BZ!
- 2 No winding of any complex energy in the mapped contour on the energy plane (by using Cauchy's Residue Principle). The OBC spectrum collapses into a line, as seen in the figure.
- 3 Skin Modes: Claim in[ZYF20]

$$\langle s | \psi(\beta) \rangle = \mathcal{O}(|\beta|^s) \quad (25)$$

# GBZ: Consequences [ZYF20] [Gon+18] [YM19]

Three Consequences.

- 1 The OBC of an infinite chain is reflected upon by evaluation on the GBZ, **not** the BZ!
- 2 No winding of any complex energy in the mapped contour on the energy plane (by using Cauchy's Residue Principle). The OBC spectrum collapses into a line, as seen in the figure.
- 3 Skin Modes: Claim in [ZYF20]

$$\langle s | \psi(\beta) \rangle = \mathcal{O}(|\beta|^s) \quad (25)$$

NHSE almost always!



# NH Polarization [Kun+18]

- Assume SSH with last unit cell broken.

# NH Polarization [Kun+18]

- Assume SSH with last unit cell broken.  $d_x \pm \imath d_y = f_{\pm} + g_{\pm} \exp\{\pm \imath ka\}$

# NH Polarization [Kun+18]

- Assume SSH with last unit cell broken.  $d_x \pm \imath d_y = f_{\pm} + g_{\pm} \exp\{\pm \imath ka\}$
- Exact states,

# NH Polarization [Kun+18]

- Assume SSH with last unit cell broken.  $d_x \pm \imath d_y = f_{\pm} + g_{\pm} \exp\{\pm \imath ka\}$
- Exact states,

$$\begin{aligned}
 |\psi_R\rangle &= \mathcal{N}_R \sum_{n=1}^N r_R^n c_{A,n}^\dagger |0\rangle \quad r_R = \frac{-f_+}{g_+} \\
 |\psi_K\rangle &= \mathcal{N}_L \sum_{n=1}^N r_L^n c_{A,n}^\dagger |0\rangle \quad r_L = \frac{-f_+^*}{g_+^*}
 \end{aligned} \tag{26}$$

# NH Polarization [Kun+18]

- Assume SSH with last unit cell broken.  $d_x \pm \imath d_y = f_{\pm} + g_{\pm} \exp\{\pm \imath ka\}$
- Exact states,

$$\begin{aligned} |\psi_R\rangle &= \mathcal{N}_R \sum_{n=1}^N r_R^n c_{A,n}^{\dagger} |0\rangle \quad r_R = \frac{-f_+}{g_+} \\ |\psi_K\rangle &= \mathcal{N}_L \sum_{n=1}^N r_L^n c_{A,n}^{\dagger} |0\rangle \quad r_L = \frac{-f_+^*}{g_+^*} \end{aligned} \quad (26)$$

$$\mathcal{N}_L^* \mathcal{N}_R = (r_L^* r_R)^{-1} (r_L^* r_R - 1) / [(r_L^* r_R)^N - 1] \quad (27)$$

# NH Polarization [Kun+18]

- Assume SSH with last unit cell broken.  $d_x \pm \imath d_y = f_{\pm} + g_{\pm} \exp\{\pm \imath ka\}$
- Exact states,

$$\begin{aligned} |\psi_R\rangle &= \mathcal{N}_R \sum_{n=1}^N r_R^n c_{A,n}^\dagger |0\rangle \quad r_R = \frac{-f_+}{g_+} \\ |\psi_K\rangle &= \mathcal{N}_L \sum_{n=1}^N r_L^n c_{A,n}^\dagger |0\rangle \quad r_L = \frac{-f_+^*}{g_+^*} \end{aligned} \quad (26)$$

- $$\mathcal{N}_L^* \mathcal{N}_R = (r_L^* r_R)^{-1} (r_L^* r_R - 1) / [(r_L^* r_R)^N - 1] \quad (27)$$

- Consider  $\Pi_n = |e_{A,n}\rangle \langle e_{A,n}| + |e_{B,n}\rangle \langle e_{B,n}|$ .

# NH Polarization [Kun+18]

- Assume SSH with last unit cell broken.  $d_x \pm \imath d_y = f_{\pm} + g_{\pm} \exp\{\pm \imath ka\}$
- Exact states,

$$\begin{aligned} |\psi_R\rangle &= \mathcal{N}_R \sum_{n=1}^N r_R^n c_{A,n}^\dagger |0\rangle \quad r_R = \frac{-f_+}{g_+} \\ |\psi_K\rangle &= \mathcal{N}_L \sum_{n=1}^N r_L^n c_{A,n}^\dagger |0\rangle \quad r_L = \frac{-f_+^*}{g_+^*} \end{aligned} \quad (26)$$

- $$\mathcal{N}_L^* \mathcal{N}_R = (r_L^* r_R)^{-1} (r_L^* r_R - 1) / [(r_L^* r_R)^N - 1] \quad (27)$$

- Consider  $\Pi_n = |e_{A,n}\rangle \langle e_{A,n}| + |e_{B,n}\rangle \langle e_{B,n}|$ .

$$\langle \Pi_n \rangle = \langle \psi_L | \Pi_n | \psi_R \rangle = \mathcal{N}_L^* \mathcal{N}_R (r_L^* r_R)^n \quad (28)$$

# NH Polarization [Kun+18]

- Assume SSH with last unit cell broken.  $d_x \pm \imath d_y = f_{\pm} + g_{\pm} \exp\{\pm \imath ka\}$
- Exact states,

$$\begin{aligned} |\psi_R\rangle &= \mathcal{N}_R \sum_{n=1}^N r_R^n c_{A,n}^\dagger |0\rangle \quad r_R = \frac{-f_+}{g_+} \\ |\psi_K\rangle &= \mathcal{N}_L \sum_{n=1}^N r_L^n c_{A,n}^\dagger |0\rangle \quad r_L = \frac{-f_+^*}{g_+^*} \end{aligned} \quad (26)$$

•

$$\mathcal{N}_L^* \mathcal{N}_R = (r_L^* r_R)^{-1} (r_L^* r_R - 1) / [(r_L^* r_R)^N - 1] \quad (27)$$

- Consider  $\Pi_n = |e_{A,n}\rangle \langle e_{A,n}| + |e_{B,n}\rangle \langle e_{B,n}|$ .

$$\langle \Pi_n \rangle = \langle \psi_L | \Pi_n | \psi_R \rangle = \mathcal{N}_L^* \mathcal{N}_R (r_L^* r_R)^n \quad (28)$$

- Critical point:  $|r_L^* r_R| = 1$ .



# NH Polarization [Kun+18]

- Assume SSH with last unit cell broken.  $d_x \pm \imath d_y = f_{\pm} + g_{\pm} \exp\{\pm \imath ka\}$
- Exact states,

$$\begin{aligned} |\psi_R\rangle &= \mathcal{N}_R \sum_{n=1}^N r_R^n c_{A,n}^\dagger |0\rangle \quad r_R = \frac{-f_+}{g_+} \\ |\psi_K\rangle &= \mathcal{N}_L \sum_{n=1}^N r_L^n c_{A,n}^\dagger |0\rangle \quad r_L = \frac{-f_+^*}{g_+^*} \end{aligned} \quad (26)$$

•

$$\mathcal{N}_L^* \mathcal{N}_R = (r_L^* r_R)^{-1} (r_L^* r_R - 1) / [(r_L^* r_R)^N - 1] \quad (27)$$

- Consider  $\Pi_n = |e_{A,n}\rangle \langle e_{A,n}| + |e_{B,n}\rangle \langle e_{B,n}|$ .

$$\langle \Pi_n \rangle = \langle \psi_L | \Pi_n | \psi_R \rangle = \mathcal{N}_L^* \mathcal{N}_R (r_L^* r_R)^n \quad (28)$$

- Critical point:  $|r_L^* r_R| = 1$ .

# NH Polarization [Kun+18]

## Biorthogonal polarization

# NH Polarization [Kun+18]

**Biorthogonal polarization**  $P$  is coined, such that,

# NH Polarization [Kun+18]


**Biorthogonal polarization**  $P$  is coined, such that,

$$P = 1 - \lim_{N \rightarrow \infty} (\langle \psi_L | \frac{\sum_n n \Pi_n}{N} | \psi_R \rangle) \quad (29)$$

# NH Polarization [Kun+18]

**Biorthogonal polarization**  $P$  is coined, such that,

$$P = 1 - \lim_{N \rightarrow \infty} (\langle \psi_L | \frac{\sum_n n \Pi_n}{N} | \psi_R \rangle) \quad (29)$$



**Figure 10:** Absolute spectra of NH SSH model. The gray lines indicate the periodic Bloch bands while the qualitatively different spectra in blue (bulk) and red (edge) correspond to the open system. The orange (dark green dashed) vertical lines indicate where  $r_L^* r_R = 1$  ( $r_L^* r_R = -1$ ) and the gray dotted-dashed lines correspond to the EPs of the periodic Bloch Hamiltonian. [Kun+18]

# Winding Number in NH SSH [Lie18]

$$\begin{aligned} H^{\text{hop}} = & v_1 \sum_{n=1}^N |n, B\rangle \langle n, A| + v_2 \sum_{n=1}^N |n, A\rangle \langle n, B| \\ & + w_1 \sum_{n=1}^{N-1} |n+1, A\rangle \langle n, B| + w_2 \sum_{n=1}^{N-1} |n, B\rangle \langle n+1, A| \end{aligned} \quad (30)$$

# Winding Number in NH SSH [Lie18]

$$\begin{aligned}
 H^{\text{hop}} = & v_1 \sum_{n=1}^N |n, B\rangle \langle n, A| + v_2 \sum_{n=1}^N |n, A\rangle \langle n, B| \\
 & + w_1 \sum_{n=1}^{N-1} |n+1, A\rangle \langle n, B| + w_2 \sum_{n=1}^{N-1} |n, B\rangle \langle n+1, A|
 \end{aligned} \tag{30}$$

$$H^{\text{pot}} = iu \sum_{n=1}^N (|n, A\rangle \langle n, A| - |n, B\rangle \langle n, B|). \tag{31}$$

# Winding Number in NH SSH [Lie18]

$$\begin{aligned}
 H^{\text{hop}} = & v_1 \sum_{n=1}^N |n, B\rangle \langle n, A| + v_2 \sum_{n=1}^N |n, A\rangle \langle n, B| \\
 & + w_1 \sum_{n=1}^{N-1} |n+1, A\rangle \langle n, B| + w_2 \sum_{n=1}^{N-1} |n, B\rangle \langle n+1, A|
 \end{aligned} \tag{30}$$

$$H^{\text{pot}} = iu \sum_{n=1}^N (|n, A\rangle \langle n, A| - |n, B\rangle \langle n, B|). \tag{31}$$

**Invariant**



# Winding Number in NH SSH [Lie18]

$$\begin{aligned}
 H^{\text{hop}} = & v_1 \sum_{n=1}^N |n, B\rangle \langle n, A| + v_2 \sum_{n=1}^N |n, A\rangle \langle n, B| \\
 & + w_1 \sum_{n=1}^{N-1} |n+1, A\rangle \langle n, B| + w_2 \sum_{n=1}^{N-1} |n, B\rangle \langle n+1, A|
 \end{aligned} \tag{30}$$

$$H^{\text{pot}} = iu \sum_{n=1}^N (|n, A\rangle \langle n, A| - |n, B\rangle \langle n, B|). \tag{31}$$

**Invariant**

$$Q_n^c = i \int_{BZ} \left\langle \psi_{L,k}^n \left| \frac{\partial}{\partial k} \right| \psi_{R,k}^n \right\rangle dk \tag{32}$$

# Winding Number in NH SSH: $u = 0$ [Lie18]

- $u = 0 \implies$  CS!

# Winding Number in NH SSH: $u = 0$ [Lie18]

- $u = 0 \implies$  CS!
- 

$$\sigma_z H(k) \sigma_z = -H(k) \quad (33)$$

# Winding Number in NH SSH: $u = 0$ [Lie18]

- $u = 0 \implies$  CS!
- 

$$\sigma_z H(k) \sigma_z = -H(k) \quad (33)$$

$$|\psi_R^\pm\rangle = \sigma_z |\psi_R^\mp\rangle \quad (34)$$

# Winding Number in NH SSH: $u = 0$ [Lie18]

- $u = 0 \implies$  CS!

- 

$$\sigma_z H(k) \sigma_z = -H(k) \quad (33)$$

$$|\psi_R^\pm\rangle = \sigma_z |\psi_R^\mp\rangle \quad (34)$$

- CS implies a conjugated-pseudo symmetry

# Winding Number in NH SSH: $u = 0$ [Lie18]

- $u = 0 \implies$  CS!

- 

$$\sigma_z H(k) \sigma_z = -H(k) \quad (33)$$

$$|\psi_R^\pm\rangle = \sigma_z |\psi_R^\mp\rangle \quad (34)$$

- CS implies a conjugated-pseudo symmetry

$$H^\dagger = \sigma_x H^* \sigma_x \quad (35)$$

## Winding Number in NH SSH: $u = 0$ [Lie18]

- $u = 0 \implies$  CS!

- 

$$\sigma_z H(k) \sigma_z = -H(k) \quad (33)$$

$$|\psi_R^\pm\rangle = \sigma_z |\psi_R^\mp\rangle \quad (34)$$

- CS implies a conjugated-pseudo symmetry

$$H^\dagger = \sigma_x H^* \sigma_x \quad (35)$$

$$|\psi_L^\pm\rangle = \sigma |\psi_R^\pm\rangle \quad (36)$$

## Winding Number in NH SSH: $u = 0$ [Lie18]

- $u = 0 \implies$  CS!

- 

$$\sigma_z H(k) \sigma_z = -H(k) \quad (33)$$

$$|\psi_R^\pm\rangle = \sigma_z |\psi_R^\mp\rangle \quad (34)$$

- CS implies a conjugated-pseudo symmetry

$$H^\dagger = \sigma_x H^* \sigma_x \quad (35)$$

$$|\psi_L^\pm\rangle = \sigma |\psi_R^\pm\rangle \quad (36)$$

- Parametrize



# Winding Number in NH SSH: $u = 0$ [Lie18]

- $u = 0 \implies$  CS!

- 

$$\sigma_z H(k) \sigma_z = -H(k) \quad (33)$$

$$|\psi_R^\pm\rangle = \sigma_z |\psi_R^\mp\rangle \quad (34)$$

- CS implies a conjugated-pseudo symmetry

$$H^\dagger = \sigma_x H^* \sigma_x \quad (35)$$

$$|\psi_L^\pm\rangle = \sigma |\psi_R^\pm\rangle \quad (36)$$

- Parametrize

$$|\psi_{R,K}^\pm\rangle = \frac{1}{\sqrt{2 \sin \theta_k \cos \theta_k}} \begin{pmatrix} e^{-i\phi_k} \cos \theta_k \\ \pm \sin \theta_k \end{pmatrix} |\psi_{L,K}^\pm\rangle = \frac{1}{\sqrt{2 \sin \theta_k \cos \theta_k}} \begin{pmatrix} e^{-i\phi_k} \sin \theta_k \\ \pm \cos \theta_k \end{pmatrix} \quad (37)$$

# Winding Number in NH SSH: $u = 0$ [Lie18]

- $u = 0 \implies$  CS!

- 

$$\sigma_z H(k) \sigma_z = -H(k) \quad (33)$$

$$|\psi_R^\pm\rangle = \sigma_z |\psi_R^\mp\rangle \quad (34)$$

- CS implies a conjugated-pseudo symmetry


$$H^\dagger = \sigma_x H^* \sigma_x \quad (35)$$

$$|\psi_L^\pm\rangle = \sigma |\psi_R^\pm\rangle \quad (36)$$

- Parametrize

$$|\psi_{R,K}^\pm\rangle = \frac{1}{\sqrt{2 \sin \theta_k \cos \theta_k}} \begin{pmatrix} e^{-i\phi_k} \cos \theta_k \\ \pm \sin \theta_k \end{pmatrix} |\psi_{L,K}^\pm\rangle = \frac{1}{\sqrt{2 \sin \theta_k \cos \theta_k}} \begin{pmatrix} e^{-i\phi_k} \sin \theta_k \\ \pm \cos \theta_k \end{pmatrix} \quad (37)$$

# Winding Number in NH SSH: $u \neq 0$ [Lie18]



**Figure 11:** One-dimensional phase diagram in  $\nu$  for the symmetric case  $v_1 = v_2 = \nu \in \mathbb{R}$ ,  $w_1 = w_2 = w \in \mathbb{R}$ . The system moves in and out of the broken phase as a function of  $\nu$ .  $\nu$  is the topological index which predicts how many pairs of gapless-real-energy, edge modes exist in the system. The bulk spectrum is gapless in the entire -broken phase. [Lie18]

# And that is all!

# And that is all!

- We focussed on reviewing the basic mechanisms underlying NH Topology.

# And that is all!

- We focussed on reviewing the basic mechanisms underlying NH Topology.
- Paves the way for more!

# And that is all!

- We focussed on reviewing the basic mechanisms underlying NH Topology.
- Paves the way for more!
  - ① [Gon+18] gives a systematic classification of NH topology based on K theory and obtained the periodic table for non-Hermitian AZ classes..

# And that is all!

- We focussed on reviewing the basic mechanisms underlying NH Topology.
- Paves the way for more!
  - 1 [Gon+18] gives a systematic classification of NH topology based on K theory and obtained the periodic table for non-Hermitian AZ classes..
  - 2 [Kaw+19] follows up a detailed analysis of symmetry in NH topology..



# And that is all!

- We focussed on reviewing the basic mechanisms underlying NH Topology.
- Paves the way for more!
  - 1 [Gon+18] gives a systematic classification of NH topology based on K theory and obtained the periodic table for non-Hermitian AZ classes..
  - 2 [Kaw+19] follows up a detailed analysis of symmetry in NH topology..
  - 3 [SAS20] proposes an alternative designing principle to realize scattering-free edge current in non-Hermitian devices..

## And that is all!

- We focussed on reviewing the basic mechanisms underlying NH Topology.
- Paves the way for more!
  - 1 [Gon+18] gives a systematic classification of NH topology based on K theory and obtained the periodic table for non-Hermitian AZ classes..
  - 2 [Kaw+19] follows up a detailed analysis of symmetry in NH topology..
  - 3 [SAS20] proposes an alternative designing principle to realize scattering-free edge current in non-Hermitian devices..
  - 4 [Edv+20] extends the biorthogonal polarization to systems with  $M$  gapless edge modes..

# And that is all!

- We focussed on reviewing the basic mechanisms underlying NH Topology.
- Paves the way for more!
  - 1 [Gon+18] gives a systematic classification of NH topology based on K theory and obtained the periodic table for non-Hermitian AZ classes..
  - 2 [Kaw+19] follows up a detailed analysis of symmetry in NH topology..
  - 3 [SAS20] proposes an alternative designing principle to realize scattering-free edge current in non-Hermitian devices..
  - 4 [Edv+20] extends the biorthogonal polarization to systems with  $M$  gapless edge modes..
  - 5 [Eza19] proposes electric circuits for NH topology models.

## And that is all!

- We focussed on reviewing the basic mechanisms underlying NH Topology.
- Paves the way for more!
  - 1 [Gon+18] gives a systematic classification of NH topology based on K theory and obtained the periodic table for non-Hermitian AZ classes..
  - 2 [Kaw+19] follows up a detailed analysis of symmetry in NH topology..
  - 3 [SAS20] proposes an alternative designing principle to realize scattering-free edge current in non-Hermitian devices..
  - 4 [Edv+20] extends the biorthogonal polarization to systems with  $M$  gapless edge modes..
  - 5 [Eza19] proposes electric circuits for NH topology models.

# References I

- [BBK21] Emil J. Bergholtz, Jan Carl Budich, and Flore K. Kunst. “Exceptional topology of non-Hermitian systems”. In: *Reviews of Modern Physics* 93.1 (2021). Publisher: APS, p. 015005.
- [Ben05] Carl M. Bender. “Introduction to PT-Symmetric Quantum Theory”. In: *Contemporary Physics* 46.4 (July 2005). arXiv:quant-ph/0501052, pp. 277–292. ISSN: 0010-7514, 1366-5812. DOI: [10.1080/00107500072632](https://doi.org/10.1080/00107500072632). URL: <http://arxiv.org/abs/quant-ph/0501052> (visited on 11/25/2022).
- [Bro14] Dorje C. Brody. “Biorthogonal Quantum Mechanics”. en. In: *Journal of Physics A: Mathematical and Theoretical* 47.3 (Jan. 2014). arXiv:1308.2609 [math-ph, physics:quant-ph], p. 035305. ISSN: 1751-8113, 1751-8121. DOI: [10.1088/1751-8113/47/3/035305](https://doi.org/10.1088/1751-8113/47/3/035305). URL: <http://arxiv.org/abs/1308.2609> (visited on 11/26/2022).

## References II

- [CW15] Hui Cao and Jan Wiersig. “Dielectric microcavities: Model systems for wave chaos and non-Hermitian physics”. In: *Reviews of Modern Physics* 87.1 (Jan. 2015). Publisher: American Physical Society, pp. 61–111. DOI: [10.1103/RevModPhys.87.61](https://doi.org/10.1103/RevModPhys.87.61). URL: <https://link.aps.org/doi/10.1103/RevModPhys.87.61> (visited on 11/26/2022).
- [Edv+20] Elisabet Edvardsson et al. “Phase Transitions and Generalized Biorthogonal Polarization in Non-Hermitian Systems”. en. In: *Physical Review Research* 2.4 (Oct. 2020). arXiv:2006.12898 [cond-mat, physics:physics, physics:quant-ph], p. 043046. ISSN: 2643-1564. DOI: [10.1103/PhysRevResearch.2.043046](https://doi.org/10.1103/PhysRevResearch.2.043046). URL: <http://arxiv.org/abs/2006.12898> (visited on 11/28/2022).
- [Eza19] Motohiko Ezawa. “Electric circuits for non-Hermitian Chern insulators”. In: *Physical Review B* 100.8 (2019). Publisher: APS, p. 081401.
- [Gon+18] Zongping Gong et al. “Topological phases of non-Hermitian systems”. In: *Physical Review X* 8.3 (2018). Publisher: APS, p. 031079.

## References III

- [Kaw+19] Kohei Kawabata et al. “Symmetry and topology in non-Hermitian physics”. In: *Physical Review X* 9.4 (2019). Publisher: APS, p. 041015.
- [Kun+18] Flore K. Kunst et al. “Biorthogonal bulk-boundary correspondence in non-Hermitian systems”. In: *Physical review letters* 121.2 (2018). Publisher: APS, p. 026808.
- [Lie18] Simon Lieu. “Topological phases in the non-Hermitian Su-Schrieffer-Heeger model”. en. In: *Physical Review B* 97.4 (Jan. 2018). arXiv:1709.03788 [cond-mat], p. 045106. ISSN: 2469-9950, 2469-9969. DOI: [10.1103/PhysRevB.97.045106](https://doi.org/10.1103/PhysRevB.97.045106). URL: <http://arxiv.org/abs/1709.03788> (visited on 11/25/2022).
- [LJS14] Ling Lu, John D. Joannopoulos, and Marin Soljačić. “Topological photonics”. en. In: *Nature Photonics* 8.11 (Nov. 2014). Number: 11 Publisher: Nature Publishing Group, pp. 821–829. ISSN: 1749-4893. DOI: [10.1038/nphoton.2014.248](https://doi.org/10.1038/nphoton.2014.248). URL: <https://www.nature.com/articles/nphoton.2014.248> (visited on 11/26/2022).

## References IV

- [MA19] Mohammad-Ali Miri and Andrea Alù. “Exceptional points in optics and photonics”. In: *Science* 363.6422 (2019). Publisher: American Association for the Advancement of Science, eaar7709.
- [Rot09] Ingrid Rotter. “A non-Hermitian Hamilton operator and the physics of open quantum systems”. en. In: *Journal of Physics A: Mathematical and Theoretical* 42.15 (Mar. 2009), p. 153001. ISSN: 1751-8121. DOI: [10.1088/1751-8113/42/15/153001](https://doi.org/10.1088/1751-8113/42/15/153001). URL: <https://dx.doi.org/10.1088/1751-8113/42/15/153001> (visited on 11/26/2022).



## References V

- [SAC06] A V Sokolov, A A Andrianov, and F Cannata. “Non-Hermitian quantum mechanics of non-diagonalizable Hamiltonians: puzzles with self-orthogonal states”. *en. In: Journal of Physics A: Mathematical and General* 39.32 (Aug. 2006), pp. 10207–10227. ISSN: 0305-4470, 1361-6447. DOI: 10.1088/0305-4470/39/32/S20. URL: <https://iopscience.iop.org/article/10.1088/0305-4470/39/32/S20> (visited on 11/26/2022).
- [SAS20] Kazuki Sone, Yuto Ashida, and Takahiro Sagawa. “Exceptional non-Hermitian topological edge mode and its application to active matter”. *en. In: Nature Communications* 11.1 (Nov. 2020). Number: 1 Publisher: Nature Publishing Group, p. 5745. ISSN: 2041-1723. DOI: 10.1038/s41467-020-19488-0. URL: <https://www.nature.com/articles/s41467-020-19488-0> (visited on 11/28/2022).

## References VI

- [SF18] Huitao Shen and Liang Fu. “Quantum Oscillation from In-Gap States and a Non-Hermitian Landau Level Problem”. In: *Physical Review Letters* 121.2 (July 2018). Publisher: American Physical Society, p. 026403. DOI: [10.1103/PhysRevLett.121.026403](https://doi.org/10.1103/PhysRevLett.121.026403). URL: <https://link.aps.org/doi/10.1103/PhysRevLett.121.026403> (visited on 11/26/2022).
- [YM19] Kazuki Yokomizo and Shuichi Murakami. “Non-Bloch Band Theory of Non-Hermitian Systems”. en. In: *Physical Review Letters* 123.6 (Aug. 2019), p. 066404. ISSN: 0031-9007, 1079-7114. DOI: [10.1103/PhysRevLett.123.066404](https://doi.org/10.1103/PhysRevLett.123.066404). URL: <https://link.aps.org/doi/10.1103/PhysRevLett.123.066404> (visited on 11/25/2022).
- [ZW19] Qi Zhang and Biao Wu. “Non-Hermitian quantum systems and their geometric phases”. In: *Physical Review A* 99.3 (2019). Publisher: APS, p. 032121.

## References VII

- [ZW20] Qi Zhang and Biao Wu. “Monopoles in non-Hermitian systems”. In: *Journal of Physics A: Mathematical and Theoretical* 53.6 (2020). Publisher: IOP Publishing, p. 065203.
- [ZYF20] Kai Zhang, Zhesen Yang, and Chen Fang. “Correspondence between Winding Numbers and Skin Modes in Non-Hermitian Systems”. en. In: *Physical Review Letters* 125.12 (Sept. 2020), p. 126402. ISSN: 0031-9007, 1079-7114. DOI: [10.1103/PhysRevLett.125.126402](https://doi.org/10.1103/PhysRevLett.125.126402). URL: <https://link.aps.org/doi/10.1103/PhysRevLett.125.126402> (visited on 11/25/2022).