


Motion in a Straight Line

Distance versus Displacement

Total length of path (ACB) covered by the particle is called distance. Displacement vector or displacement is the minimum distance (AB) and directed from initial position to final position.




Displacement is Change of Position Vector

From ΔOAB $\Delta \vec{r} = \vec{r}_B - \vec{r}_A$

$$\vec{r}_B = x_2\hat{i} + y_2\hat{j} + z_2\hat{k}$$

and $\vec{r}_A = x_1\hat{i} + y_1\hat{j} + z_1\hat{k}$

$$\Delta \vec{r} = (x_2 - x_1)\hat{i} + (y_2 - y_1)\hat{j} + (z_2 - z_1)\hat{k}$$



$$\text{Average velocity} = \frac{\text{Displacement}}{\text{Time interval}} \Rightarrow \vec{v}_{av} = \frac{\Delta \vec{r}}{\Delta t}$$

$$\text{Average speed} = \frac{\text{Distance travelled}}{\text{Time interval}}$$

For uniform motion

Average speed = | average velocity | = | instantaneous velocity |

$$\text{Velocity } \vec{v} = \frac{d\vec{r}}{dt} = \frac{d}{dt}(x\hat{i} + y\hat{j} + z\hat{k})$$

$$= \frac{dx}{dt}\hat{i} + \frac{dy}{dt}\hat{j} + \frac{dz}{dt}\hat{k} = v_x\hat{i} + v_y\hat{j} + v_z\hat{k}$$

$$\text{Average Acceleration} = \frac{\text{Total change in velocity}}{\text{Total time taken}} = \vec{a}_{av} = \frac{\Delta \vec{v}}{\Delta t}$$

Acceleration

$$\vec{a} = \frac{d\vec{v}}{dt} = \frac{d}{dt}(v_x\hat{i} + v_y\hat{j} + v_z\hat{k})$$

$$= \frac{dv_x}{dt}\hat{i} + \frac{dv_y}{dt}\hat{j} + \frac{dv_z}{dt}\hat{k} = a_x\hat{i} + a_y\hat{j} + a_z\hat{k}$$

Important Points About 1D Motion

- Distance $\geq$ | displacement | and Average speed $\geq$ | average velocity |
- If distance $>$ | displacement | this implies
(a) atleast at one point in path, velocity is zero.



Motion with Constant Acceleration: Equations of Motion

❖ In vector form

$$\vec{v} = \vec{u} + \vec{a}t \text{ and}$$

$$\Delta \vec{r} = \vec{r}_2 - \vec{r}_1, \vec{s} = \left(\frac{\vec{u} + \vec{v}}{2} \right) t = \vec{u}t + \frac{1}{2}\vec{a}t^2 = \vec{v}t - \frac{1}{2}\vec{a}t^2$$

$$v^2 = u^2 + 2\vec{a} \cdot \vec{s} \text{ and } \vec{s}_{n^{\text{th}}} = \vec{u} + \frac{\vec{a}}{2}(2n-1)$$

($S_{n^{\text{th}}}$ $\rightarrow$ displacement in n^{th} second)

❖ In scalar form (for one dimensional motion):






$$v = u + at \quad s = \left(\frac{u + v}{2} \right) t = ut + \frac{1}{2}at^2 = vt - \frac{1}{2}at^2$$







$$v^2 = u^2 + 2as \quad s_n = u + \frac{a}{2}(2n-1)$$

Uniform Motion

If an object is moving along the straight line covers equal distance in equal interval of time, it is said to be in uniform motion along a straight line.




Different Graphs of Motion

Displacement-Time Graph		
(a) For a stationary body Displacement 	(b) Body moving with a constant velocity Displacement 	(c) Body moving with a constant acceleration Displacement 
(d) Body moving with a constant retardation Displacement 	(e) Body moving with infinite velocity. But such motion of body is never possible Displacement 	

Velocity-Time Graph		
(a) Moving with a constant velocity Velocity 	(b) Moving with a constant acceleration having zero initial velocity Velocity 	(c) Body moving with a constant retardation and its initial velocity is not zero Velocity 
(d) Moving with negative acceleration with zero initial velocity 	(e) Moving with increasing acceleration Velocity 	(f) Moving with decreasing retardation Velocity 

Note: Slope of velocity-time graph gives acceleration.

Acceleration-Time Graph

(a) When object is moving with constant acceleration Acceleration 	(b) When object is moving with constant increasing acceleration Acceleration 	(c) When object is moving with constant decreasing acceleration Acceleration 
--	--	--

Motion under Gravity (No Air Resistance)


If an object is falling freely under gravity and downward direction is taken as positive, then equations of motion becomes

$$(i) v = u + gt \quad (ii) h = ut + \frac{1}{2}gt^2 \quad (iii) v^2 = u^2 + 2gh$$

Note: If upward direction is taken as positive then g is replaced by $-g$ in above three equations.


If a body is thrown vertically up with a velocity u in the uniform gravitational field then

- (i) Maximum height attained $H = \frac{u^2}{2g}$
- (ii) Time of ascent = time of descent = $\frac{u}{g}$
- (iii) Total time of flight = $\frac{2u}{g}$
- (iv) Final velocity at the point of projection = u (downwards)
- (v) **Gallileo's law of odd numbers:** For a freely falling body ratio of successive distance covered in equal time interval ' t '



$$S_1 : S_2 : S_3 : \dots, S_n = 1 : 3 : 5 : \dots, 2n - 1$$

- (vi) At any point on its path the body will have same speed for upward journey and downward journey.



- (vii) If a body thrown upwards crosses a point in time t_1 and t_2 respectively then height of point $h = \frac{1}{2} g t_1 t_2$. Maximum height $H = \frac{1}{2} g (t_1 + t_2)^2$.

- (viii) A body is thrown upward, downward and horizontally with same speed takes time t_1 , t_2 and t_3 respectively to reach the ground then $t_3 = \sqrt{t_1 t_2}$ and height from where the particle was thrown is $H = \frac{1}{2} g t_1 t_2$.

