Total No. of Questions—8]

[Total No. of Printed Pages—4

Seat	
No.	3

[5559]-195

S.E. (Comp/IT) (II Semester) EXAMINATION, 2019

ENGINEERING MATHEMATICS—III

(2015 **PATTERN**)

Time: Three Hours

Maximum Marks: 60

- Neat diagrams must be drawn wherever necessary.
 - Figures to the right indicate full marks.
 - Use of electronic pocket calculator is allowed.
 - Assume suitable data, if necessary. (iv)
- Solve any two differential equations: 1. (a)

(i)
$$\frac{d^2y}{dx^2} - 4\frac{dy}{dx} + 4y = e^{2x}\sin 4x + 2^{3x} + 6$$

(ii)
$$x^2 \frac{d^2y}{dx^2} + x \frac{dy}{dx} - y = x^4 + 3x + 1$$

- (ii) $x^2 \frac{d^2y}{dx^2} + x \frac{dy}{dx} y = x^4 + 3x + 1$ (iii) $\frac{d^2y}{dx^2} + 9y = \tan 3x$, by using the method of variation of parameters.
- Solve the integral equation: (*b*)

[4]

the integral equation :
$$\int_{0}^{\infty} f(x) \cos \lambda x \ dx = \begin{cases} 2 - \lambda, \ 0 \le \lambda \le 2 \\ 0, \ \lambda > 2 \end{cases}$$

P.T.O.

[8]

- 2. (a) An inductor of 0.25 henries, with negligible resistance, a capacitor of 0.04 farads are connected in series and having an alternating voltage [12 sin 6t]. Find the current and charge at any time t. [4]
 - (b) Solve any one of the following : $(i) \quad \text{Obtain } z[4^k e^{-6k}], \ k \geq 0.$
 - (ii) Obtain $z^{-1} \left[\frac{13z}{(5z+1)(4z+1)} \right]$.
 - (c) Solve the difference equation : f(k+2) 7f(k+1) + 12f(k) = 0 where f(0) = 0, f(1) = 3, $k \ge 0$.
- 3. (a) The first four moments of a distribution about the value 5 are 2, 20, 40 and 50. Obtain the first four central moments, β_1 and β_2 . [4]
 - (b) Fit a straight line of the form Y = aX + b to the following data by the least square method: [4]

X	1	3	4	5	6	8
Y	-3	1	3	5	7	11

(c) A riddle is given to three students whose probabilities of solving it are $\frac{1}{2}$, $\frac{1}{3}$ and $\frac{1}{4}$ respectively. Find the probability that the riddle is solved. [4]

[5559]-195

4. (a) In a sample of 1,000 cases, the mean of a certain examination is 14 and standard deviation is 2.5. Assuming the distribution to be normal. Find the number of students scoring between 12 and 15.

[Given: $Z_1 = 0.4$, $A_1 = 0.1554$, $Z_2 = 0.8$, $A_2 = 0.2881$]

- (b) During working hours, on an average 3 phone calls are coming into a company within an hour. Using Poisson distribution, find the probability that during a particular working hour, there will be at the most one phone call. [4]
- (c) For a bivariate data, the regression equation of Y on X is 8x 10y = -66 and the regression equation of X on Y is 40x 18y = 214. Find the mean values of X and Y. Also, find the correlation coefficient between X and Y. [4]
- 5. (a) Find the directional derivative of $\phi = xy^2 + yz^2 + zx^2$ at (1, 1, 1) along the line 2(x-2) = y + 1 = z 1. [4]
 - (b) Find constants a, b, c so that

$$\overline{\mathbf{F}} = (x + 2y + az)\overline{i} + (bx - 3y - z)\overline{j} + (4x + cy + 2z)\overline{k}$$

is irrotational. [4]

(c) Find the workdone by the force

$$\overline{\mathbf{F}} = (x^2 - yz)\overline{i} + (y^2 - zx)\overline{j} + (z^2 - xy)\overline{k}$$

in taking a particle from (0, 0, 0) to (1, 2, 1). [5]

[5559]-195 3 P.T.O.

Show that (any one) **6.** (a)[4]

$$(i) \qquad \nabla \cdot \left(\frac{\overline{a} \times \overline{r}}{r^n}\right) = 0 \qquad (ii) \qquad \nabla^2 \left[\nabla \cdot \left(\frac{\overline{r}}{r^2}\right)\right] = \frac{2}{r^4} .$$

- Find the directional derivative of $\phi = 4xz^3 3x^2y^2$ (*b*) (2, -1, 2) along the tangent to the curve $x = e^t \cos t$, $y = e^t \sin t$, $z = e^t \cot t = 0$. [4]
- Find the workdone by, $\overline{F} = 2xy^2\overline{i} + (2x^2y + y)\overline{j}$ in taking a (c)particle from $(0,\ 0,\ 0)$ to $(2,\ 4,\ 0)$ along the parabola $y = x^2, z = 0.$ [5]
- Determine the analytic function f(z) = u + iv if u =**7.** (a) $2x - x^3 + 3xy^2.$ [4]
 - Find the bilinear transformation that maps to points (*b*) z = -i, 0, i into the points W = 1, 0, ∞ .
 - Evaluate $\int \frac{z^3}{z^2-4} dz$, where c is the circle |z| = 3. (c)

- the analytic function $f(z) = u + 2x^2 y^3 2y^2$. 8. (a) $u = 3x^2y + 2x^2 - y^3 - 2y^2.$ [4]
 - Find image of the circle |z 2i| = 2, under the mapping (*b*) $w = \frac{1}{2}$ [4]
 - Evaluate $\int \frac{2z^2+z}{z^2-1} dz$, where c is the circle $|z| = \frac{3}{2}$. [5] (*c*)