03/12/2026

ABHA ID -

Mr. Dummy 1,

50/M, 70 kg,

Patna, Bihar.

C/O - Follow-up after 2 months. Polyuria and thirst reduced. No hypoglycemic symptoms.

I → FBS – 124 mg/dl; PPBS – 178 mg/dl; HbA1c – 6.5%.

Δ → **Type 2 Diabetes**

Adv :-

1) Continue Tab. Metformin 500 mg oral before meal, 1 × 3 tab.

2) Low calorie diet.

3) Brisk walking / regular physical activity.

4) Avoid high sugar-content foods.

5) Follow-up after 3 months.

Dr. XYZ

Reg. 12345

RESEARCH USE ONLY — NOT A REAL PRESCRIPTION

03/03/2027

ABHA ID -

Mr. Dummy 1,

50/M, 70 kg,

Patna, Bihar.

C/O - Generally well. Occasional dietary indiscretion. No polyuria/polydipsia.

I → FBS – 118 mg/dl; PPBS – 165 mg/dl; HbA1c – 6.3%. Renal function – within reference range.

Δ → **Type 2 Diabetes**

Adv :-

1) Continue existing metformin regimen.

2) Continue low calorie diet and brisk walking.

3) Monitor blood glucose periodically.

4) Continue avoidance of high-sugar foods.

5) Follow-up after 3 months.

Dr. XYZ

Reg. 12345

03/06/2027

ABHA ID -

Mr. Dummy 1,

50/M, 70 kg,

Patna, Bihar.

C/O - No polyuria or polydipsia. No new complaints.

I → FBS – 112 mg/dl; PPBS – 154 mg/dl; HbA1c – 6.1%.

Δ → **Type 2 Diabetes**

Adv :-

1) Continue existing metformin regimen.
2) Continue dietary modification.
3) Regular brisk walking / physical activity.
4) Periodic HbA1c and renal profile.
5) Follow-up after 3 months.

Dr. XYZ

Reg. 12345

RESEARCH USE ONLY — NOT A REAL PRESCRIPTION

03/09/2027

ABHA ID -

Mr. Dummy 1,

50/M, 70 kg,

Patna, Bihar.

C/O - No new complaints. Clinically stable.

I → FBS – 109 mg/dl; PPBS – 148 mg/dl; HbA1c – 6.0%.

Δ → **Type 2 Diabetes**

Adv :-

1) Continue existing treatment.
2) Continue low calorie diet.
3) Continue regular physical activity.
4) Avoid high sugar-content foods.
5) Follow-up after 3 months.

Dr. XYZ

Reg. 12345

RESEARCH USE ONLY — NOT A REAL PRESCRIPTION

03/12/2027

ABHA ID -

Mr. Dummy 1,

50/M, 70 kg,

Patna, Bihar.

C/O - Asymptomatic. No polyuria or polydipsia.

I → FBS – 106 mg/dl; PPBS – 142 mg/dl; HbA1c – 5.9%.

Δ → **Type 2 Diabetes**

Adv :-

1) Continue existing treatment.
2) Continue dietary modification and physical activity.
3) Periodic diabetes monitoring.
4) Continue avoidance of high-sugar foods.
5) Follow-up as advised.

Dr. XYZ

Reg. 12345

RESEARCH USE ONLY — NOT A REAL PRESCRIPTION