


Worm up


A worm is crawling accross the table. The path of the worm looks something like this:



True or False?

The position of the worm is a function.

Worm function




A worm is crawling accross the table.

For any time t , let $f(t)$ be the position of the worm.

This defines a function f .

1. What is the domain of f ?
2. What is the codomain of f ?

Finding a Restricted Domain on which a Function is Invertible




1. Find the largest interval containing 0 on which f is invertible.
2. Find the largest interval containing 1 on which f is invertible.

Fill in the Blank

Given that f is an invertible function, fill in the blanks.

1. If $f(-1) = 0$, then $f^{-1}(0) = \text{——}$.
2. If $f^{-1}(2) = 1$, then $f(1) = \text{——}$.
3. If $(2, 3)$ is on the graph of f , then —— is on the graph of f^{-1} .
4. If —— is on the graph of f , then $(-2, 4)$ is on the graph of f^{-1} .

Inverse function from a graph



Calculate:

1. $f(2)$
2. $f(0)$
3. $f^{-1}(2)$
4. $f^{-1}(0)$
5. $f^{-1}(-1)$

Let

$$h(x) = x|x| + 1$$

1. Calculate $h^{-1}(-8)$.

Let

$$h(x) = x|x| + 1$$

1. Calculate $h^{-1}(-8)$.
2. Sketch the graph of h .
3. Find an equation for h^{-1} .
4. Sketch the graph of h^{-1} .
5. Verify that
 - for every $t \in \boxed{???}$, $h(h^{-1}(t)) = t$.
 - for every $t \in \boxed{???}$, $h^{-1}(h(t)) = t$.

Functions, inverses, and graphs

Sketch the graph of a function g satisfying all the following properties:

1. The domain of g is \mathbb{R} .
2. g is continuous everywhere except at -2 .
3. g is differentiable everywhere except at -2 and 1 .
4. g has an inverse function.
5. $g(0) = 2$
6. $g'(0) = 2$
7. $(g^{-1})'(-3) = -2$.

Functions, inverses, and graphs - 2

Draw the graph of a function f satisfying all of the following:

1. The domain of f is \mathbb{R} .
2. f is differentiable everywhere.
3. The restriction of f to $[0, \infty)$ is one-to-one, and its INVERSE has a vertical tangent line at 2.
4. The restriction of f to $(-\infty, 0]$ is one-to-one, and its INVERSE has derivative 2 at 2.

Composition and inverses

Assume for simplicity that all functions in this problem have domain \mathbb{R} .

Let f and g be functions. Assume they each have an inverse.

Is $(f \circ g)^{-1} = f^{-1} \circ g^{-1}$?

- If YES, prove it.
- If NO, fix the statement.

Composition and inverses

Assume for simplicity that all functions in this problem have domain \mathbb{R} .

Let f and g be functions. Assume they each have an inverse.

Is $(f \circ g)^{-1} = f^{-1} \circ g^{-1}$?

- If YES, prove it.
- If NO, fix the statement.

If you do not know how to start, experiment with the functions

$$f(x) = x + 1, \quad g(x) = 2x.$$

Composition of one-to-one functions

Assume for simplicity that all functions in this problem have domain \mathbb{R} . Prove the following theorem.

Theorem A

Let f and g be functions.

IF f and g are one-to-one,

THEN $f \circ g$ is one-to-one.

Composition of one-to-one functions

Assume for simplicity that all functions in this problem have domain \mathbb{R} . Prove the following theorem.

Theorem A

Let f and g be functions.

IF f and g are one-to-one,

THEN $f \circ g$ is one-to-one.

Suggestion:

1. Write the definition of what you want to prove.
2. Figure out the formal structure of the proof.
3. Complete the proof (use the hypotheses!)

Composition of one-to-one functions – 2

Assume for simplicity that all functions in this problem have domain \mathbb{R} . Prove the following theorem.

Theorem B

Let f and g be functions.

IF $f \circ g$ is one-to-one, THEN g is one-to-one.

Composition of one-to-one functions – 2

Assume for simplicity that all functions in this problem have domain \mathbb{R} . Prove the following theorem.

Theorem B

Let f and g be functions.

IF $f \circ g$ is one-to-one, THEN g is one-to-one.

Suggestion:

1. Transform the “ $P \implies Q$ ” theorem into an equivalent “(not Q) \implies (not P)” theorem. You will prove that one instead.

Composition of one-to-one functions – 2

Assume for simplicity that all functions in this problem have domain \mathbb{R} . Prove the following theorem.

Theorem B

Let f and g be functions.

IF $f \circ g$ is one-to-one, THEN g is one-to-one.

Suggestion:

1. Transform the " $P \implies Q$ " theorem into an equivalent " $(\text{not } Q) \implies (\text{not } P)$ " theorem. You will prove that one instead.
2. Write the definition of the hypotheses and of the conclusion.
3. Write the proof.

Composition of one-to-one functions – 3

Assume for simplicity that all functions in this problem have domain \mathbb{R} .

Prove the following claim is FALSE with a counterexample.

Claim

Let f and g be functions.

IF $f \circ g$ is one-to-one,

THEN f is one-to-one.

Definition

Let f be a function with domain D . We say that f is *increasing* on D when

$$\forall x_1, x_2 \in D, \quad x_1 < x_2 \implies f(x_1) < f(x_2).$$

1. Prove that if a function is increasing, then it is one-to-one.

Definition

Let f be a function with domain D . We say that f is *increasing* on D when

$$\forall x_1, x_2 \in D, \quad x_1 < x_2 \implies f(x_1) < f(x_2).$$

1. Prove that if a function is increasing, then it is one-to-one.
2. Use this to show that $g(x) = x^5 + 4x^3 + 2x + 1$ has an inverse.
3. Find $(g^{-1})'(1)$.

Where is the error?

- We know that $(f^{-1})' = \frac{1}{f'}$
- Let $f(x) = x^2$, restricted to the domain $x \in (0, \infty)$

$$f'(x) = 2x \quad \text{and} \quad f'(4) = 8$$

- Then $f^{-1}(x) = \sqrt{x}$

$$(f^{-1})'(x) = \frac{1}{2\sqrt{x}} \quad \text{and} \quad (f^{-1})'(4) = \frac{1}{4}$$

- So $(f^{-1})'(4) \neq \frac{1}{f'(4)}$

Derivatives of the inverse function

Let f be a one-to-one function.

Let $a, b \in \mathbb{R}$ such that $b = f(a)$.

1. Obtain a formula for $(f^{-1})'(b)$ in terms of $f'(a)$.

Hint: This was done in Video 4.4

Take $\frac{d}{dy}$ of both sides of $f(f^{-1}(y)) = y$.

Derivatives of the inverse function

Let f be a one-to-one function.

Let $a, b \in \mathbb{R}$ such that $b = f(a)$.

1. Obtain a formula for $(f^{-1})'(b)$ in terms of $f'(a)$.

Hint: This was done in Video 4.4

Take $\frac{d}{dy}$ of both sides of $f(f^{-1}(y)) = y$.

2. Obtain a formula for $(f^{-1})''(b)$ in terms of $f'(a)$ and $f''(a)$.

Derivatives of the inverse function

Let f be a one-to-one function.

Let $a, b \in \mathbb{R}$ such that $b = f(a)$.

1. Obtain a formula for $(f^{-1})'(b)$ in terms of $f'(a)$.

Hint: This was done in Video 4.4

Take $\frac{d}{dy}$ of both sides of $f(f^{-1}(y)) = y$.

2. Obtain a formula for $(f^{-1})''(b)$ in terms of $f'(a)$ and $f''(a)$.
3. *Challenge:* Obtain a formula for $(f^{-1})'''(b)$ in terms of $f'(a)$, $f''(a)$, and $f'''(a)$.

Compute the derivative of the following functions:

1. $f(x) = e^{\sin x + \cos x} \ln x$

2. $f(x) = \pi^{\tan x}$

3. $f(x) = \ln [e^x + \ln \ln \ln x]$

4. $f(x) = \log_{10} (2x + 3)$

Logarithm and Absolute Value

The function F is defined by the equation

$$F(x) = \ln |x|.$$

What is its derivative?

1. $F'(x) = \frac{1}{x}$

2. $F'(x) = \frac{1}{|x|}$

3. F is not differentiable

A different type of logarithm

Calculate the derivative of

$$f(x) = \log_{x+1}(x^2 + 1)$$

Note: This is a new function. We have not given you a formula for it yet, That is on purpose.

A different type of logarithm

Calculate the derivative of

$$f(x) = \log_{x+1}(x^2 + 1)$$

Note: This is a new function. We have not given you a formula for it yet, That is on purpose.

Hint: If you do not know where to start, remember the definition of logarithm:

$$\log_a b = c \iff a^c = b.$$

Calculate the derivative of

$$g(x) = x^{\tan x}.$$

More logarithmic differentiation

Calculate the derivative of

$$f(x) = (\sin x)^{\cos x} + (\cos x)^{\sin x}.$$

More logarithmic differentiation

Calculate the derivative of

$$f(x) = (\sin x)^{\cos x} + (\cos x)^{\sin x}.$$

What is wrong with this answer?

$$\ln f(x) = (\cos x) \ln(\sin x) + (\sin x)(\ln \cos x)$$

$$\frac{d}{dx} [\ln f(x)] = \frac{d}{dx} [(\cos x) \ln(\sin x)] + \frac{d}{dx} [(\sin x)(\ln \cos x)]$$

$$\begin{aligned} \frac{f'(x)}{f(x)} &= -(\sin x) \ln(\sin x) + (\cos x) \frac{\cos x}{\sin x} \\ &\quad + (\cos x) \ln(\cos x) + (\sin x) \frac{-\sin x}{\cos x} \end{aligned}$$

$$f'(x) = f(x) \left[-(\sin x) \ln(\sin x) + (\cos x) \ln(\cos x) + \frac{\cos^2 x}{\sin x} - \frac{\sin^2 x}{\cos x} \right]$$

Calculate the derivative of

$$h(x) = \sqrt[3]{\frac{(\sin^6 x) \sqrt{x^7 + 6x + 2}}{3^x (x^{10} + 2x)^{10}}}$$

An Implicit Function

Find y' if $x^y = y^x$.

Definition of arctan

1. Sketch the graph of \tan .
2. Prove that \tan is not one-to-one.
3. Select the largest interval containing 0 such that the restriction of \tan to it is one-to-one. We define \arctan as the inverse of this restriction. Let $x, y \in \mathbb{R}$

$$\arctan y = x \quad \Longleftrightarrow \quad ???$$

4. What is the domain of \arctan ? What is the range of \arctan ? Sketch the graph of \arctan .
5. Compute

5.1 $\arctan(\tan(1))$

5.2 $\arctan(\tan(3))$

5.3 $\arctan\left(\tan\left(\frac{\pi}{2}\right)\right)$

5.4 $\arctan(\tan(-6))$

5.5 $\tan(\arctan(0))$

5.6 $\tan(\arctan(10))$

Derivative of arctan

Obtain (and prove) a formula for the derivative of arctan.

Hint: Call $f(t) = \arctan t$ and differentiate

$$\forall t \in \dots \quad \tan(f(t)) = t$$

Computations - Inverse trig functions

Compute the derivatives of these functions, and simplify them as much as possible:

1. $f(x) = \arcsin \left(x^{3/2} \right)$

2. $f(x) = 2x^2 \arctan(x^2) - \ln(x^4 + 1)$

Find simple expressions for these quantities and state the domain on which they are valid:

1. $\sin (\arccos x)$

3. $\sec (\arctan x)$

2. $\sec (\arccos x)$

4. $\tan (\operatorname{arcsec} x)$

Trig-inverse-trig

Find simple expressions for these quantities and state the domain on which they are valid:

1. $\sin (\arccos x)$

3. $\sec (\arctan x)$

2. $\sec (\arccos x)$

4. $\tan (\operatorname{arcsec} x)$

Hint: There are two standard ways to attack these problems:

- Use a trig identity
e.g.: a trig identity relating \sin and \cos for (1)
- Or draw a right triangle with side lengths 1 and x
e.g.: with an angle θ such that $\cos \theta = x$ for (1)

If you need to take a square root, you must justify which branch (+ or -) you are choosing.

1. Complete: “We define arcsec as the inverse of the restriction of sec to ...”
Hint: Sketch the graph of sec.
2. What are the domain and range of arcsec?
Sketch its graph.
3. Obtain (and prove) a formula for the derivative of arcsec in the same way you did for arctan.
4. Now obtain the same formula in a different way: use $\sec x = \frac{1}{\cos x}$ to write arcsec in terms of arccos.