

PHYS 357 Pset 3. Due 11:59 PM Thursday Sep. 26

1. Starting with the angular momentum operators (J_x, J_y, J_z) you worked out in the last problem set (which should have been

$$J_z = \frac{\hbar}{2} \begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix}, \quad J_x = \frac{\hbar}{2} \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}, \quad J_y = \frac{\hbar}{2} \begin{bmatrix} 0 & -i \\ i & 0 \end{bmatrix} \quad (1)$$

show that you get the canonical commutation relations

$$[J_x, J_y] = i\hbar J_z, \quad [J_y, J_z] = i\hbar J_x, \quad [J_z, J_x] = i\hbar J_y \quad (2)$$

where $[A, B] = AB - BA$ for matrices A and B.

2. Consider two coordinate systems (x, y, z) and (x', y', z) that have the same z-axis, but have their x- and y-axes rotated by some angle γ relative to each other (note the lack of a prime on z). Show that if I rotate a vector in the (x, y, z) coordinate system through some angle θ about the z-axis, I get the same physical vector if I transform the vector to the (x', y', z) coordinate system and carry out the rotation there.
3. If I have a set of arbitrary basis vectors (x', y', z') , where you have the (x, y, z) representation of each basis vector, write down the matrix that converts a vector in the (x, y, z) basis to one in the (x', y', z') basis, and vice-versa. How are these two matrices related? I suggest you write a function

`genbasis(xyz)`

that returns a valid matrix of basis vectors where xyz is one of the principal axes (a reminder that this isn't unique so you'll have to make a choice. The results of Problem 2 show that this choice won't matter so just do something easy.)

4. We're now ready to work out an rotation matrix in 3D about an arbitrary axis. Start by picking a random rotation axis in the \hat{n} , direction. Start with random θ, ϕ and convert that to a vector in (x, y, z) coordinates. Then generate a pair of basis vectors to make an

orthogonal space (hint - if you take the cross product of two vectors, you get something that's perpendicular to both, so e.g. $\hat{z} \times \hat{n}$ will be perpendicular to both \hat{z} and \hat{n}). Don't forget that basis vectors must have unit length. Now write out the matrices that convert from (x, y, z) to the \hat{n} basis, rotate by an angle γ in that basis, then convert back to (x, y, z) . I suggest you do this on a computer, and write a function

```
genrot(xyz,gamma)
```

that generates the individual matrices, multiplies them together, and returns the final rotation matrix in the (x, y, z) coordinate system. What is the rotation matrix for $\theta = \pi/4$, $\phi = \pi/6$, and $\text{gamma}=0.01$?

5. Pretend, in a shocking turn of events, the Earth's crust gets rotated so that the Royal Greenwich Observatory, England (current latitude 51.476852, longitude=0.000) moves to the north pole. What are Montreal's new latitude and longitude? (current: 45.50884, -73.58781). Verify that the current angle between Montreal and Greenwich corresponds to Montreal's new latitude (as it must, because the distance to the pole depends only on latitude, not longitude, and your rotation matrix had better not change relative angles between vectors). Hint - think what axis we have to be rotating about to move Greenwich to the North pole.
6. Use your the results (and code!) to show numerically that the rotation commutation relations hold in an arbitrary coordinate system. If you haven't done this sort of thing before, you can show that as you make gamma smaller, the largest term in the error shrinks faster than the largest term in the commutator. I found gammas in the range of 0.01-0.001 were sufficient to show this clearly.