

PHYS 357 Pset 8. Due 11:59 PM Thursday Nov. 14

1. Townsend 6.1.
2. Townsend 6.3.
3. Size of the hydrogen atom. The potential between a proton and electron is $U = -\frac{e^2}{4\pi\epsilon_0 r}$. If we work in a frame where $\langle x \rangle = \langle p \rangle = 0$, then the Heisenberg uncertainty principle tells us we'll have some minimum kinetic energy for a fixed r (remember, the electron can "move" in angular directions, so even if we know say it's at fixed radius, there's still uncertainty in the 3-D position). Find the radius r that minimizes the total energy, in terms of elementary constants, and a constant of order unity (at this level of accuracy, reasonable people may get different values for this constant). Evaluate your radius, and compare to the accepted value for the radius of the ground-state hydrogen, $5.3 \times 10^{-11}m$.

Note - Like many things in quantum mechanics, what we mean by "the" radius is rather fuzzy. This is the Bohr radius, which is the average distance of a ground-state electron from the nucleus for an isolated atom. You'd get a different number if you made molecular hydrogen (H_2), and yet another number if you froze hydrogen into a solid and measured the per-atom volume of that solid.

4. Fourier transform of a Gaussian. Gaussian wave packets get used repeatedly in QM, and so practice with integrals is very valuable. The Fourier transform $F(k)$ of $f(x)$ is defined to be $\int f(x) \exp(ikx) dx$. For a Gaussian, we have $f(x) = \exp(-x^2/2\sigma^2)$. By completing the square, show that

$$\int_{-\infty}^{\infty} \exp(ikx) \exp(-x^2/2\sigma^2) dx = \exp(-k^2\sigma^2) \int (\text{something}) dx$$

Then show that $\int (\text{something}) dx = \int_{-\infty}^{\infty} \exp(-x^2/2\sigma^2) dx$. As a reminder, all closed path integrals in the complex plane are zero if the function has no poles. That integral evaluates to $\sqrt{2\pi\sigma^2}$, giving us the Fourier transform of a Gaussian.

5. Discrete Fourier transform. The discrete Fourier transform of $f(x)$ is defined to be

$$F(k) \equiv \sum_{x=0}^{N-1} f(x) \exp(-2\pi i k x / N)$$

for some integer N , and x, k integers running from 0 to $N - 1$. First, show that

$$\sum_{x=0}^{N-1} \exp(-2\pi i k x / N) = 0$$

unless $k = 0$ (technically, any integer multiple of N , but we restricted k to be between 0 and $N - 1$). As a reminder, the sum of a geometric series $\sum_{m=m_0}^{\infty} r^m = r^{m_0} / (1 - r)$.

Now show that we can get $F(k) = Ff(x)$ for the symmetric matrix F that has $F_{m,n} = \exp(-2\pi i mn / N)$.

Finally, show that $F^{-1} = \frac{1}{N} F^\dagger$. This means for the discrete case, the inverse Fourier transform is just the conjugate of the Fourier transform, with a $1/N$ normalization factor. We'll be able to use this to go easily between position and momentum representations of wave functions.