


Hybrid particle-field molecular dynamics for biological systems

Hyllerås seminar


Sigbjørn Løland Bore
University of Oslo, Norway

Friday, September 6, 2019


Biological scale





Biological scale




Coarse-graining



Coarse-graining




Coarse-graining





$$Z = \int d\Gamma e^{-\beta H(\Gamma)}$$

↓

$$Z \simeq \int d\Gamma_{CG} e^{-\beta H(\Gamma_{CG})}$$



Coarse-graining



$$Z = \int d\Gamma e^{-\beta H(\Gamma)}$$

↓

$$Z \simeq \int d\Gamma_{CG} e^{-\beta H(\Gamma_{CG})}$$

- ▶ Fewer degrees of freedom
- ▶ Speedup of dynamics

The hybrid particle-field method

$$H(\{\mathbf{r}\}) = H_0(\{\mathbf{r}\}) + W[\{\phi(\mathbf{r})\}]$$

Particle-particle-Hamiltonian


Density-field interaction-energy

The hybrid particle-field method

$$H(\{\mathbf{r}\}) = H_0(\{\mathbf{r}\}) + W[\{\phi(\mathbf{r})\}]$$

Particle-particle-Hamiltonian

Density-field interaction-energy




$\{\mathbf{r}\} \equiv \{\mathbf{r}_1, \dots, \mathbf{r}_N\}$, *particle positions*.

The hybrid particle-field method

$$H(\{\mathbf{r}\}) = H_0(\{\mathbf{r}\}) + W[\{\phi(\mathbf{r})\}]$$

Particle-particle-Hamiltonian

Density-field interaction-energy



$\{\mathbf{r}\} \equiv \{\mathbf{r}_1, \dots, \mathbf{r}_N\}$, *particle positions*.


$\{\phi\} \equiv \{\phi_1, \dots, \phi_M\}$, *particle-type number densities*.

The hybrid particle-field method

$$H(\{\mathbf{r}\}) = H_0(\{\mathbf{r}\}) + W[\{\phi(\mathbf{r})\}]$$

Particle-particle-Hamiltonian

Density-field interaction-energy



$$V_k(\mathbf{r}) = \frac{\delta W[\{\phi\}]}{\delta \phi_k(\mathbf{r})}$$


$$\mathbf{F}_i = -\nabla_i V_k(\mathbf{r}_i)$$

$\{\mathbf{r}\} \equiv \{\mathbf{r}_1, \dots, \mathbf{r}_N\}$, particle positions.

$\{\phi\} \equiv \{\phi_1, \dots, \phi_M\}$, particle-type number densities.


Computation of forces

Particle-mesh



Computation of forces

Particle-mesh




1) Linear interpolation:

$$\{\mathbf{r}\} \rightarrow \{\phi_{nml}\}$$

Computation of forces

Particle-mesh



1) Linear interpolation:


$$\{\mathbf{r}\} \rightarrow \{\phi_{nml}\}$$

2) Finite-differences:

$$\{\phi_{nml}\} \rightarrow \{\nabla \phi_{nml}\} \rightarrow \{\nabla V_{nml}\}$$

Computation of forces

Particle-mesh



1) Linear interpolation:

$$\{\mathbf{r}\} \rightarrow \{\phi_{nml}\}$$


2) Finite-differences:

$$\{\phi_{nml}\} \rightarrow \{\nabla \phi_{nml}\} \rightarrow \{\nabla V_{nml}\}$$


3) Force interpolation:

$$\{\nabla V_{nml}\} \rightarrow \mathbf{F}_i$$


Implementation and parallelization



Implementation and parallelization




Implementation and parallelization



Excellent scaling for small and large systems!



No communication



Interaction energy: Polymer-theory

$$W[\phi] = \int d\mathbf{r} \frac{1}{2\rho_0} \left(\sum_{k\ell} \tilde{\chi}_{k\ell} \phi_k(\mathbf{r}) \phi_\ell(\mathbf{r}) + \frac{1}{\kappa} \left(\sum_\ell \phi_\ell(\mathbf{r}) - \rho_0 \right)^2 \right)$$

ρ_0 : density-parameter related to the volume per bead.

Interaction energy: Polymer-theory

$$W[\phi] = \int d\mathbf{r} \frac{1}{2\rho_0} \left(\underbrace{\sum_{k\ell} \tilde{\chi}_{k\ell} \phi_k(\mathbf{r}) \phi_\ell(\mathbf{r})}_{\text{Mixing}} + \frac{1}{\kappa} \left(\sum_\ell \phi_\ell(\mathbf{r}) - \rho_0 \right)^2 \right)$$

$\tilde{\chi}_{k\ell} > 0 \rightarrow$ Likes not to mix

$\tilde{\chi}_{k\ell} \leq 0 \rightarrow$ Likes to mix

ρ_0 : density-parameter related to the volume per bead.

Interaction energy: Polymer-theory

$$W[\phi] = \int d\mathbf{r} \frac{1}{2\rho_0} \left(\underbrace{\sum_{k\ell} \tilde{\chi}_{k\ell} \phi_k(\mathbf{r}) \phi_\ell(\mathbf{r})}_{\text{Mixing}} + \underbrace{\frac{1}{\kappa} \left(\sum_\ell \phi_\ell(\mathbf{r}) - \rho_0 \right)^2}_{\text{Compressibility}} \right)$$

$\tilde{\chi}_{k\ell} > 0 \rightarrow$ Likes not to mix

$\kappa \sim 0 \rightarrow$ incompressible

$\tilde{\chi}_{k\ell} \leq 0 \rightarrow$ Likes to mix

$\kappa \gg 0 \rightarrow$ very compressible

ρ_0 : density-parameter related to the volume per bead.

Interaction energy: Polymer-theory

$$W[\phi] = \int d\mathbf{r} \frac{1}{2\rho_0} \left(\underbrace{\sum_{k\ell} \tilde{\chi}_{k\ell} \phi_k(\mathbf{r}) \phi_\ell(\mathbf{r})}_{\text{Mixing}} + \underbrace{\frac{1}{\kappa} \left(\sum_\ell \phi_\ell(\mathbf{r}) - \rho_0 \right)^2}_{\text{Compressibility}} \right)$$

$\tilde{\chi}_{k\ell} > 0 \rightarrow$ Likes not to mix

$\kappa \sim 0 \rightarrow$ incompressible


$\tilde{\chi}_{k\ell} \leq 0 \rightarrow$ Likes to mix

$\kappa \gg 0 \rightarrow$ very compressible


Net effect $\Rightarrow V_k(\mathbf{r}) = \frac{1}{\rho_0} \left(\sum_\ell \tilde{\chi}_{k\ell} \phi_\ell(\mathbf{r}) + \frac{1}{\kappa} \left(\sum_\ell \phi_\ell(\mathbf{r}) - \rho_0 \right) \right)$

ρ_0 : density-parameter related to the volume per bead.

Hybrid Particle-Field Model for Conformational Dynamics of Peptide Chains



hPF-MD



$$H = H_0(\{\mathbf{r}\}) + W[\rho(\mathbf{r})]$$

S.L Bore et al., JCTC, 2018

Previous work



JCTC 2008

JCTC 2010

JCTC 2013

Previous work




JCTC 2008

JCTC 2010

JCTC 2013

C α -representation



Previous work




JCTC 2008


JCTC 2010

JCTC 2013

C_α-representation




Reconstruction of dipole




Two-bead model

$$V(\gamma, \phi) = V(\gamma) + V(\phi)?$$




Two-bead model

$$V(\gamma, \phi) = V(\gamma) + V(\phi)?$$




4-Alanine




Two-bead model


$$V(\gamma, \phi) = V(\gamma) + V(\phi)?$$



4-Alanine




Metadynamics \rightarrow




Two-bead model


$$V(\gamma, \phi) = V(\gamma) + V(\phi)?$$



4-Alanine




Metadynamics →




$$\rightarrow V(\gamma, \phi) = \frac{1}{2}k(\phi)(\gamma - \gamma_0(\phi))^2 + V_{\text{prop}}(\phi, \lambda)$$

Propensity potential




Propensity potential

Boltzmann Inversion



Propensity potential


Boltzmann Inversion



$$V_{\text{prop}}(\phi, \lambda) = \frac{1}{2} ((|\lambda| - \lambda) V_\alpha(\phi) + (|\lambda| + \lambda) V_\beta(\phi) + (1 - |\lambda|) V_0(\phi))$$

Propensity potential


Boltzmann Inversion



$$V_{\text{prop}}(\phi, \lambda) = \frac{1}{2} ((|\lambda| - \lambda) V_\alpha(\phi) + (|\lambda| + \lambda) V_\beta(\phi) + (1 - |\lambda|) V_0(\phi))$$

Propensity potential

Boltzmann Inversion




$-1 \leq \lambda < 0 \implies \text{Helical}$

$0 < \lambda < 1 \implies \text{Extended}$

$$V_{\text{prop}}(\phi, \lambda) = \frac{1}{2} ((|\lambda| - \lambda) V_\alpha(\phi) + (|\lambda| + \lambda) V_\beta(\phi) + (1 - |\lambda|) V_0(\phi))$$


Propensity potential

Boltzmann Inversion




$$V_{\text{prop}}(\phi, \lambda) = \frac{1}{2} ((|\lambda| - \lambda) V_\alpha(\phi) + (|\lambda| + \lambda) V_\beta(\phi) + (1 - |\lambda|) V_0(\phi))$$

Phase-diagram: homo-poly-peptide






Phase-diagram: homo-poly-peptide

$\tilde{\chi}_{kl}$	CB	H ₂ O
CB	0	α
H ₂ O	α	0



Phase-diagram: homo-poly-peptide

$\tilde{\chi}_{k\ell}$	CB	H ₂ O
CB	0	α
H ₂ O	α	0



i: Random coil ii: α -helix iii: β -hairpin iv: Extended v: Helix-coil-helix vi: β -floor/helix vii: Helical bundle

HP-model

$\tilde{\chi}_{k\ell}$	CB _P	CB _H	H ₂ O
CB _P	0	α	0
CB _H	α	0	α
H ₂ O	0	α	0

HP-model

$\tilde{\chi}_{k\ell}$	CB _P	CB _H	H ₂ O
CB _P	0	α	0
CB _H	α	0	α
H ₂ O	0	α	0

α -part: PHPPHHPPHPHPPHHHP


β -part: PHPHPHPHPHPHPHPH

HP-model



$\tilde{\chi}_{k\ell}$	CB _P	CB _H	H ₂ O
CB _P	0	α	0
CB _H	α	0	α
H ₂ O	0	α	0

α -part: PHPPHHPPHPPPHHP

β -part: PHPHPHPHPHPHPHPH



HP-polymer interacting with membrane



Outlook

- ▶ Parametrization: Toy-model → 20 amino-acids

Outlook

- ▶ Parametrization: Toy-model → 20 amino-acids
 - ▶ Chemical specific $\tilde{\chi}_{kl}$ -parameter
 - ▶ Strategies for modeling $V(\phi, \gamma)$
 - ▶ Machine-learning?
 - ▶ New PhD-student Manuel Carrera

Outlook

- ▶ Parametrization: Toy-model → 20 amino-acids
 - ▶ Chemical specific $\tilde{\chi}_{k\ell}$ -parameter
 - ▶ Strategies for modeling $V(\phi, \gamma)$
 - ▶ Machine-learning?
 - ▶ New PhD-student Manuel Carrera
- ▶ Electrostatics
 - ▶ Particle-field method for electrostatics


Outlook

- ▶ Parametrization: Toy-model → 20 amino-acids
 - ▶ Chemical specific $\tilde{\chi}_{k\ell}$ -parameter
 - ▶ Strategies for modeling $V(\phi, \gamma)$
 - ▶ Machine-learning?
 - ▶ New PhD-student Manuel Carrera
- ▶ Electrostatics
 - ▶ Particle-field method for electrostatics
- ▶ Application of current model
 - ▶ Tsudo Yamanaka, Yamagata University, Japan




Outlook

- ▶ Parametrization: Toy-model → 20 amino-acids
 - ▶ Chemical specific $\tilde{\chi}_{ke}$ -parameter
 - ▶ Strategies for modeling $V(\phi, \gamma)$
 - ▶ Machine-learning?
 - ▶ New PhD-student Manuel Carrera
- ▶ Electrostatics
 - ▶ Particle-field method for electrostatics
- ▶ Application of current model
 - ▶ Tsudo Yamanaka, Yamagata University, Japan




Outlook

- ▶ Parametrization: Toy-model → 20 amino-acids
 - ▶ Chemical specific $\tilde{\chi}_{k\ell}$ -parameter
 - ▶ Strategies for modeling $V(\phi, \gamma)$
 - ▶ Machine-learning?
 - ▶ New PhD-student Manuel Carrera
- ▶ Electrostatics
 - ▶ Particle-field method for electrostatics
- ▶ Application of current model
 - ▶ Tsudo Yamanaka, Yamagata University, Japan




Electrostatics in Hybrid particle-field

hPF-MD + electrostatics




H.B. Kolli et al. JCTC, 2018



S.L. Bore et al., JCTC, 2018

Electrostatic screening: Atomistic vs coarse-grained


Atomistic molecular dynamics:



Electrostatic screening: Atomistic vs coarse-grained

Atomistic molecular dynamics:

- ▶ *Charges are resolved*
- ▶ *Screening is modeled directly*




Electrostatic screening: Atomistic vs coarse-grained

Atomistic molecular dynamics:

- ▶ *Charges are resolved*
- ▶ *Screening is modeled directly*

Coarse-grained molecular dynamics:

- ▶ *Charge resolution is lost*
- ▶ *Screening modeled modelled indirectly*




Electrostatic screening: Atomistic vs coarse-grained

Atomistic molecular dynamics:

- ▶ *Charges are resolved*
- ▶ *Screening is modeled directly*

Coarse-grained molecular dynamics:

- ▶ *Charge resolution is lost*
- ▶ *Screening modeled modelled indirectly*



Idea : $\nabla \cdot (\epsilon(\mathbf{r}) \nabla \psi(\mathbf{r})) = -\rho(\mathbf{r})$ (Generalized Poisson equation)

External potential in a density dependent dielectric

Electrostatic interaction energy:

$$W_{\text{elec}}[\{\phi(\mathbf{r})\}] = \frac{1}{2} \int d\mathbf{r} \frac{\mathbf{D}(\mathbf{r}) \cdot \mathbf{D}(\mathbf{r})}{\epsilon(\mathbf{r})},$$

$\{\phi\}$: number densities. \mathbf{D} : displacement field. ϵ : permittivity.

External potential in a density dependent dielectric

Electrostatic interaction energy:

$$W_{\text{elec}}[\{\phi(\mathbf{r})\}] = \frac{1}{2} \int d\mathbf{r} \frac{\mathbf{D}(\mathbf{r}) \cdot \mathbf{D}(\mathbf{r})}{\epsilon(\mathbf{r})},$$

$\{\phi\}$: number densities. \mathbf{D} : displacement field. ϵ : permittivity.

Potential felt by particles of type K :

$$V_{\text{ext},K}(\mathbf{r}) = \frac{\delta W_{\text{elec}}}{\delta \phi_K(\mathbf{r})}$$

External potential in a density dependent dielectric

Electrostatic interaction energy:

$$W_{\text{elec}}[\{\phi(\mathbf{r})\}] = \frac{1}{2} \int d\mathbf{r} \frac{\mathbf{D}(\mathbf{r}) \cdot \mathbf{D}(\mathbf{r})}{\epsilon(\mathbf{r})},$$

$\{\phi\}$: number densities. \mathbf{D} : displacement field. ϵ : permittivity.

Potential felt by particles of type K :

$$V_{\text{ext},K}(\mathbf{r}) = \frac{\delta W_{\text{elec}}}{\delta \phi_K(\mathbf{r})} = \underbrace{\int d\mathbf{r}' \frac{\delta W_{\text{elec}}}{\delta \mathbf{D}(\mathbf{r}')} \frac{\delta \mathbf{D}(\mathbf{r}')}{\delta \phi_K(\mathbf{r})}} + \underbrace{\frac{\delta W_{\text{elec}}}{\delta \epsilon(\mathbf{r})} \frac{\partial \epsilon(\mathbf{r})}{\partial \phi_K(\mathbf{r})}}$$

External potential in a density dependent dielectric

Electrostatic interaction energy:

$$W_{\text{elec}}[\{\phi(\mathbf{r})\}] = \frac{1}{2} \int d\mathbf{r} \frac{\mathbf{D}(\mathbf{r}) \cdot \mathbf{D}(\mathbf{r})}{\epsilon(\mathbf{r})},$$

$\{\phi\}$: number densities. \mathbf{D} : displacement field. ϵ : permittivity.

Potential felt by particles of type K :

$$V_{\text{ext},K}(\mathbf{r}) = \frac{\delta W_{\text{elec}}}{\delta \phi_K(\mathbf{r})} = \underbrace{\int d\mathbf{r}' \frac{\delta W_{\text{elec}}}{\delta \mathbf{D}(\mathbf{r}')} \frac{\delta \mathbf{D}(\mathbf{r}')}{\delta \phi_K(\mathbf{r})}}_{q_K \psi(\mathbf{r})} + \underbrace{\frac{\delta W_{\text{elec}}}{\delta \epsilon(\mathbf{r})} \frac{\partial \epsilon(\mathbf{r})}{\partial \phi_K(\mathbf{r})}}$$

ψ : electrostatic potential. \mathbf{E} : electrostatic field ($\mathbf{E} = -\nabla \psi = \epsilon \mathbf{D}$).

External potential in a density dependent dielectric

Electrostatic interaction energy:

$$W_{\text{elec}}[\{\phi(\mathbf{r})\}] = \frac{1}{2} \int d\mathbf{r} \frac{\mathbf{D}(\mathbf{r}) \cdot \mathbf{D}(\mathbf{r})}{\epsilon(\mathbf{r})},$$

$\{\phi\}$: number densities. \mathbf{D} : displacement field. ϵ : permittivity.

Potential felt by particles of type K :

$$V_{\text{ext},K}(\mathbf{r}) = \frac{\delta W_{\text{elec}}}{\delta \phi_K(\mathbf{r})} = \underbrace{\int d\mathbf{r}' \frac{\delta W_{\text{elec}}}{\delta \mathbf{D}(\mathbf{r}')} \frac{\delta \mathbf{D}(\mathbf{r}')}{\delta \phi_K(\mathbf{r})}}_{q_K \psi(\mathbf{r})} + \underbrace{\frac{\delta W_{\text{elec}}}{\delta \epsilon(\mathbf{r})} \frac{\partial \epsilon(\mathbf{r})}{\partial \phi_K(\mathbf{r})}}_{-\frac{1}{2} \frac{\partial \epsilon(\mathbf{r})}{\partial \phi_K(\mathbf{r})} |\mathbf{E}(\mathbf{r})|^2}$$

ψ : electrostatic potential. \mathbf{E} : electrostatic field ($\mathbf{E} = -\nabla \psi = \epsilon \mathbf{D}$).

Modelling: density dependence of the dielectric

Modelling: density dependence of the dielectric

Density weighted average:

$$\epsilon(\{\phi(\mathbf{r})\}) = \frac{\sum_K^M \epsilon_K \phi_K(\mathbf{r})}{\phi_0(\mathbf{r})},$$

ϵ_K : dielectric of particle type K . ϕ_0 : local total particle density.

Modelling: density dependence of the dielectric

Density weighted average:

$$\epsilon(\{\phi(\mathbf{r})\}) = \frac{\sum_K^M \epsilon_K \phi_K(\mathbf{r})}{\phi_0(\mathbf{r})},$$

ϵ_K : dielectric of particle type K . ϕ_0 : local total particle density.

Potential felt by particle of type K :

$$V_{\text{ext},K}(\mathbf{r}) = q_K \psi(\mathbf{r}) - \frac{1}{2} \frac{\epsilon_K - \epsilon(\mathbf{r})}{\phi_0(\mathbf{r})} |\mathbf{E}(\mathbf{r})|^2,$$

Modelling: density dependence of the dielectric

Density weighted average:

$$\epsilon(\{\phi(\mathbf{r})\}) = \frac{\sum_K^M \epsilon_K \phi_K(\mathbf{r})}{\phi_0(\mathbf{r})},$$

ϵ_K : dielectric of particle type K . ϕ_0 : local total particle density.

Potential felt by particle of type K :


$$V_{\text{ext},K}(\mathbf{r}) = q_K \psi(\mathbf{r}) - \frac{1}{2} \frac{\epsilon_K - \epsilon(\mathbf{r})}{\phi_0(\mathbf{r})} |\mathbf{E}(\mathbf{r})|^2,$$

Forces on particle of type K :

$$\mathbf{F}_K = -\nabla V_{\text{ext},K}(\mathbf{r}) = q_K \mathbf{E}(\mathbf{r}) + \frac{1}{2} \nabla \left(\frac{\epsilon_K - \epsilon(\mathbf{r})}{\phi_0(\mathbf{r})} |\mathbf{E}(\mathbf{r})|^2 \right)$$


Force computation and molecular dynamics

Force computation and molecular dynamics



Partitioning of ions (1)

Ions in a phase separated oil/water mixture of ϵ_o and ϵ_w .
($RT \times \chi_{ow} = 30 \text{ kJ mol}^{-1}$)




Partitioning of ions (1)

Ions in a phase separated oil/water mixture of ϵ_o and ϵ_w . ($RT \times \chi_{ow} = 30 \text{ kJ mol}^{-1}$)



Distribution coefficient:

$$D_{o/w} = \frac{c_o}{c_w}$$

c_o and c_w : concentration of ions within each phase.





Partitioning of ions (2)



$$D_{o/w} = f(c, P_{o,w}^\pm, P_{o,w}^{\text{ip}}, K_w)$$

c: concentration of ions.

Partitioning of ions (2)




$$D_{o/w} = f(c, P_{o,w}^\pm, P_{o,w}^{\text{ip}}, K_w)$$

c: concentration of ions.

Born theory of ions:

$$\log P_{o/w}^\pm = \gamma \left(\frac{1}{\epsilon_w} - \frac{1}{\epsilon_o} \right)$$



Outlook

- ▶ Implementation
 - ▶ Parallel version
 - ▶ Improve on accuracy

Outlook

- ▶ Implementation
 - ▶ Parallel version
 - ▶ Improve on accuracy
- ▶ Application
 - ▶ Antonio De Nicola: Charged lipids
 - ▶ Ken Schafer: Molecular packing of SDS
 - ▶ Victoria Ariel Bjørnestad


Outlook

- ▶ Implementation
 - ▶ Parallel version
 - ▶ Improve on accuracy
- ▶ Application
 - ▶ Antonio De Nicola: Charged lipids
 - ▶ Ken Schafer: Molecular packing of SDS
 - ▶ Victoria Ariel Bjørnestad




Outlook

- ▶ Implementation
 - ▶ Parallel version
 - ▶ Improve on accuracy
- ▶ Application
 - ▶ Antonio De Nicola: Charged lipids
 - ▶ Ken Schafer: Molecular packing of SDS
 - ▶ Victoria Ariel Bjørnestad



Outlook

- ▶ Implementation
 - ▶ Parallel version
 - ▶ Improve on accuracy
- ▶ Application
 - ▶ Antonio De Nicola: Charged lipids
 - ▶ Ken Schafer: Molecular packing of SDS
 - ▶ Victoria Ariel Bjørnestad



Acknowledgements

University of Oslo, Norway:

Michele Cascella

Hima Bindu Kolli

Morten Ledum

Victoria Ariel Bjørnestad

Reidar Lund



OCCAM
Molecular Dynamics

The logo for Hylleraas consists of two overlapping circles, one light blue and one light red, followed by the word 'Hylleraas' in a bold, black, sans-serif font.

Yamagata University, Japan:

Giuseppe Milano

Antonio De Nicola

Tsudo Yamanaka

Sendai University, Japan:

Toshihiro Kawakatsu



UiO The logo for the University of Oslo (UiO) consists of the letters 'UiO' in a large, black, sans-serif font, with a red circle to the right.