

Advances in the Hybrid Particle-Field Approach Towards Biological Systems

Sigbjørn Løland Bore

Hylleraas Centre for Quantum Molecular Sciences,
Department of Chemistry, University of Oslo, Norway

PhD lecture, Oslo, 23 April 2020




UiO • **University of Oslo**




Hylleraas


Scope of the work




Scope of the work




Coarse-grained methods



Coarse-grained methods



Coarse-grained methods




$$Z = \int d\Gamma e^{-\beta H(\Gamma)}$$

↓

$$Z \simeq \int d\Gamma_{CG} e^{-\beta H(\Gamma_{CG})}$$


Coarse-grained methods



$$Z = \int d\Gamma e^{-\beta H(\Gamma)}$$

↓

$$Z \simeq \int d\Gamma_{CG} e^{-\beta H(\Gamma_{CG})}$$



Large systems and
long time scales!

The hybrid particle-field method

$$H(\{\mathbf{r}\}) = \sum_{m=1}^{N_{\text{mol}}} \underbrace{H_0(\{\mathbf{r}_m\})}_{\text{Intramolecular}} + \underbrace{W[\{\phi(\mathbf{r})\}]}_{\text{Intermolecular}}$$

($\{\mathbf{r}\} \equiv \{\mathbf{r}_1, \dots, \mathbf{r}_N\}$, *particle positions*)


($\{\phi\} \equiv \{\phi_1, \dots, \phi_M\}$, *particle-type number densities*)

The hybrid particle-field method

$$H(\{\mathbf{r}\}) = \sum_{m=1}^{N_{\text{mol}}} \underbrace{H_0(\{\mathbf{r}_m\})}_{\text{Intramolecular}} + \underbrace{W[\{\phi(\mathbf{r})\}]}_{\text{Intermolecular}}$$

Intermolecular interactions

Particle-particle



($\{\mathbf{r}\} \equiv \{\mathbf{r}_1, \dots, \mathbf{r}_N\}$, *particle positions*)


($\{\phi\} \equiv \{\phi_1, \dots, \phi_M\}$, *particle-type number densities*)

The hybrid particle-field method


$$H(\{\mathbf{r}\}) = \sum_{m=1}^{N_{\text{mol}}} \underbrace{H_0(\{\mathbf{r}_m\})}_{\text{Intramolecular}} + \underbrace{W[\{\phi(\mathbf{r})\}]}_{\text{Intermolecular}}$$

Intermolecular interactions

Particle-particle



Particle-field



($\{\mathbf{r}\} \equiv \{\mathbf{r}_1, \dots, \mathbf{r}_N\}$, *particle positions*)


($\{\phi\} \equiv \{\phi_1, \dots, \phi_M\}$, *particle-type number densities*)

The hybrid particle-field method


$$H(\{\mathbf{r}\}) = \sum_{m=1}^{N_{\text{mol}}} \underbrace{H_0(\{\mathbf{r}_m\})}_{\text{Intramolecular}} + \underbrace{W[\{\phi(\mathbf{r})\}]}_{\text{Intermolecular}}$$

Intermolecular interactions

Particle-particle



Particle-field



$$V_k(\mathbf{r}) = \frac{\delta W[\{\phi\}]}{\delta \phi_k(\mathbf{r})}$$

$$\mathbf{F}_i = -\nabla_i V_k(\mathbf{r}_i)$$

($\{\mathbf{r}\} \equiv \{\mathbf{r}_1, \dots, \mathbf{r}_N\}$, *particle positions*)

($\{\phi\} \equiv \{\phi_1, \dots, \phi_M\}$, *particle-type number densities*)

Interaction energy from polymer theory

$$W[\phi] = \int d\mathbf{r} \frac{1}{\rho_0} \left(\sum_{k\ell} \frac{\tilde{\chi}_{k\ell}}{2} \phi_k(\mathbf{r}) \phi_\ell(\mathbf{r}) + \frac{1}{\kappa} \left(\sum_\ell \phi_\ell(\mathbf{r}) - \rho_0 \right)^2 \right)$$

(ρ_0 : Density parameter related to the volume per bead)

Interaction energy from polymer theory

$$W[\phi] = \int d\mathbf{r} \frac{1}{\rho_0} \left(\underbrace{\sum_{k\ell} \frac{\tilde{\chi}_{k\ell}}{2} \phi_k(\mathbf{r}) \phi_\ell(\mathbf{r})}_{\text{Mixing}} + \frac{1}{\kappa} \left(\sum_\ell \phi_\ell(\mathbf{r}) - \rho_0 \right)^2 \right)$$

$\tilde{\chi}_{k\ell} > 0 \rightarrow$ Likes not to mix

$\tilde{\chi}_{k\ell} \leq 0 \rightarrow$ Likes to mix

(ρ_0 : Density parameter related to the volume per bead)

Interaction energy from polymer theory

$$W[\phi] = \int d\mathbf{r} \frac{1}{\rho_0} \left(\underbrace{\sum_{k\ell} \frac{\tilde{\chi}_{k\ell}}{2} \phi_k(\mathbf{r}) \phi_\ell(\mathbf{r})}_{\text{Mixing}} + \underbrace{\frac{1}{\kappa} \left(\sum_\ell \phi_\ell(\mathbf{r}) - \rho_0 \right)^2}_{\text{Compressibility}} \right)$$

$\tilde{\chi}_{k\ell} > 0 \rightarrow$ Likes not to mix

$\tilde{\chi}_{k\ell} \leq 0 \rightarrow$ Likes to mix

$\kappa \sim 0 \rightarrow$ incompressible

$\kappa \gg 0 \rightarrow$ very compressible

(ρ_0 : Density parameter related to the volume per bead)

Interaction energy from polymer theory

$$W[\phi] = \int d\mathbf{r} \frac{1}{\rho_0} \left(\underbrace{\sum_{k\ell} \frac{\tilde{\chi}_{k\ell}}{2} \phi_k(\mathbf{r}) \phi_\ell(\mathbf{r})}_{\text{Mixing}} + \underbrace{\frac{1}{\kappa} \left(\sum_\ell \phi_\ell(\mathbf{r}) - \rho_0 \right)^2}_{\text{Compressibility}} \right)$$

$\tilde{\chi}_{k\ell} > 0 \rightarrow$ Likes not to mix

$\kappa \sim 0 \rightarrow$ incompressible


$\tilde{\chi}_{k\ell} \leq 0 \rightarrow$ Likes to mix

$\kappa \gg 0 \rightarrow$ very compressible


Net effect $V_k(\mathbf{r}) = \frac{1}{\rho_0} \left(\sum_\ell \tilde{\chi}_{k\ell} \phi_\ell(\mathbf{r}) + \frac{1}{\kappa} \left(\sum_\ell \phi_\ell(\mathbf{r}) - \rho_0 \right) \right)$

(ρ_0 : Density parameter related to the volume per bead)

Prototypic applications: Relaxation of polymer melts



Prototypic applications: Phospholipid aggregation



Research Goals

Overall objective:

- ▶ *Develop new hPF methods and models that can be used in the study of macromolecular biological systems*

Research Goals

Overall objective:

- ▶ *Develop new hPF methods and models that can be used in the study of macromolecular biological systems*

Extend the capability of hPF simulations to representing:

- ▶ Proteins
- ▶ Electrostatics
- ▶ Multiphase electrolytes
- ▶ Constant-pressure simulations

Papers included in the thesis


- I: *Hybrid Particle-Field Model for Conformational Dynamics of Peptide Chains* **Sigbjørn Løland Bore** et al. JCTC **14**, 1120–1130 (2018)
- II: *Hybrid Particle-Field Molecular Dynamics Simulations of Charged Amphiphiles in Aqueous Environment* Hima Bindu Kolli et al. JCTC **14**, 4928–4937 (2018)
- III: *Mesoscale Electrostatics Driving Particle Dynamics in Nonhomogeneous Dielectrics* **Sigbjørn Løland Bore** et al. JCTC **15**, 2033-2041 (2019)
- IV: *Aggregation of Lipid A Variants: a Hybrid Particle-Field Model* Antonio De Nicola et al. BBA, in press (2020)
- V: *Beyond the Molecular Packing Model: Understanding Morphological Transitions of Charged Surfactant Micelles* Ken Schäfer et al. (Submitted for peer-review)
- VI: *Hybrid Particle-Field Molecular Dynamics Under Constant Pressure* **Sigbjørn Løland Bore** et al. JCP, in press (2020)

Papers included in the thesis


- I: *Hybrid Particle-Field Model for Conformational Dynamics of Peptide Chains* **Sigbjørn Løland Bore** et al. JCTC **14**, 1120–1130 (2018)
- II: *Hybrid Particle-Field Molecular Dynamics Simulations of Charged Amphiphiles in Aqueous Environment* Hima Bindu Kolli et al. JCTC **14**, 4928–4937 (2018)
- III: *Mesoscale Electrostatics Driving Particle Dynamics in Nonhomogeneous Dielectrics* **Sigbjørn Løland Bore** et al. JCTC **15**, 2033-2041 (2019)
- IV: *Aggregation of Lipid A Variants: a Hybrid Particle-Field Model* Antonio De Nicola et al. BBA, in press (2020)
- V: *Beyond the Molecular Packing Model: Understanding Morphological Transitions of Charged Surfactant Micelles* Ken Schäfer et al. (Submitted for peer-review)
- VI: *Hybrid Particle-Field Molecular Dynamics Under Constant Pressure* **Sigbjørn Løland Bore** et al. JCP, in press (2020)

Proteins, Electrostatics, Multiphase electrolytes, Constant-pressure


Hybrid Particle-Field Model for Proteins



Intramolecular model: H_0




Intramolecular model: H_0



$$V(\gamma, \phi) = \frac{1}{2}k(\gamma - \gamma_0(\phi))^2 + V_{\text{propensity}}(\phi, \lambda)$$


Intramolecular model: H_0




$$V(\gamma, \phi) = \frac{1}{2}k(\gamma - \gamma_0(\phi))^2 + V_{\text{propensity}}(\phi, \lambda)$$

$$\lambda = -1$$

Helical



Intramolecular model: H_0





$$V(\gamma, \phi) = \frac{1}{2}k(\gamma - \gamma_0(\phi))^2 + V_{\text{propensity}}(\phi, \lambda)$$

$$\lambda = -1$$


Helical

$$\lambda = 1$$

Extended




Intramolecular model: H_0



$$V(\gamma, \phi) = \frac{1}{2}k(\gamma - \gamma_0(\phi))^2 + V_{\text{propensity}}(\phi, \lambda)$$


$$\lambda = -1$$

Helical



$$\lambda = 1$$

Extended




$$\lambda = 0$$

Random



Intermolecular model: W

Hydrophobic



Polar


$4 \cdot \text{H}_2\text{O}$



$$V_k(\mathbf{r}) = \frac{1}{\rho_0} \left(\sum_{\ell} \tilde{\chi}_{k\ell} \phi_{\ell}(\mathbf{r}) + \frac{1}{\kappa} \left(\sum_{\ell} \phi_{\ell}(\mathbf{r}) - \rho_0 \right) \right)$$


Intermolecular model: W

Hydrophobic




$$V_k(\mathbf{r}) = \frac{1}{\rho_0} \left(\sum_{\ell} \tilde{\chi}_{k\ell} \phi_{\ell}(\mathbf{r}) + \frac{1}{\kappa} \left(\sum_{\ell} \phi_{\ell}(\mathbf{r}) - \rho_0 \right) \right)$$

Phase-diagram: Solvated homo-poly-peptide






Phase-diagram: Solvated homo-poly-peptide

$\tilde{\chi}_{kl}$	CB	$4 \cdot H_2O$
CB	0	α
$4 \cdot H_2O$	α	0



Phase-diagram: Solvated homo-poly-peptide




$\tilde{\chi}_{k\ell}$	CB	$4 \cdot H_2O$
CB	0	α
$4 \cdot H_2O$	α	0



i: Random coil ii: α -helix iii: β -hairpin iv: Extended

Phase-diagram: Solvated homo-poly-peptide




$\tilde{\chi}_{k\ell}$	CB	$4 \cdot H_2O$
CB	0	α
$4 \cdot H_2O$	α	0



i: Random coil ii: α -helix iii: β -hairpin iv: Extended v: Helix-coil-helix vi: β -floor/helix


Phase-diagram: Solvated homo-poly-peptide

$\tilde{\chi}_{k\ell}$	CB	$4 \cdot H_2O$
CB	0	α
$4 \cdot H_2O$	α	0



Electrostatics Coarse-Grained Simulations


All-atom molecular dynamics:



Electrostatics Coarse-Grained Simulations

All-atom molecular dynamics:

- ▶ *Charges are resolved*
- ▶ *Screening is modeled directly*




Electrostatics Coarse-Grained Simulations

All-atom molecular dynamics:

- ▶ *Charges are resolved*
- ▶ *Screening is modeled directly*

Coarse-grained molecular dynamics:

- ▶ *Charge resolution is lost*
- ▶ *Dielectric screening modeled modelled indirectly*




Electrostatics Coarse-Grained Simulations

All-atom molecular dynamics:

- ▶ *Charges are resolved*
- ▶ *Screening is modeled directly*

Coarse-grained molecular dynamics:

- ▶ *Charge resolution is lost*
- ▶ *Dielectric screening modeled modelled indirectly*



How do we model electrostatics?

Constant dielectrics approach

Model of paper II:

$$\epsilon \nabla^2 \psi = -\rho(\mathbf{r})$$



External potential:

$$V_K(\mathbf{r}) = q_K(\psi_S(\mathbf{r}) + \psi_L(\mathbf{r}))$$

Solution for ψ :

Particle-Mesh-Ewald by FFT


hPF-MD + electrostatics



Y.L Zhu et al., *Phys. Chem. Chem. Phys.* **18**, 9799–9808 (2016)
H.B. Kolli et al., *J. Chem. Theory Comput.* **14**, 4928–4937 (2018)


Application: Hybrid particle-field model for lipid A

CG-Representation



Application: Hybrid particle-field model for lipid A

CG-Representation




Free parameters:

$$\epsilon \nabla^2 \psi = -\rho(\mathbf{r})$$

$$V_k(\mathbf{r}) = \frac{1}{\rho_0} \sum_{\ell} \left(\tilde{x}_{k\ell} + \frac{1}{\kappa} \right) \phi_{\ell}(\mathbf{r})$$

Application: Hybrid particle-field model for lipid A

CG-Representation




Free parameters:

$$\epsilon \nabla^2 \psi = -\rho(\mathbf{r})$$






$$V_k(\mathbf{r}) = \frac{1}{\rho_0} \sum_{\ell} \left(\tilde{\chi}_{k\ell} + \frac{1}{\kappa} \right) \phi_{\ell}(\mathbf{r})$$

Parameterization

all-atom



$$\epsilon_r = 15$$



Tuning of ϵ_r and χ

Lipid A aggregation

Variable dielectrics approach

Model of paper III:


$$\nabla \cdot (\epsilon(\mathbf{r}) \nabla \psi(\mathbf{r})) = -\rho(\mathbf{r})$$

Solution for ψ :

Successive Over-Relaxation


External potential:

$$V_K(\mathbf{r}) = q_K \psi(\mathbf{r}) - \frac{1}{2} \frac{\partial \epsilon(r)}{\partial \phi_k} |\nabla \psi(\mathbf{r})|^2$$



Application in paper III: Partitioning of ions (1)

Ions in a phase separated oil/water mixture of ϵ_o and ϵ_w . ($RT \times \chi_{ow} = 30 \text{ kJ mol}^{-1}$)




Application in paper III: Partitioning of ions (1)

Ions in a phase separated oil/water mixture of ϵ_o and ϵ_w . ($RT \times \chi_{ow} = 30 \text{ kJ mol}^{-1}$)



Distribution coefficient:

$$D_{o/w} = \frac{c_o}{c_w}$$

(c_o and c_w : concentration of ions within each phase)





Application in paper III: Partitioning of ions (2)



$$D_{o/w} = f(c, P_{o,w}^\pm, P_{o,w}^{\text{ip}}, K_w)$$

c : concentration of ions.

Application in paper III: Partitioning of ions (2)




$$D_{o/w} = f(c, P_{o,w}^\pm, P_{o,w}^{\text{ip}}, K_w)$$


c: concentration of ions.

Born theory of ions:

$$\log P_{o/w}^\pm = \gamma \left(\frac{1}{\epsilon_w} - \frac{1}{\epsilon_o} \right)$$



Constant-Pressure Simulations



Pressure for particle-field

Hybrid particle field interaction-energy:

$$W_0[\phi] = \int d\mathbf{r} \frac{1}{\rho_0} \left(\sum_{k\ell} \frac{\tilde{\chi}_{k\ell}}{2} \phi_k(\mathbf{r}) \phi_\ell(\mathbf{r}) + \frac{1}{\kappa} \left(\sum_\ell \phi_\ell(\mathbf{r}) - a \right)^2 \right)$$

(a : equation of state parameter.)

Pressure for particle-field

Hybrid particle field interaction-energy:

$$W_0[\phi] = \int d\mathbf{r} \frac{1}{\rho_0} \left(\sum_{k\ell} \frac{\tilde{\chi}_{k\ell}}{2} \phi_k(\mathbf{r}) \phi_\ell(\mathbf{r}) + \frac{1}{\kappa} \left(\sum_\ell \phi_\ell(\mathbf{r}) - a \right)^2 \right)$$

We use $P_\mu = -\frac{1}{V} \left\langle L_\mu \frac{\partial W}{\partial L_\mu} \right\rangle_{T,N}$:

(a : equation of state parameter.)

Pressure for particle-field

Hybrid particle field interaction-energy:





$$W_0[\phi] = \int d\mathbf{r} \frac{1}{\rho_0} \left(\sum_{k\ell} \frac{\tilde{\chi}_{k\ell}}{2} \phi_k(\mathbf{r}) \phi_\ell(\mathbf{r}) + \frac{1}{\kappa} \left(\sum_\ell \phi_\ell(\mathbf{r}) - a \right)^2 \right)$$

We use $P_\mu = -\frac{1}{V} \left\langle L_\mu \frac{\partial W}{\partial L_\mu} \right\rangle_{T,N}$:

$$P_{0,\mu} = \frac{1}{V} \int d\mathbf{r} \frac{1}{\rho_0} \left(\sum_{k\ell} \frac{\tilde{\chi}_{k\ell}}{2} \phi_k(\mathbf{r}) \phi_\ell(\mathbf{r}) + \frac{1}{2\kappa} \left(\phi(\mathbf{r})^2 - a^2 \right) \right),$$

(a : equation of state parameter.)

Parameterization of water model



Conclusions and outlook

New methodology for:

- ▶ Electrostatics
- ▶ Constant-pressure simulations

Conclusions and outlook

New methodology for:

- ▶ Electrostatics
- ▶ Constant-pressure simulations

New models for:

- ▶ Proteins
- ▶ Charged lipids

Conclusions and outlook

New methodology for:

- ▶ Electrostatics
- ▶ Constant-pressure simulations

New models for:

- ▶ Proteins
- ▶ Charged lipids

Outlook:

- ▶ Chemical specific models
- ▶ Parameterization of models
- ▶ OpenMP parallelization

Conclusions and outlook

New methodology for:

- ▶ Electrostatics
- ▶ Constant-pressure simulations

New models for:

- ▶ Proteins
- ▶ Charged lipids

Outlook:

- ▶ Chemical specific models
 - ▶ Parameterization of models
 - ▶ OpenMP parallelization
- ⇒ Realistic representation of large biological systems

Acknowledgements

University of Oslo, Norway:

Michele Cascella

Hima Bindu Kolli

Morten Ledum

Simen Kvaal

Victoria Ariel Bjørnestad

Reidar Lund



OCCAM
Molecular Dynamics

The logo for Hylleraas consists of two overlapping circles, one light blue and one light red, followed by the word "Hylleraas" in a bold, black, sans-serif font.

Yamagata University, Japan:

Giuseppe Milano

Antonio De Nicola

Tsudo Yamanaka

Sendai University, Japan:

Toshihiro Kawakatsu

The logo for notur features a circular pattern of vertical bars of varying heights, resembling a stylized globe or a bar chart, positioned to the left of the word "notur" in a large, bold, dark blue sans-serif font.



The logo for the University of Oslo (UiO) consists of the letters "UiO" in a large, bold, black sans-serif font, with a red dot positioned to the right of the letter "O".