

Quantum Maimonides

Charitable Giving in Anonymous Sub-Atomic Networks

Moses ben Maimon

Department of Natural Philosophy

Universidad de Córdoba

Cordova, Andalusia, Reino de España

mm@uco.es

ABSTRACT

Maimonides' Ladder of Tzedakah is a 12th Century guidance from *Laws of Obligatory Gifts to the Poor*, in the 7th book of the Mishnah Torah - Zera'im (Seeds) - in the second section titled "Aniyim." In it, he prioritizes alms (or giving) in terms of "charitability" over "random" charity. Nonetheless, acts of blind charity are essential, especially in absence of conjunctural hardships from proximate contacts (such as relatives or friends) in well-to-do networks. This paper will explore a "subterranean-like" network of charitable acts and donations that use quantum tunneling effects along existing quantum field theory (QFT) and putative superstring theory, with a comparative analysis of quantum currencies: Neuromancer/Neuralink, cryptowallets, and luminiferous aether.

KEYWORDS

Quantum finance, Philanthropy, Quantum Field Theory, Majorana particles

Introduction

Alms giving is a millennium old practice, spanning over 35 centuries from the time of the Torah (Deuteronomy 15:7-11), to the Vedic period (1500 BCE), to Buddha in the 5th Century BCE¹, to the Apostolic era in the 1st Century, and Islam in the 6th Century. While alms giving is only known to the extent of written history, its practice extends probably much earlier than the beginning of recorded history.

Modern philanthropy has evolved significantly since its earliest origins. Mite boxes were used in the time of Jehoiada². Tithes were used in various religions. Maimonides' *Laws of Gifts to the Poor* was written after considerable family hardship in 1177, and formed the basis of what can be called reasoned, or subjective giving³. Further advancements in economic systems allowed for tax-deductible donations, often to 501(c)(3) organizations. The advent of quantum computing has opened new avenues for philanthropy, allowing for anonymous donations to be administered via secure quantum transport layers, using

fermionic particles serving as error correction codes (ECC) [Figure 1]^{4,5,6,7}.




Figure 1: A Majorana fermion (Palacio-Morales et al. Science Advances, 2019)⁴

Theorem

A secure quantum donation network cannot exist without an accurate mechanism to safeguard from quantum transmission errors, therefore its infrastructure and transport layer certificates must be fully verified before deployment.

1.1 Maimonides' Quantum OSI Layer

The 8 Levels of Giving in the Ladder of Tzedakah mirrors the OSI Internet X.200 architecture⁸, with the exception of the highest layer (Enabling self-sufficiency: loans/business partnerships) being excluded due to it not needing quantum encryption (often done in person, or via established intermediaries). As Table 1 shows, only Layers 5-7 are capable of 2D anyonic braiding, which are necessary for secure and anonymous donations, similar to slipping an envelope with cash under a door, regardless of direction. Fermions and Bosons are suggested to exist in both 1 dimensional and 3 dimensional planes, while the dimensionless applications of neutrinos are believed to exist in supersymmetric space. All layers, regardless of whether they are anonymous, are considered untamperable and satisfying the measurement problem¹⁶. However, in the anonymous categories,

limits to the maximum donations can be set by the benefactor and/or the beneficiary. This is to prevent overwithdrawing from a public coffer intended for multiple recipients (as in Level 5). A unique, but anonymous wallet ID, similar to a virtual debit card number, can be used by a recipient so that the Unknown donor does not deposit more than their maximum donation amount. This differs from “giving inadequately” (Level 2), as the benefactor may have intended to donate a large sum to a number of recipients, rather than insufficiently to a number of the same subjects.

Table 1: OSI Transport Layers with Quantum ECC Braids

Layer		Unit	Function
Anonymous giving	7	Anonymous to Unknown	Anyons (2D) ^{9, 10}
	6	Known to Anonymous	
	5	Unknown to Known	
	4	Giving Unasked	Fermion (1/3D) ¹¹ Boson (1/3D) ^{12, 13}
Overt giving	3	Giving After Asking	Fermion (1/3D) Boson (1/3D)
	2	Giving Inadequately	Fermion (1/3D) Boson (1/3D)
	1	Giving Reluctantly	Neutrino (?D) ^{14,15}

1.2 Comparative Analysis of Digital Currencies

In recent decades, cryptocurrencies such as Bitcoin and Ethereum have attempted to secure — or at least make transparent — digital transactions with a public ledger, or, in Monero’s case, an anonymous ledger. The limits to bitcoin value are thought to be energy utility investments (mining) and policing currency exchanges, whereas the limits of a modern quantum currency system depend only on the infrastructure investment for its adoption and maintenance. All technology eventually becomes obsolete, therefore the cost of manufacturing quantum wallets and quantum banking apps do depend on a modern semiconductor manufacturing supply chain. However, as costs become externalized to startups such as Nanotechnology giant Neuralink, alternative options to trading securities can bypass the brain stem via consensual backdoors. Opponents to Neuralink such as #NoBackdoorsForMyBrainStem may opt for

natural public spaces for exchanging securities, using luminiferous aether, tech yet to be commercialized^{17,18,19}.

ACKNOWLEDGMENTS

Acknowledgments are made only to my anonymous supporter(s). Thank you. You know who you are :) 😊

REFERENCES

- [1] https://en.wikipedia.org/wiki/Book_of_Deuteronomy
- [2] https://en.wikipedia.org/wiki/Poor_box#cite_note-Carrera1828-1
- [3] <https://sapirjournal.org/money/2025/jewish-masterpiece-laws-of-gifts-to-the-poor/>
- [4] <https://www.sciencealert.com/look-scientists-have-imaged-a-majorana-fermion>
- [5] <https://arxiv.org/abs/0707.1889> Nayak, Chetan; Simon, Steven H.; Stern, Ady; Freedman, Michael; Sarma, Sankar Das (27 March 2008). "Non-Abelian anyons and topological quantum computation". *Reviews of Modern Physics*. **80** (3): 1083–1159
- [6] <https://arxiv.org/abs/1501.02813> Sarma, Sankar Das; Freedman, Michael; Nayak, Chetan (27 October 2015). "Majorana zero modes and topological quantum computation" (1): 15001.
- [7] <https://arxiv.org/abs/1408.3379> Hastings, M.B.; Geller, A. (2015). "Reduced space-time and time costs using dislocation codes and arbitrary ancillas". *Quantum Information and Computation*. **15** (11–12): 0962–0986.
- [8] https://en.wikipedia.org/wiki/OSI_model#Layer_architecture
- [9] <https://phys.org/news/2020-07-evidence-anyons.html>
- [10] <https://arxiv.org/abs/quant-ph/9707021> A. Yu. Kitaev, "Fault-tolerant quantum computation by anyons" Jul 1997
- [11] <https://arxiv.org/abs/2401.09549> "Interferometric Single-Shot Parity Measurement in an InAs-Al Hybrid Device" *Nature* 638, 651 (2025)
- [12] <https://www.pnas.org/doi/10.1073/pnas.2304294120> "Fermionic quantum processing with programmable neutral atom arrays" 2023
- [13] <https://www.nature.com/articles/s41534-025-01001-4> "Digital-analog quantum computing of fermion-boson models in superconducting circuits" *npj Quantum Information* volume11 (2025)
- [14] https://en.wikipedia.org/wiki/Neutrino#Scientific_interest
- [15] <https://web.archive.org/web/20170329063317/https://www.scientificamerican.com/article/dimensional-shortcuts/>
- [16] https://en.wikipedia.org/wiki/Masurement_problem
- [17] <https://philsci-archive.pitt.edu/4000/1/LT.pdf> "Relativity without tears", Z. K. Silagadze, 2008
- [18] <https://www.newscientist.com/article/mg24432543-300-einstein-killed-the-aether-now-the-idea-is-back-to-save-relativity/>
- [19] <https://ui.adsabs.harvard.edu/abs/2024PhyEs..37....9L/abstract> <https://vixra.org/pdf/2401.0050v1.pdf> "The quantum luminiferous aether," 2024