

The Heart

1

2.1 Normal Heart (brief review)

2.2 Principles of cardiac dysfunction

2.2.1 Heart failure

2.2.1.1 Cardiac hypertrophy:
Pathophysiology & progression to failure

2.2.1.2 Left-sided heart failure

2.2.1.3 Right-sided heart failure

• 2.3 Types of heart disease

✦ **Congenital Heart disease**

- Left –to- right shunts – Late cyanosis
 - ASD, VSD, PDA, atrioventricular septal defect
- Right – to – left shunts – Early cyanosis
 - Tetralogy of Fallot, transposition of great arteries, truncus arteriosus, tricuspid atresia, total anomalous pulmonary venous connection
- Obstructive congenital anomalies
 - coarctation of aorta, pulmonary stenosis & atresia, aortic stenosis & atresia.

✦ **Ischemic heart disease**

- Angina pectoris
- Myocardial infarction
- Chronic ischemic heart disease
- Sudden cardiac death

✦ **Hypertensive heart disease**

- Systemic (left-sided) hypertensive heart disease
- Pulmonary (right-sided) hypertensive heart disease (cor pulmonale)

✦ **Valvular heart disease**

- Rheumatic fever & rheumatic heart disease
- Infective endocarditis
- Noninfected vegetations
- Carcinoid heart disease
- Valvular degenerations caused by calcification
- Mitral valve prolapse

✦ **Myocardial disease**


- Cardiomyopathies
- Myocarditis

✦ **Pericardial disease**


- Pericardial effusion & hemopericardium
- Pericarditis
- Rheumatoid heart disease


✦ **Neoplastic Heart disease**

- Primary cardiac tumors (myxoma, lipoma, etc...)





Copyright © 2004 Pearson Education, Inc., publishing as Benjamin Cummings.






Copyright © 2004 Pearson Education, Inc., publishing as Benjamin Cummings.








(b)

Copyright © 2001 Benjamin Cummings, an imprint of Addison Wesley Longman, Inc.




(d)

Copyright © 2001 Benjamin Cummings, an imprint of Addison Wesley Longman, Inc.




Copyright © 2004 Pearson Education, Inc., publishing as Benjamin Cummings.




(e)

Copyright © 2001 Benjamin Cummings, an imprint of Addison Wesley Longman, Inc.








Copyright © 2004 Pearson Education, Inc., publishing as Benjamin Cummings.



Copyright © 2004 Pearson Education, Inc., publishing as Benjamin Cummings.







- Cardiovascular dysfunction results from one or more of five principal mechanisms


- Failure of the pump
- An obstruction to flow
- Regurgitant flow
- Disorders of cardiac conduction
- Disruption of the continuity of the circulatory system

CONGESTIVE HEART FAILURE

- Is a condition when cardiac output insufficient for metabolic requirements of the body
- Heart failure is the end result of many different forms of heart disease
- CHF is a common & recurrent condition with a poor prognosis

- Heart failure is usually consequence of progressive deterioration of myocardial contractile function (**systolic dysfunction**) as often occurs in ischemic injury & pressure or volume overload (as in hypertension)

- Heart failure can also result from an inability of the heart chamber to relax, expand & fill sufficiently during diastole to accommodate an adequate ventricular blood volume (**diastolic dysfunction**)
Eg- massive left ventricular hypertrophy , myocardial fibrosis or constrictive pericarditis



- Heart failure may result because of greatly increased demands for blood by the tissues, a process sometimes referred to as *high output failure*.

- Inadequate cardiac output, also termed *forward failure*, is almost always accompanied by increased congestion of the venous circulation (*backward failure*), because the failing ventricle is unable to eject the normal volume of venous blood delivered to it during diastole.

- Congestive heart failure may involve the left or right side of the heart or all of the cardiac chambers

Pathophysiology & Progression To Failure

- As the heart begins to fail, a number of local *adaptive responses* are triggered in an attempt to maintain normal cardiac output.
- These include *neurohumoral reactions* as well as *molecular and morphologic changes* within the heart.

➤ ***Activation of neurohumoral systems***

- Release of the norepinephrine by adrenergic cardiac nerves (which increase heart rate & augments myocardial contractility & vascular resistance)
- Activation of the renin-angiotensin-aldosterone system
- Release of atrial natriuretic peptide

➤ ***The Frank-Starling mechanism***

- The increased preload of dilation helps to sustain cardiac performance by enhancing contractility.

- ***Myocardial structural changes, including augmented muscle mass(hypertrophy) with or without cardiac chamber dilation, in which the mass of contractile tissue is augmented.***

- Greater heart weight or ventricular thickness indicates hypertrophy & an enlarged chamber size implies dilatation .

Cardiac Hypertrophy: Pathophysiology & Progression To failure

33


- The major initial structural adaptation to a chronically increased workload is hypertrophy of individual muscle fibers.

- Increased mechanical work due to pressure or volume overload or trophic signals (eg hyperthyroidism through stimulation of beta-adrenergic receptors) increases rate of protein synthesis & lead to an increase in dimension & mass of myocyte & consequently size of the heart.

- In the case of a cardiac chamber subjected to a pure pressure load (e.g., hypertension, valvular stenosis), the hypertrophy is characterized by an increase in the diameter of individual muscle fibers. Cross-sectional area of myocytes is expanded but not cell length

This pattern of fiber enlargement results in the development of classic *concentric hypertrophy*, in which the thickness of the ventricular wall increases without an increase in the size of the chamber

- When the heart is subjected to an *abnormal volume load*, as opposed to a pressure load (e.g., valvular regurgitation or abnormal shunts), the length of individual muscle fibers also increases. This pattern of hypertrophy, sometimes referred to as *eccentric hypertrophy*, is characterized by an increase in heart size as well as an increase in wall thickness.



- The development of hypertrophy initially serves as a positive, adaptive response. However, sustained cardiac hypertrophy often evolves to cardiac failure.

- Despite the potential hemodynamic benefits associated with hypertrophy, the development of hypertrophy comes at a significant cost to the cell. Oxygen requirements of the hypertrophic myocardium are increased, owing to increased myocardial cell mass and increased tension of the ventricular wall accompanied by decreased capillary density.

- Additional evidence suggests that molecular signals that lead to the development of hypertrophy may also be accompanied by the expression of certain proteins that lead to impaired myocyte contractility, increased fibrous tissue and even to premature myocyte death.

HYPERTENSION

Pressure
overload

VALVULAR
DISEASE

Pressure and/or
volume overload

MYOCARDIAL
INFARCTION

Regional dysfunction
with volume overload

↑ Cardiac work

↑ Wall stress

Cell stretch

Hypertrophy and/or dilation

- Characterized by
- ↑ heart size and mass
 - ↑ protein synthesis
 - induction of immediate-early genes
 - induction of fetal gene program
 - abnormal proteins
 - fibrosis
 - inadequate vasculature

Cardiac dysfunction

- Characterized by
- heart failure (systolic/diastolic)
 - arrhythmias
 - neurohumoral stimulation

Left-Sided Heart Failure

- *The most common causes of left-sided cardiac failure are systemic hypertension, mitral or aortic valve disease, ischemic heart disease, and primary diseases of the myocardium*

- The morphologic & clinical effects of left-sided CHF primarily result from progressive damming of blood within the pulmonary circulation & the consequences of diminished peripheral blood perfusion & flow.

Morphology

Heart


- ▣ Depending on the cause of the disease process, abnormalities such as myocardial infarction or valvular deformity may be present.
- ▣ The failing **left ventricle is dilated** and is usually hypertrophied as well

- Secondary enlargement of the left atrium with resultant atrial fibrillation may compromise stroke volume or cause blood stasis & possible thrombus formation


Lungs

- Pressure in the pulmonary veins is transmitted retrograde to the capillaries & arteries. The result is pulmonary congestion & edema

- The **lungs** are boggy and congested and the cut surface exudes a frothy mixture of surfactant-rich fluid and blood.
- Microscopically, the pulmonary alveolar capillaries are congested. There is transudation of fluid, initially limited to perivascular interstitial spaces, causing widening of the alveolar septa. In time, it overflows into the alveoli (**pulmonary edema**).
- The protein-poor edema fluid stains pale pink when viewed under the microscope.



- With persistent elevation of pulmonary venous pressure, the capillaries may become tortuous and may rupture to produce small hemorrhages into the alveolar spaces. Alveolar macrophages phagocytose red blood cells and eventually become filled with hemosiderin. These pigmented macrophages are called **heart failure cells**.



- Persistence of septal edema may induce fibrosis within the alveolar walls, which, together with the accumulation of hemosiderin, is characteristic of **chronic venous congestion** of lungs. The lungs thus become dark brown and firm, an appearance that has been referred to as **brown induration of the lungs**.

Kidneys

- Decreased cardiac output causes a reduction in renal perfusion, which activates the renin-angiotensin-aldosterone system, inducing retention of water & salt with consequent expansion of interstitial fluid & blood volume.
- If the perfusion deficit of the kidney is severe, impaired excretion of nitrogenous products may cause azotemia (prerenal)

Brain

- Cerebral hypoxia may give rise hypoxic encephalopathy

Right- Sided Heart Failure


- ☐ *The most common cause of right-sided heart failure is left ventricular failure, with its associated pulmonary congestion and elevation in pulmonary arterial pressure.*


- Right-sided failure may also occur in the absence of left-sided heart failure in patients with intrinsic diseases of the lung parenchyma and/or pulmonary vasculature (cor pulmonale) and in patients with pulmonic or tricuspid valve disease. It sometimes follows congenital heart diseases, in which there is a left-to-right shunt.

Morphology

Liver & portal system

- ❑ Liver is increased in size & weight (congestive hepatomegaly)
- ❑ Changes in the **liver** include chronic passive congestion, characterized by atrophy of hepatocytes around the central veins that gives rise to a nutmeg-like appearance on the cut surface of the organ .
- ❑ Hemorrhagic necrosis of centrilobular hepatocytes is common in severe cases, particularly in patients with concomitant left ventricular failure. With long-standing cardiac failure, the liver may become fibrotic and, in extreme cases, frankly cirrhotic (cardiac cirrhosis).





Spleen

- ☐ Right sided heart failure leads to elevated pressure in the portal vein & its tributaries.
- ☐ Congestion results in a tense, enlarged spleen (congestive splenomegaly)

Subcutaneous Tissues

- Peripheral edema of dependent portions of the body (pedal & pretibial edema) is the hallmark of right-sided heart failure.
- Generalized massive edema is called *anasarca*.

Clinical features

Left-sided heart failure

- The most common & earliest manifestation of left ventricular failure is *Dyspnea*, or a sense of breathlessness.
- This is caused predominantly by decreased lung compliance resulting from pulmonary edema and congestion and Dyspnea is most noticeable during periods of physical activity (*exertional dyspnea*). It is also prominent when the person is lying down (*orthopnea*) because of the increased amount of venous blood returned to the thorax from the lower extremities and because the diaphragm is elevated in this position.

- *Paroxysmal nocturnal dyspnea* is a dramatic form of dyspnea that awakens the patient with sudden, severe shortness of breath, accompanied by coughing, a choking sensation, and wheezing.

Right sided heart failure

- ▣ Its major consequences are *systemic venous congestion* and *soft tissue edema*.
- ▣ Systemic venous congestion is manifested clinically by distended neck veins and an enlarged liver & spleen. It is also associated with an increased frequency of deep venous thrombi and pulmonary embolism.
- ▣ Edema causes weight gain and usually becomes apparent first in the dependent areas of the body, such as the feet and lower legs. With more severe degrees of ventricular failure, the edema may become generalized.
- ▣ Pleural effusions are common, particularly on the right side, and may be accompanied by pericardial effusions and ascites.

- In many cases of chronic CHF, patients present with the clinical syndromes of both right-sided & left-sided heart failure.

Heart disease

☐ Five categories of heart disease include

- ✓ ***Congenital heart disease***
- ✓ ***Ischemic heart disease***
- ✓ ***Hypertensive heart disease***
- ✓ ***Valvular heart disease***
- ✓ ***Nonischemic (primary) myocardial disease***

Congenital Heart Disease

- Abnormalities of the heart or great vessels that are present from birth.
- Congenital abnormalities of the heart occur in about 8 per 1000 live births, making them one of the most common types of congenital malformations

- Most disorders arise from faulty emryogenesis during gestational weeks 3 through 8, when major cardiovascular structures develop.

- Some forms produce manifestation soon after birth. Others don't necessarily become evident until adulthood.

Etiology & Pathogenesis

- ***Genetic Factors***

- Trisomy 21 (associated with Down syndrome) is the most common known genetic cause of congenital heart disease.

- ***Environmental factors***
Congenital rubella infection
Teratogens

- Multifactorial genetic, environmental & maternal factors account for majority of cases (90%) in which cause is unknown.

- Congenital heart diseases can be subdivided into three major groups
 - Malformations causing a *left-to-right shunt*
 - Malformations causing a *right-to-left shunt* (cyanotic congenital heart diseases)
 - Malformations causing *obstruction*

RELATIVE FREQUENCIES OF CARDIAC MALFORMATIONS AT BIRTH

73

Malformation	Congenital Heart Disease (%)
Ventricular septal defect	42
Atrial septal defect	10
Pulmonary stenosis	8
Patent ductus arteriosus	7
Tetralogy of Fallot	5
Coarctation of the aorta	5
Atrioventricular septal defect	4
Aortic stenosis	4
Transposition of the great arteries	4
Truncus arteriosus	1

Left-to-Right Shunts

- Left-to-right shunts, or abnormal communications permitting blood to flow from the left to the right cardiac chambers, represent the most common type of congenital cardiac malformation.
- They include *atrial septal defects (ASDs)*, *ventricular septal defects (VSDs)*, and *patent (persistent) ductus arteriosus (PDA)*.

- These malformations may be asymptomatic at birth, or they may cause fulminant congestive heart failure.


- Left-to-right shunts increase pulmonary blood flow. when pulmonary circulation is exposed to increased pressure and volume , it can result in right ventricular hypertrophy and, potentially, failure.

- Pulmonary arteries respond to increased pressure and volume by medial hypertrophy and vasoconstriction.
- Prolonged pulmonary arterial vasoconstriction stimulates development of irreversible obstructive intimal lesions
- Eventually pulmonary vascular resistance increases toward systemic levels, reversing the shunt to right-to-left with unoxygenated blood in the systemic circulation *late cyanotic congenital heart disease or Eisenmenger syndrome.*

- Once significant pulmonary hypertension develops, the structural defects of congenital heart disease are considered irreparable.

ATRIAL SEPTAL DEFECTS


- **ASD** is an abnormal opening in the atrial septum permitting free communication of blood between the left and right atria..



There are three types


The *ostium secundum ASD*

- ▣ The most common (90% of cases)
- ▣ appears as a smooth-walled defect in the vicinity of the foramen ovale



82

ATRIAL SEPTAL DEFECT Ostium Secundum



Ostium primum ASDs are


- less common (5% of cases)
- they occur adjacent to the AV valves .
- This defect is often associated with abnormalities in other structures derived from the endocardial cushions, such as the mitral and tricuspid valves.

Sinus venosus ASD

- ▢ the least common
- ▢ **Sinus venosus ASDs** are located high in the atrial septum and are often accompanied by anomalous drainage of the pulmonary veins into the right atrium or superior vena cava.

ATRIAL SEPTAL DEFECT

Sinus Venosus




Clinical Features

- ASDs are the most common congenital cardiac malformations first diagnosed in adults
- ▣ ASDs cause left-to-right shunts after birth, owing to the lower pressures in the pulmonary circulation and right side of the heart.
- ▣ Ostium secundum defects are the most common, and in general these defects are well tolerated, especially if they are less than 1 cm in diameter, but even larger lesions do not produce any symptoms in childhood because the flow of blood is from left to right.

- With time, pulmonary vascular resistance increases, culminating in the development of pulmonary hypertension in some patients. This, in turn, causes reversal of the left-to-right shunt, manifested by the development of cyanosis and congestive heart failure.
- Ostium primum defects may be initially asymptomatic, but they are more likely to be associated with evidence of congestive heart failure, in part because of the high frequency of associated mitral insufficiency

VENTRICULAR SEPTAL DEFECTS

- an abnormal opening in the ventricular septum, which allows free communication between the Rt & Lt ventricles. Accounts for 25% of CHD



- *VSDs are the most common congenital heart defects at birth*
- Like ASDs, they may occur in isolation (around 30% of cases), but they are usually encountered in association with other cardiac malformations

Types

- Perimembranous (or membranous) – Most common.
- Infundibular (subpulmonary or supracristal VSD) – involves the RV outflow tract.
- Muscular VSD – can be single or multiple.

MORPHOLOGY


- The size and location of VSDs are variable, ranging from minute defects in the muscular or membranous portions of the septum to large defects involving virtually the entire septum .
- In defects associated with a significant left-to-right shunt, the right ventricle is hypertrophied and often dilated. The diameter of the pulmonary artery is increased, owing to the increased volume ejected by the right ventricle. Vascular changes typical of pulmonary hypertension are common.


Clinical Features

- Small VSDs may be asymptomatic, and those in the muscular portion of the septum may close spontaneously during infancy or childhood.
- Larger defects, however, cause a severe left-to-right shunt, often complicated by pulmonary hypertension and congestive heart failure.
- Progressive pulmonary hypertension, with resultant reversal of the shunt and cyanosis, occurs earlier and more frequently in patients with VSDs than in those with ASDs
- A particular risk for those with small or medium-sized defects that produce jet lesions in the right ventricle is superimposed infective endocarditis.

PATENT DUCTUS ARTERIOSUS

- Persistence of the normal fetal vessel that joins the PA to the aorta after birth (ductus arteriosus).
- Isolated PDA accounts for about 7% of cases of congenital heart disease. This lesion may also occur in combination with other anomalies, particularly VSDs





Clinical Features

- ❏ PDA causes a high-pressure left-to-right shunt, audible as a harsh waxing and waning murmur sometimes referred to as a "machinery" murmur.
- ❏ A small PDA causes no symptoms; with larger defects, symptoms develop in childhood or adulthood.
- ❏ As with other left-to-right shunts, the development of pulmonary hypertension is announced by the appearance of cyanosis and congestive heart failure.
- ❏ The high-pressure shunt also predisposes affected individuals to the development of infective endocarditis.

Right-to-left shunts

- When blood from the right side of the heart enters the left side, cyanosis & dusky blueness of the skin and mucus membrane develops because there is diminished pulmonary blood flow and poorly oxygenated blood enters the systemic circulation & cyanotic congenital heart disease.
- The important examples include *tetralogy of Fallot*, *transposition of the great vessels*, *persistent truncus arteriosus*, *tricuspid atresia* and *total anomalous pulmonary venous connection*.

- Bland and septic emboli arising in veins can directly enter the systemic circulation (paradoxical embolism) and result in brain infarction and abscess

- Clinical features associated with severe, long standing cyanosis include clubbing of the tips of the fingers & toes & polycythemia.



Normal angle
of nail bed



Distorted angle
of nail bed

Clubbed fingers




ADAM.




TETRALOGY OF FALLOT


- Accounting for about 5% of all congenital cardiac malformations, tetralogy of Fallot is *the most common cause of cyanotic congenital heart disease*

- The four components of the tetralogy are
 - VSD
 - "dextraposed" aortic root that overrides the VSD
 - Right ventricular outflow obstruction (subpulmonary stenosis)
 - Right ventricular hypertrophy






A Classic Tetralogy of Fallot



With VSD



Without VSD

B Complete Transposition

Clinical Features


- The hemodynamic consequences of tetralogy of Fallot are a right-to-left shunt, decreased blood flow to the lungs, and increased blood flow through the aorta. *The extent of shunting is determined by the degree of right ventricular outflow obstruction*
- Marked stenosis causes significant cyanosis early in life. Thus, most patients with tetralogy of Fallot are cyanotic from birth or soon thereafter

- Because of the right-to-left shunt, patients with tetralogy of Fallot are also at increased risk for infective endocarditis, systemic emboli, and brain abscesses.
- These patients develop complications of chronic cyanosis, such as erythrocytosis with attendant hyperviscosity, and digital clubbing


TRANSPOSITION OF THE GREAT ARTERIES

- Transposition of the great arteries is the second leading cause of cyanotic congenital heart disease.
- Because of abnormal truncal septation, *the aorta arises from the right ventricle and the pulmonary artery arises from the left ventricle.*


- In its complete form, the pulmonary and systemic circulations are entirely separate, and there is no shunting of blood. However, such a condition is incompatible with extrauterine life; therefore, those who survive after birth must have some type of shunt, such as an ASD, VSD, or PDA, that allows oxygenated blood to reach the aorta.



A Classic Tetralogy of Fallot




With VSD



Without VSD

B Complete Transposition



Congenital Obstructive Lesions

- A number of malformations cause obstruction of blood flow. In some cases, these are isolated lesions, as in congenital valvular aortic stenosis. In other instances, they are one component of a more complex malformation


COARCTATION OF THE AORTA


- Coarctation of the aorta, defined as an abnormal narrowing of the aortic lumen, is one of the most important forms of obstructive congenital heart disease.
- Females with Turner syndrome frequently have coarctation of aorta

- Most cases of coarctation of the aorta can be placed into one of two major categories: preductal and postductal. The postductal type is the more common.


Preductal coarctation, formerly referred to as infantile coarctation, is characterized by **narrowing of the so-called aortic isthmus**, the segment of aorta that lies between the left subclavian artery and the point of entry of the ductus arteriosus.

- The ductus arteriosus is usually patent and is the main source of blood delivered to the distal aorta.





In the more common **postductal**, or **adult-type**, **coarctation**, the aorta is constricted by a sharply defined ridge of tissue at, or just distal to, the obliterated ductus arteriosus (the ligamentum arteriosum)






Clinical Features

- ❏ Patients with *preductal coarctation* usually present in infancy.
- ❏ Classic features include congestive heart failure and selective cyanosis of the lower extremities, the latter caused by perfusion of the lower part of the body by poorly oxygenated blood delivered via the ductus arteriosus.
- ❏ The femoral pulses are almost invariably weaker than those of the upper extremities. These patients do not survive the neonatal period without surgical correction.

- *Postductal coarctation*, in contrast, is more likely to present as signs and symptoms in older children and adults..
- *Hypertension of the upper extremities is noted in most cases.* By contrast, blood pressure is low and pulses are weak in the lower extremities.
- Increased collateral circulation through intercostal arteries can result in "rib notching".



Ischemic heart disease

- Ischemic heart disease refers to a group of closely related syndromes resulting from myocardial ischemia caused by an imbalance between the myocardial oxygen demand and the blood supply.

- The most common cause of ischemic heart disease is narrowing of the lumina of the coronary arteries by atherosclerosis, and hence ischemic heart disease is often termed *coronary heart disease* or *coronary artery disease*.

126

☰ Depending on the rate and severity of coronary artery narrowing and the myocardial response, one of four syndromes may develop

(1) Various forms of *angina pectoris* (chest pain)

(2) *Myocardial infarction (MI)*

(3) *Sudden cardiac death*

(4) *Chronic ischemic heart disease with congestive heart failure*

- The term *acute coronary syndromes* is applied to the spectrum of three acute catastrophic manifestations of ischemic heart disease-unstable angina, acute MI, and sudden cardiac death.


Pathogenesis

- The dominant influence in the causation of the IHD syndromes is diminished coronary perfusion relative to myocardial demand , due to complex & dynamic interaction among fixed atherosclerotic narrowing of the epicardial coronary arteries, intraluminal thrombosis overlying disrupted atherosclerotic plaque, platelet aggregation & vasospasm .

- Severe and chronic atherosclerosis that causes narrowing of the lumen of one or more coronary arteries is the fundamental disorder underlying ischemic heart disease.
- A fixed 75% or greater reduction in the lumen of the coronary artery is defined as "critical stenosis".
- The onset of symptoms and the prognosis of ischemic heart disease, however, depend not only on the extent and severity of fixed, chronic anatomic disease but also critically on dynamic changes in the morphology of the coronary plaque.

Acute Plaque Changes

- Myocardial ischemia underlying the acute coronary syndrome is precipitated by abrupt changes in plaque followed by thrombosis.
- Acute changes in the morphology of chronic atherosclerotic plaques include fissuring, hemorrhage into the plaque, and overt plaque rupture with embolization of atheromatous debris into distal coronary vessels.
- Local disruption of plaque increases the risk of platelet aggregation and thrombosis at that site. Such acute changes often develop in plaques associated with lesser degrees of stenosis than the critical 75% stenosis.



Coronary Artery Thrombosis

- Plaque rupture exposes thrombogenic lipids and subendothelial collagen, thus initiating a wave of platelet aggregation, thrombin generation, and, ultimately, thrombus formation
- If the vessel is completely occluded by the thrombus overlying the ruptured plaque, MI occurs.
- In contrast, if the luminal obstruction by the thrombus is incomplete and dynamic, the patient may develop unstable angina or a lethal arrhythmia, giving rise to sudden cardiac death.

Coronary Artery Vasospasm

133

- This usually occurs in patients with at least some preexisting atherosclerosis.
- At the site of plaque disruption, it may be induced by the release of vasospastic mediators such as thromboxane A₂ from platelet aggregates. Endothelial dysfunction may also precipitate vasospasm by reduced elaboration of endothelial cell-derived relaxing factors.

Other Pathologic Processes

- ☐ Uncommonly, processes other than atherosclerosis or its complications may compromise blood flow through the coronary arteries
- ✓ Emboli originating from vegetations on the aortic or mitral valves and coronary vasculitis.
- ✓ Severe systemic hypotension
- ✓ Increased myocardial oxygen demand may also contribute to the development of myocardial ischemia. This often occurs because of left ventricular myocardial hypertrophy, as might be encountered in patients with systemic hypertension or diseases of the heart valves.

Angina Pectoris

- Symptom complex of IHD characterized by paroxysmal & usually recurrent attacks of substernal or precordial chest discomfort caused by transient (15 seconds to 15 minutes) myocardial ischemia that falls short of infarction.

- Three major variants of angina pectoris are recognized
- Typical (stable) angina pectoris
- Prinzmetal (variant) angina
- Unstable angina pectoris.

Typical or stable angina pectoris

- Episodic chest pain associated with exertion or some other form of stress.
- The pain is classically described as a crushing or squeezing substernal sensation, which may radiate down the left arm.
- Stable angina pectoris is usually associated with a *fixed atherosclerotic narrowing* (usually 75% or greater) of one or more coronary arteries.
- With this degree of obstruction ("critical stenosis"), the myocardial oxygen demand may be adequate under basal conditions but cannot be augmented sufficiently to meet the increased requirements imposed by exercise or other conditions that stress the heart.
- The pain is usually relieved by rest (reducing demand) or by administration of nitroglycerin. This vasodilator reduces venous blood delivered to the heart (and hence cardiac work) by causing venous dilation; in larger doses, it may increase blood supply to the myocardium by coronary vasodilation.

Prinzmetal, or variant, angina

- ▣ refers to uncommon pattern of angina that occurs at rest
- ▣ It is associated with coronary artery spasm. Although the spasm usually occurs near an atherosclerotic plaque, it may affect a normal vessel.
- ▣ they respond to the administration of vasodilators.

Unstable angina pectoris, sometimes called *crescendo angina*

139

- characterized by the increased frequency of anginal pain. The attacks tend to be precipitated by progressively less exertion, and they are more intense and often last longer than episodes of stable angina pectoris.
- Unstable angina is a harbinger of more serious, potentially irreversible myocardial ischemia and hence is sometimes referred to as *preinfarction angina*.
- In most patients it is induced by acute plaque change with superimposed partial thrombosis, distal embolization of the thrombus, and/or vasospasm.

Myocardial Infarction

- The term *myocardial infarction* indicates the *development of an area of myocardial necrosis caused by local ischemia*.
- MI, also known as "heart attack," is the single most common cause of death in industrialized nations

Transmural versus Subendocardial infarction

141

- ☐ *Transmural* – when necrosis involves the full thickness of the ventricular wall. This pattern is usually associated with coronary atherosclerosis, acute plaque change & superimposed thrombosis
- *Subendocardial (nontransmural) infarct* – an area of necrosis limited to the inner one third or half of the ventricular wall. It can result from acute plaque change, from sufficiently prolonged & severe reduction in systemic blood pressure(eg shock) often on superimposed on chronic coronary stenosis.

- The major risk factors for MI are the same as those discussed previously for coronary atherosclerosis

Pathogenesis

☐ *MIIs are caused by coronary artery thrombosis.*

Disruption of an underlying atherosclerotic plaque (e.g., fissure formation) serves as the nidus for the generation of the thrombus in many cases.

Vasospasm and platelet aggregation may contribute to coronary artery occlusion, but they are seldom, if ever, the sole cause of the occlusion.

- In 10% of cases, other mechanisms may cause MI
- ❖ Vasospasm
- ❖ Emboli

Myocardial response

- The loss of critical blood supply to myocardium induces profound functional, biochemical & morphologic consequences.

Approximate Time of Onset of Key Events in Ischemic Cardiac Myocytes

Feature	Time
Onset of ATP depletion	Seconds
Loss of contractility	<2 min
ATP reduced	
to 50% of normal	10 min
to 10% of normal	40 min
Irreversible cell injury	20–40 min
Microvascular injury	>1 hr

- Classic myocardial necrosis with extensive damage begins at approximately 30 minutes after coronary occlusion. Classic acute MI occurs when perfusion of the myocardium is reduced severely below its needs for an extended interval (at least 2 hrs)

- 148
- The precise location, size & specific morphologic features of an acute myocardial infarction depend on
 - The location , severity, rate of development of coronary atherosclerotic obstructions
 - The duration of the occlusion
 - The metabolic /oxygen needs of the myocardium at risk
 - The extent of collateral blood vessels
 - The presence, site & severity of coronary arterial spasm
 - Other factors – change in blood pressure, heart rate & cardiac rhythm

Morphology


149

- *The location of an MI is determined by the site of the vascular occlusion and by the anatomy of the coronary circulation.*

- Occlusion of the left anterior descending coronary artery (40% to 50%) typically causes an infarct in the anterior and apical areas of the left ventricle and anterior ventricular septum (anteroapical MI).

- Occlusion of the right coronary artery (30% to 40%) is responsible for most infarcts involving the posterior and basal portions of the left ventricle, posterior portion of ventricular septum, free wall of right ventricle .

- Left circumflex coronary artery (15% to 20%):
infarct involves lateral wall of left ventricle except at
apex




- The gross & microscopic appearance of an infarct at autopsy depends on the duration of survival of the patient following the MI.

- The area of damage undergo a progressive sequence of morphologic changes that consists of ischemic coagulative necrosis, followed by inflammation & repair.

0-4hrs


No gross or microscopic changes



4-12 hrs

Gross – occasionally dark mottling


Mic- beginning coagulation necrosis ,edema & hemorrhage



12-24hrs

Gross- dark mottling


Mic- Ongoing coagulation necrosis; pyknosis of nuclei;
myocyte hypereosinophilia; marginal contraction
band necrosis; beginning neutrophilic infiltrate




1-3days

Gross- Pallor with some hyperemia

Mic- Total loss of nuclei and striations along with heavy neutrophilic infiltrate






3-7 days

Gross - Hyperemic border with central yellowing

Mic- Macrophage and mononuclear infiltration begin,
fibrovascular response begins





7-10days


Gross – maximally yellow & soft with depressed red tan margins

Mic- well developed phagocytosis of dead cells , early formation of granulation tissue

10-14days

Gross – Red-gray depressed infarct borders

Mic- well established granulation tissue



2-8wks


Gross- Gray-white scar


Mic- increased collagen deposition, with decreased cellularity

>2 mo

Gross – scarring complete

Mic- dense collagenous scar





Clinical Features

- MI is diagnosed classically by typical symptoms , biochemical evidence & by the ECG pattern

- The onset of MI is usually accompanied by severe, crushing substernal chest pain, which may radiate to the neck, jaw, epigastrium, shoulder, or left arm.
- In contrast to the pain of angina pectoris, the pain associated with an MI typically lasts several hours to days and is not significantly relieved by nitroglycerin.

- The pulse is generally rapid and weak, and patients are often diaphoretic(sweating profusely).
- Dyspnea is common and is caused by impaired contractility of the ischemic myocardium, with resultant pulmonary congestion and edema.
- With massive MIs involving over 40% of the left ventricle, cardiogenic shock develops.

- In minority of patients (20% to 30%) the MI does not cause chest pain. Such "silent" MIs are particularly common in patients with underlying diabetes mellitus and hypertension and in elderly patients

Laboratory evaluation

- It is based on measuring the blood levels of intracellular macromolecules that leak out of fatally injured myocardial cells through damaged cell membranes.
- These molecules include myoglobin, cardiac troponins T & I, creatine kinase (CK), lactate dehydrogenase

Creatine kinase (CK)

- an enzyme that is highly concentrated in brain, myocardium, and skeletal muscle.
- The enzyme is composed of two dimers designated "M" and "B." CK-MB principally from myocardium, although variable amounts of the MB form are also present in skeletal muscle.
- ▣ *Total CK activity* begins to rise within 2 to 4 hours of an MI, peaks at 24 hours, and returns to normal within approximately 72 hours.
- ▣ Although *total CK activity is one of the most sensitive determinants of acute myocardial necrosis, it is not specific*, because CK is also elevated in other conditions, such as skeletal muscle injury.
- ▣ The specificity for the detection of MI is enhanced by measurement of the CK-MB fraction. CK-MB rises within 2 to 4 hours of the onset of an MI, peaks at 18 hours, and usually disappears by 48 hours.
- An absence of a change in the levels of CK and CK-MB during the first 2 days after the development of chest pain essentially excludes the diagnosis of MI

☐ *Troponins*


☐ A group of proteins found in both human skeletal and cardiac muscle. They regulate calcium-mediated contraction of muscles.


- Cardiac troponin I is found only in heart muscle and is thus more specific than CK-MB, which is also found in skeletal muscle in small proportion.
- After an acute MI, both cTnT and cTnI levels rise at about the same time as CK-MB. The diagnostic sensitivity of cardiac troponin measurements is similar to that of CK-MB in the early stages of acute MI. In contrast to CK-MB levels, however, troponin levels remain elevated for 4 to 7 days after the acute event, allowing the diagnosis of an acute MI long after CK-MB levels have returned to normal.


Consequences & complications of MI

182

- Cardiac arrhythmias
- Left ventricular failure with mild to severe pulmonary edema
- Cardiogenic shock
- Rupture of free wall, septum, or papillary muscle
- Thromboembolism
- Pericarditis
- Ventricular aneurysm







Chronic Ischemic Heart Disease

186

- The term *chronic ischemic heart disease*, sometimes called *ischemic cardiomyopathy*, is used to describe the development of progressive congestive heart failure as a consequence of long-term ischemic myocardial injury.
- Many cases are associated with a history of angina pectoris and may be preceded by recognized infarcts. In other cases the condition may develop more insidiously.

Sudden Cardiac Death

- It is defined as unexpected death from cardiac causes early after symptom onset (usually within 1 hr) or without symptoms

- *The most common cause of sudden cardiac death is ischemic heart disease.*
- Chronic ischemia predisposes the myocardium to the development of lethal ventricular arrhythmias, usually in the form of ventricular fibrillation, which is the most common cause of sudden death in these cases.
- Sudden death is the initial manifestation of ischemic heart disease in about 50% of patients with the disease.

Nonatherosclerotic causes of SCD include

- Developmental abnormalities (anomalous origin, hypoplasia of coronary artery)
- Aortic stenosis and other forms of left ventricular outflow obstruction
- Mitral valve prolapse
- Myocarditis
- Cardiomyopathies
- Conduction System Abnormalities
- Isolated hypertrophy, hypertensive or unknown cause

SYSTEMIC (LEFT SIDED) HYPERTENSIVE HEART DISEASE


- It is the response of the heart to the increased demands induced by systemic hypertension .

- The diagnosis of hypertensive heart disease is based on the presence of *left ventricular hypertrophy in an individual with a history of hypertension* and in whom other causes of ventricular hypertrophy (e.g., aortic stenosis or primary hypertrophic cardiomyopathy) have been excluded.

MORPHOLOGY

- The essential feature of hypertensive heart disease is left ventricular hypertrophy. The weight of the heart usually exceeds 450 g. The hypertrophy typically involves the ventricular wall in a symmetric, circumferential pattern termed concentric hypertrophy .
- The size of the chamber is normal in the early stages of hypertensive heart disease, but in long-standing cases myocardial dysfunction , cardiac dilation, CHF & sudden death occur.





Clinical Features

- In its early stages, while the cardiac output is maintained at normal levels, hypertensive heart disease may cause no symptoms. In these patients, the diagnosis is usually based on the detection of left ventricular enlargement in chest radiographs or echocardiograms or by electrocardiographic evidence of left ventricular hypertrophy.
- As the left ventricle begins to fail, the clinical manifestations of heart failure appear. Heart failure in the setting of hypertension is associated with a poor prognosis.

- There is substantial evidence that effective control of hypertension can prevent or lead to regression of hypertrophy and its associated risks

PULMONARY HYPERTENSIVE HEART DISEASE(COR PULMONALE)


- The term *cor pulmonale*, or *pulmonary heart disease*, is used to describe disease of the right-sided cardiac chambers caused by pulmonary hypertension resulting from pulmonary parenchymal or pulmonary vascular disease.

- The condition may be acute or chronic.

- *Acute cor pulmonale* is most often caused by pulmonary embolism. When emboli acutely obstruct more than 50% of the pulmonary vascular bed, the sudden increase in the burden on the right side of the heart causes right ventricular failure. The right ventricle is typically dilated but is not hypertrophied.

- *Chronic cor pulmonale* implies right ventricular hypertrophy (and dilation) secondary to prolonged pressure overload caused by obstruction of the pulmonary arteries or compression or obliteration of septal capillaries (due to primary pulmonary hypertension or emphysema)

Chronic cor pulmonale is characterized by right ventricular, and often right atrial, hypertrophy. When ventricular failure develops, the right ventricle and atrium may also be dilated




VALVULAR HEART DISEASES

- Valvular involvement by disease causes stenosis, insufficiency (regurgitation or incompetence) or both.

Stenosis is the failure of a valve to open completely impeding forward flow.

Regurgitation – results from failure a valve to close completely allowing reversed flow.



- Deformed cardiac valves may cause disease by two major mechanisms.
- ✓ *They impose a major hemodynamic burden on the cardiac chambers by causing obstruction (stenosis) or regurgitation (incompetence) or sometimes a combination of the two.*
- ✓ *The abnormal valves are more susceptible to infection and thus predispose patients to infective endocarditis and its many complications*

- Abnormalities of flow often produce abnormal heart sounds known as murmurs

- Diseases of the heart valves include a diverse group of acquired and congenital lesions. Some of these occur in isolation, and others occur in association with other heart diseases

■ The most frequent causes of the major functional valvular lesions include

209

- Aortic stenosis : calcification of anatomically & congenitally bicuspid aortic valves
- Aortic insufficiency: dilation of the ascending aorta, related to hypertension & aging
- Mitral stenosis: rheumatic heart disease
- Mitral insufficiency: myxomatous degeneration (mitral valve prolapse)

Calcific Aortic Stenosis



- Aortic stenosis is the most common of all valvular abnormalities.
- It is usually the consequence of calcification due to progressive & advanced age associated 'wear & tear' of either previously anatomically normal aortic valves or congenitally bicuspid valves.

MORPHOLOGY

- In degenerative calcific aortic stenosis, the aortic valve leaflets are rigid and deformed by irregular calcified masses.
- The calcium deposits lie behind the valve cusps and extend into the sinus of Valsalva preventing the opening of the cusps.
- Concentric hypertrophy of the left ventricle and ventricular dilation are commonly seen with long-standing disease.



A: Tricuspid Aortic Valve – Senile B: Bicuspid Aortic Valve
C & D: Mitral Valve –



Clinical Features

213

- Degenerative calcific aortic stenosis arising in a previously normal valve is usually asymptomatic until the eighth or ninth decade.
- *In congenitally bicuspid* aortic valves, lesions become clinically manifested 10 to 20 years earlier than degenerative calcific aortic stenosis.

- The cardinal manifestations of degenerative calcific aortic stenosis include angina pectoris, syncope, and (in later stages) congestive heart failure.


- *Angina* occurs because of the increased oxygen requirements of the hypertrophied myocardium coupled with reduced aortic outflow and *Syncope* reflects poor perfusion of the brain, particularly during episodes of exertion.


- Untreated aortic stenosis usually causes death within 3 to 4 years of the onset of symptoms, either from congestive heart failure or from a lethal arrhythmia




Mitral Valve Prolapse (myxomatous degeneration of mitral valve)

- One of the common causes of isolated mitral regurgitation is a disorder known as *mitral valve prolapse*.
- Most cases are discovered between the ages of 20 and 40; the disease is more common in women than in men.

- It is characterized by an accumulation of loose ground substance (myxomatous) within the leaflets and chordae of the mitral valve, which causes the valve to become "floppy" and prolapse into the left atrium during systole making it incompetent.







Pathogenesis

- It is proposed that there is an underlying developmental defect of connective tissue
- Mitral valve prolapse may arise as a complication of Marfan syndrome or similar connective tissue disorders

Clinical Features

- Most patients with mitral valve prolapse are asymptomatic.

- Approximately 3% of patients experience one of several complications. These include hemodynamically significant mitral regurgitation and congestive heart failure, particularly if the chordae or valve leaflets rupture.
- Patients with mitral valve prolapse are also at increased risk for the development of infective endocarditis and sudden death caused by ventricular arrhythmias. Stroke or other systemic infarction may occur from embolism of thrombi formed in the left atrium.


Rheumatic Fever and Heart Disease

- *Rheumatic fever is an acute, immunologically mediated, multisystem inflammatory disease that follows an episode of group A streptococcal pharyngitis after an interval of a few weeks.*

- The most important consequence of RF are chronic valvular deformities characterized principally by deforming fibrotic valvular disease (particularly mitral stenosis)- chronic rheumatic heart disease
- It produces permanent dysfunction & severe cardiac problems decades later.

Pathogenesis

- It is strongly suspected that *acute rheumatic fever* is a *hypersensitivity reaction induced by group A streptococci*.
- It is proposed that antibodies directed against the M proteins of group A streptococci cross-react with normal proteins present in the heart, joints, and other tissues.



- Rheumatic fever doesn't follow infections by streptococci at other sites such as skin.
- Only minority of infected patients develop RF, indicating genetic susceptibility influences the hypersensitivity reaction.

- The chronic sequelae result from progressive fibrosis due to both healing of the acute inflammatory lesions & the turbulence induced by ongoing valvular deformity.


MORPHOLOGY

- In acute rheumatic fever, inflammatory infiltrates may occur in a wide range of sites, including synovium, joints, skin, and (most importantly) the heart.


- This provokes a mixed inflammatory response, which may take the form of either a diffuse cellular infiltrate or a localized aggregation of cells that resembles a granuloma.
- Areas of fibrosis eventually develop at sites of inflammation. Fibrosis is particularly common in cardiac tissues, where it is responsible for the valvular deformities seen in chronic rheumatic heart disease.


- Acute rheumatic carditis is characterized by inflammatory changes in all three layers of the heart, and thus it is appropriately designated a pancarditis.
- The hallmark of acute rheumatic carditis is the presence of multiple foci of inflammation within the connective tissues of the heart, called **Aschoff bodies**

- They contain a central focus of fibrinoid necrosis surrounded by a chronic mononuclear inflammatory infiltrate and occasional large macrophages with vesicular nuclei and abundant basophilic cytoplasm, called **Anitschkow cells**. Aschoff bodies may be found anywhere in the connective tissues of the heart




- **Aschoff body:**
Fibrinoid degeneration
T lymphocytes
Plasma cells
Macrophages
Anitschkow cells
Aschoff giant cells
Around blood vessel.





Pericarditis


- The inflammation is accompanied by fibrinous pericardial exudate , resulting in 'bread-and-butter' pericarditis which resolves without sequelae.






- The myocardial involvement- myocaditis- takes the form scattered Aschoff bodies in the interstitium.





- Involvement of the endocardium & the left sided valves results in fibrinoid necrosis on which sit small vegetations—verrucae—along the lines of closure.






- **Chronic rheumatic heart disease** is characterized by organization of the acute inflammation & subsequent fibrosis.

- The major anatomic changes of the valves in chronic RHD are leaflet thickening, commissural fusion (resulting in 'fish mouth' shape in mitral valve) & shorting & thickening & fusion of the tendinous cords.









© Elsevier 2005





Microscopically, there is diffuse fibrosis & neovascularization that obliterate the originally avascular valves.



- The mitral valve is abnormal in approximately 95% of cases of chronic rheumatic heart disease, and combined aortic and mitral valve disease is present in about 25% of patients.
- Right-sided valvular disease is relatively uncommon.

- Scarring of the valve leaflets may cause a reduction in the diameter of the valve orifice (**stenosis**), or it may prevent proper closure of the valve leaflets, resulting in **regurgitation** of blood during diastole. Sometimes stenosis and regurgitation coexist, although one hemodynamic defect usually predominates.

- In mitral stenosis, the left atrium progressively dilates & may harbor mural thrombosis
- Longstanding congestive changes in the lungs may induce pulmonary vascular & parenchymal changes & lead to right ventricular hypertrophy.
- Damage to the valves also predisposes the patients to superimposed **infective endocarditis**.

Clinical Features

- *Acute rheumatic fever* occurs anywhere from 10 days to 6 weeks after an episode of pharyngitis caused by group A streptococci in about 3% of patients.
- The peak incidence is between the ages of 5 and 15, although younger children and adults may also develop the disease.


- Although pharyngeal cultures for streptococci are negative by the time the illness begins, antibodies to one or more streptococcal enzymes, such as streptolysin O and DNase B, are present in the sera of most patients.

Criteria for diagnosis of RF include major & minor criterias

Major manifestations

- Migratory polyarthrititis of large joints
- Carditis
- Subcutaneous nodules
- Erythema marginatum of the skin
- Sydenham chorea, neurologic disorder with involuntary purposeless movements

© Images Paediatr Cardiol



Minor manifestation

Non specific sign & symptoms that include fever, arthralgia or elevated blood levels of acute phase reactants)

- The diagnosis of RF is established by so called **Jones criteria** : evidence of a preceding group A streptococcal infection, with the presence of two of the major manifestation OR one major & two minor manifestation.

- The predominant clinical manifestation are those of arthritis & carditis.

Polyarthritis- multiple joints are affected one after another & become painful & swollen

Acute carditis – pericardial friction rub, weak heart sound, tachycardia & arrhythmia. The prognosis is good.

- After an initial attack, there is increased vulnerability to reactivation of the disease with subsequent pharyngeal infections & the same manifestations are likely to appear with each recurrence.

- *Chronic rheumatic carditis* usually does not cause clinical manifestations for years or even decades after the initial episode of rheumatic fever.
- The signs and symptoms of valvular disease depend on which cardiac valve or valves are involved.
- patients with chronic rheumatic heart disease may suffer from cardiac hypertrophy and dilation, and congestive heart failure, arrhythmias (particularly atrial fibrillation in the setting of mitral stenosis), thromboembolic complications, and infective endocarditis

Infective Endocarditis

- The term *infective endocarditis* designates infection of the cardiac valves or mural surface of the endocardium, resulting in the formation of an adherent, bulky mass of thrombotic debris and organisms, termed a *vegetation*.
- Most cases are caused by bacteria.

- Infective endocarditis has traditionally been subdivided into acute and subacute forms.

- *Acute endocarditis* is characterized by infection of the valves by organisms of high virulence, such as *Staphylococcus aureus*. Such organisms are capable of infecting even structurally **normal valves** and cause rapidly destructive infection, with little accompanying local host reaction.

- *Subacute endocarditis*, in contrast, is typically associated with infection of previously abnormal valves by organisms of lower virulence, such as α -hemolytic streptococci. The resultant infections tend to progress somewhat more slowly and are often accompanied by the development of a local inflammatory reaction and granulation tissue in the affected valve.

Etiology and Pathogenesis

- ❑ Infection occurs when organisms are implanted on the endocardial surface during episodes of bacteremia.
- ❑ The portal of entry of the agent into the blood stream may be an obvious infection or procedures (surgical or dental).

- Conditions that increase the risk of infective endocarditis include
 - A number of cardiac abnormalities that causes increased hemodynamic trauma to the endocardial surface, such as high pressure shunts within the heart (e.g., small ventricular septal defects) or chronic valvular diseases (e.g., chronic rheumatic heart disease, degenerative calcific aortic stenosis, mitral valve prolapse), prosthetic heart valves.

- Host factors such as neutropenia, immunodeficiency, malignancy, diabetes mellitus & intravenous drug abuse


- Endocarditis of abnormal valves is caused most commonly by streptococcus viridans (50-60%)
- ▣ S. aureus attacks both healthy & deformed valves & are responsible for 10% to 20% of cases.
- ▣ Other causes include bacteria such as enterococci, HACEK group (Haemophilus, Actinobacillus, Cardiobacterium, Eikenella & Kingella)


MORPHOLOGY


- The hallmark of infective endocarditis is the presence of valvular vegetations containing bacteria or other organisms.
- The vegetation is friable bulky, potentially destructive containing fibrin, inflammatory cells & bacteria .


- The aortic and mitral valves are the most common sites of infection, although the valves of the right side of the heart may also be involved, particularly in cases of endocarditis occurring in intravenous drug abusers . The vegetations may be single or multiple and may involve more than one valve.

- Vegetations sometimes erode into the myocardium & cause an abscess cavity – ring abscess




















- **Systemic emboli** may occur at any time because of the friable nature of the vegetations, and they may cause infarcts in the brain, kidneys, myocardium, and other tissues. Because the embolic fragments contain large numbers of virulent organisms, abscesses often develop at the sites of such emboli

Clinical Features

- The onset of infective endocarditis may be gradual or explosive, depending on the organism responsible for the infection. Low-grade fever, malaise, and weight loss are characteristic of cases caused by organisms of low virulence, while more acute cases, in contrast, typically present as high fevers, shaking chills, and other evidence of overt septicemia.



- The *spleen is often enlarged*, and clubbing of the digits may be seen, particularly in subacute cases.

- *Systemic emboli* are very common in all forms of infective endocarditis, manifesting as neurologic deficits, retinal abnormalities, necrosis of the digits, and infarcts of the myocardium and other viscera.
- Entrapment of infected emboli in the walls of blood vessels may cause local infection and weakening of the vessel wall, with the formation of so-called *mycotic* aneurysms.

Uncommon manifestations

- Petechiae (small hemorrhages) may be seen on the skin or mucosal surfaces
- splinter hemorrhages - These hemorrhages are subungual, linear, dark red streaks
- Osler node – subcutaneous nodules in the pulp of digits



- *Blood cultures* should be done the moment the possibility of endocarditis is considered

Nonbacterial Thrombotic Endocarditis

- ▣ *NBTE is characterized by the deposition of small masses of fibrin, platelets, and other blood components on the leaflets of the cardiac valves.*
- ▣ The valvular lesions of NBTE are sterile and do not contain microorganisms.


Pathogenesis

294

- It is thought that hypercoagulable states predispose to its development.
- Malignancies, particularly adenocarcinomas, have been identified in up to 50% of patients with NBTE.
- Although NBTE may occur in otherwise healthy individuals, a wide variety of diseases associated with general debility or wasting are associated with an increased risk of NBTE. The term *marantic endocarditis* has also been used to describe this entity, in recognition of the increased frequency of NBTE in cachectic patients.

Clinical Features

- NBTE is usually asymptomatic. Sometimes, particularly in patients with larger lesions, fragments of the vegetations may embolize and cause infarcts in the brain and other organs.



PRIMARY MYOCARDIAL DISEASES

- Diseases that are intrinsic to myocardial fibers.

▢ Primary myocardial diseases are a diverse group that includes inflammatory disorders (*myocarditis*), immunologic diseases (e.g., rheumatic fever), systemic metabolic disorders (e.g., hemochromatosis), muscular dystrophies (e.g., Duchenne muscular dystrophy), and category of idiopathic diseases termed *cardiomyopathies*.

Cardiomyopathies

- The term *cardiomyopathy* (literally, heart muscle disease) is used to describe *heart disease resulting from a primary abnormality in the myocardium.*

- The diagnosis is made only when all other causes of cardiac failure such as hypertension, ischemic heart disease, valvular & congenital heart disease have been excluded.

There are three clinical, functional & pathologic patterns

- ✓ Dilated cardiomyopathy – most common
- ✓ Hypertrophic cardiomyopathy
- ✓ Restrictive cardiomyopathy

DILATED CARDIOMYOPATHY

- A form of cardiomyopathy characterized by *progressive cardiac hypertrophy, dilation, and contractile (systolic) dysfunction*. It is sometimes called congestive cardiomyopathy.

Pathogenesis

- This clinicopathologic picture can result from a large number of different myocardial insults.

- *Inherited genetic abnormalities* are responsible for a significant percentage (25% to 35%) of cases of dilated cardiomyopathy, including many cases previously consigned to the "idiopathic" category.
- *Mutations in genes coding for cytoskeletal proteins (dystrophin), in particular, appear to be responsible for many cases of hereditary dilated cardiomyopathy.*

- Number of acquired myocardial insults can yield similar clinicopathologic pattern.

✓ Myocarditis

Dilated cardiomyopathy may represent a late stage of myocarditis

✓ Alcohol or other toxicity

Alcohol abuse is also strongly associated with the development of dilated cardiomyopathy, raising the possibility that ethanol toxicity or a secondary nutritional disturbance may be the cause of the myocardial injury.


Nonalcoholic toxic insults such as chemotherapeutic agents (adriamycin) can be cause of myocardial insult


- ✓ Peripartum cardiomyopathy
 - occurs late in pregnancy or several weeks to months post partum. The cause of peripartum cardiomyopathy is poorly understood.


✓ Idiopathic


Morphology

- The heart is enlarged and flabby, with weights often exceeding 900 g. The enlargement is caused by a combination of dilation and hypertrophy of all chambers.
- The microscopic features are nonspecific and include myocyte hypertrophy, interstitial fibrosis, wavy fiber change, and (in some cases) a scanty mononuclear inflammatory infiltrate.









Clinical features

- Dilated cardiomyopathy may occur at any age but is most common between the ages of 20 and 60 years
- Dilated cardiomyopathy is the most common form of cardiomyopathy, accounting for about 90% of cases.
- *The fundamental defect in dilated cardiomyopathy is ineffective contraction.* Hence, the clinical picture is that of progressive congestive heart failure, which becomes refractory to therapy.
-

HYPERTROPHIC CARDIOMYOPATHY




-It is also referred to as *asymmetric septal hypertrophy* and *idiopathic hypertrophic subaortic stenosis*, is a cardiac disorder characterized by *myocardial hypertrophy*, *abnormal diastolic filling*, and (in many cases) *intermittent ventricular outflow obstruction*.


- In about 50% of cases, the disorder is inherited as an autosomal dominant trait with variable expression, *abnormalities in genes encoding sarcomeric contractile proteins appear to play a critical role in the development of hypertrophic cardiomyopathies.*

MORPHOLOGY


- The essential feature of hypertrophic cardiomyopathy is myocardial hypertrophy without ventricular dilation which is most pronounced in the left ventricle and interventricular septum.
- In most cases the interventricular septum is substantially thicker than the free (lateral) wall of the left ventricle

- Although disproportionate hypertrophy can involve the entire septum, septal hypertrophy is usually most conspicuous in the subaortic region. Asymmetric septal hypertrophy is often associated with a significant degree of ventricular outflow obstruction during systole, a feature emphasized in the older designation idiopathic **hypertrophic subaortic stenosis**.





- Microscopically, hypertrophic cardiomyopathy is characterized by a haphazard arrangement of hypertrophied, abnormally branching myocytes surrounded by increased connective tissue . This change is most pronounced in sections taken from the interventricular septum. In later stages of the disease, considerable myocardial fibrosis may develop.



Clinical Features

- The basic physiologic abnormality in hypertrophic cardiomyopathy is *inability to fill a hypertrophic left ventricle during diastole*.
- The limitation of cardiac output and a secondary increase in pulmonary venous pressure cause exertional dyspnea.
- Myocardial ischemia is common, even in the absence of concomitant coronary artery disease, and thus anginal pain is frequent.

- Hypertrophic cardiomyopathy is associated with an increased incidence of ventricular arrhythmias and sudden death and is one of the most common causes of sudden unexplained death in young athletes.


RESTRICTIVE CARDIOMYOPATHY


- ☐ Restrictive cardiomyopathy is a disorder characterized by a *primary decrease in ventricular compliance, resulting in impaired ventricular filling during diastole.*
- ☐ It is considerably less common than either dilated or hypertrophic cardiomyopathy.


- Restrictive cardiomyopathy can be idiopathic or associated with distinct diseases that affect the myocardium , principally, radiation fibrosis, amyloidosis, sarcoidosis, hemochromatosis or metastatic tumor

Morphology

- In restrictive cardiomyopathy , the ventricles are of normal size & the cavities are not dilated & the myocardium is firm.
- The appearance of the heart varies with the cause of the restrictive cardiomyopathy.









Clinical Features


- The physiologic problem in restrictive cardiomyopathy is *a stiff and inelastic ventricle that can be filled only with great effort*.
- As with other cardiomyopathies, the symptoms in patients with restrictive cardiomyopathy include fatigue, exertional dyspnea, and chest pain.
- The course is often complicated by arrhythmias and atrioventricular blocks, particularly when the myocardial fibrosis or infiltrative process encroaches on the conduction system.
- Myocardial contractility, although often normal early in the course of the disease, usually declines, causing congestive heart failure in later stages.




Normal



Dilated
cardiomyopathy



Hypertrophic
cardiomyopathy



Restrictive
cardiomyopathy

Myocarditis

- A group of inflammatory processes of the myocardium that result in injury to the cardiac myocytes.
- The inflammatory process is the cause rather than a response to myocardial injury.

The chief causes of myocarditis


- ❖ Viral infections - the most common cause of myocarditis eg. Coxsackie groups A & B, influenza, echovirus, EBV, HIV
- ❖ Bacterial infection eg diphtheria, leptospirosis, meningococcus
- ❖ Parasitic infections eg chagas disease, toxoplasmosis
- ❖ Ionising radiation
- ❖ Drugs eg doxorubicin

Morphology


- The heart may be of normal size, but more commonly it is dilated.
- The myocardium is flabby and pale. In cases of myocarditis caused by bacteria, frank abscesses may be visible


The histologic appearance varies considerably, depending on the cause of the myocarditis.


- In acute viral myocarditis, the myocardium is edematous and contains an inflammatory infiltrate dominated by lymphocytes and other mononuclear cells . Some degree of myocyte degeneration and/or necrosis is almost always present.



- In patients with bacterial myocarditis, the myocardium contains a neutrophilic infiltrate, sometimes complicated by abscess formation.







Clinical Features

- The clinical manifestations of myocarditis range from an asymptomatic state to severe congestive heart failure. Arrhythmias are common
- In most cases, myocarditis appears to be self-limited, although some patients develop chronic congestive heart failure months to years after the initial insult. If seen for the first time at such a late stage, they may be considered to have dilated cardiomyopathy

PERICARDIAL DISEASES


- Pericardial diseases most frequently are seen in conjunction with local myocardial or mediastinal diseases and in patients with certain systemic conditions.

Pericardial effusion & Hemopericardium

- The parietal pericardium can be distended by fluid of variable composition (pericardial effusion), blood(hemopericardium) or pus (purulent)

- The consequence depend on the ability of the parital percardium to stretch based on the speed of accumulation & the amount of fluid.

- Hemopericardium caused by ruptured MI, traumatic perforation, infective endocarditis, or ruptured aortic aneurysm may produce compression of the atria, vena cavae or the ventricles. As result, cardiac filling is restricted – Cardiac tamponade



- Signs and symptoms of cardiac tamponade include faint distant heart sounds, distended neck veins, declining cardiac output, and shock.

Pericarditis

- Primary pericarditis is uncommon and usually infectious in origin. *Viruses are responsible for most cases*, although it may also be caused by other organisms, including pyogenic bacteria, mycobacteria, and fungi.

- More often the pericarditis is secondary to acute MI, cardiac surgery, or radiation to the mediastinum.
- *Uremia* is probably the most common systemic disorder associated with pericarditis.
- Less common secondary causes include rheumatic fever, systemic lupus erythematosus, and metastatic malignancies. The latter are usually associated with a bloody effusion.

Pericarditis may

- cause immediate hemodynamic complications if a significant effusion is present,
- resolve without significant sequelae, or
- progress to a chronic fibrosing process.

Acute pericarditis



The appearance of acute pericarditis varies with its cause.

Serous pericarditis


- ▣ Serous inflammatory exudates are characteristically caused by noninfectious inflammations such as RF, SLE, tumors & uremia.
- ▣ Sometimes bacterial & viral

Fibrinous & serofibrinous pericarditis

- These are the most frequent type of pericarditis & are composed of serous fluid mixed with a fibrinous exudate.
- Common causes include acute MI, uremia, RF, SLE & trauma
- The exudate is typically fibrinous and imparts a shaggy, irregular appearance to the pericardial surface ("bread and butter" pericarditis)









- The fibrin may be may be digested or become organized
- Loud pericardial friction rub is the most striking characteristics of fibrinous pericarditis

Hemorrhagic pericarditis

- It is fibrinous pericarditis with hemorrhage
- The surface of the heart with hemorrhagic pericarditis demonstrates a roughened and red appearance. Hemorrhagic pericarditis is most likely to occur with metastatic tumor and with tuberculosis (TB).





Purulent or suppurative pericarditis

- The invasion of the pericardial space by infective organisms , bacteria is usually implicated
- The infection typically spreads from the lungs.




Chronic pericarditis

- Organization can result in obliteration of the pericardial sac

Constrictive pericarditis

- The heart may be encased in dense fibrous or fibrocalcific scar that limits diastolic expansion & restricts cardiac output (resembling restrictive cardiomyopathy)

- The heart is surrounded by a dense, adherent layer of scar with or without calcification (0.5 to 1cm)
- The major therapy is pericardiectomy



CARDIAC TUMORS


- The heart is usually spared the ugliness of tumors. Those that occur are usually metastatic

Primary Neoplasms


- Primary tumors of the heart are rare. The most common types encountered, in descending order of frequency, are myxomas, fibromas, lipomas, papillary elastofibromas, rhabdomyomas, angiosarcomas and rhabdomyosarcomas.

Myxomas

- ❏ benign neoplasms that arise most frequently in the left atrium near the fossa ovalis.
- ❏ They appear as sessile or pedunculated lesions, usually covered by an intact endocardium.
- ❏ They occur most often in adults but may arise at any age.



Microscopically, they are composed of multinucleated stellate cells suspended in an edematous, mucopolysaccharide-rich stroma. Immature and mature smooth muscle cells may also be present.



- Although histologically benign, myxomas may cause significant problems for the patient, either because of the tendency of some lesions to fragment and embolize or because they may cause a "ball-valve" obstruction of the atrioventricular valves, with resultant syncopal episodes and even sudden death.

Metastatic Neoplasms

- Metastatic neoplasms involving the heart are far more common than primary cardiac tumors.
- Metastases to the heart occur in up to 10% of patients with disseminated cancer.
- Most often, they involve the pericardium, where they may cause pericarditis and hemorrhagic pericardial effusions.
- The most common primary neoplasms that metastasize to the heart are carcinomas of the lung and breast, malignant melanomas, and hematopoietic malignancies (lymphomas and leukemias). Metastases may reach the heart via lymphatic, venous, or arterial channels.

