

MCQ

Select the single best answer

1. The nerve which innervates a muscle that acts on both the hip and knee joints is:
 - a. Ilioinguinal nerve
 - b. Femoral nerve
 - c. Saphenous nerve
 - d. Common peroneal nerve
 - e. Superficial peroneal nerve
2. During walking, the pelvis is stabilized by which of the following muscles:
 - a. Gluteus maximus
 - b. Obturator internus
 - c. Gluteus medius
 - d. Obturator externus
 - e. Quadratus femoris
3. Which of the following muscles is the main flexor of the thigh?
 - a. Superior gemellus
 - b. Adductor longus
 - c. Gracilis
 - d. Psoas
 - e. Obturator internus
4. Which of the following muscles dorsiflexes the foot at the ankle joint?
 - a. Peroneus longus
 - b. Extensor digitorum brevis
 - c. Tibialis posterior
 - d. Extensor hallucis brevis
 - e. Tibialis anterior
5. In the femoral canal, the neck of the femoral hernia lies:
 - a. Below and lateral to the pubic tubercle.
 - b. Above and medial to the pubic tubercle.
 - c. At the saphenous opening.
 - d. In the obturator canal.
 - e. Lateral to the iliacus muscle.
6. The peroneal artery is a branch of the:
 - a. Anterior tibial artery.
 - b. Popliteal artery.
 - c. Posterior tibial artery.
 - d. Arcuate artery.
 - e. Lateral plantar artery.
7. Locking of the knee joint at the end stage of extension is caused by the action of the:
 - a. Vastus medialis muscle.
 - b. Articularis genu muscle.
 - c. Gastrocnemius muscle.
 - d. Biceps femoris muscle.
 - e. Popliteus muscle.

- Unlocking of the knee joint to permit flexion is caused by the action of the:
- Vastus medialis muscle.
 - Articularis genu muscle.
 - Gastrocnemius muscle.
 - Biceps femoris muscle.
 - Popliteus muscle.
- In the adult, the chief arterial supply to the head of the femur is from the:
- Superior circumflex iliac artery.
 - Obturator artery.
 - Branches of the medial and lateral circumflex femoral arteries.
 - Deep external pudendal artery.
 - Inferior gluteal artery.
- The lymph drainage of the skin covering the ball of the big toe is into the:
- Vertical group of superficial inguinal nodes.
 - Horizontal group of superficial inguinal nodes.
 - Popliteal nodes.
 - Axillary nodes.
 - Internal iliac nodes.
- The lymph drainage of the skin covering the medial side of the knee joint is into the:
- Popliteal nodes.
 - Internal iliac nodes.
 - Vertical group of superficial inguinal nodes.
 - Horizontal group of superficial inguinal nodes.
 - Obturator nodes.
- The lymph drainage of the skin of the buttock (gluteal region) is into the:
- Axillary nodes.
 - Superior gluteal nodes.
 - Vertical group of superficial inguinal nodes.
 - Horizontal group of superficial inguinal nodes.
 - Internal iliac nodes.
- The lymph drainage of the skin of the calf (back of the leg) is into the:
- Vertical group of superficial inguinal nodes.
 - Internal iliac nodes.
 - Horizontal group of superficial inguinal nodes.
 - Popliteal lymph nodes.
 - Obturator lymph nodes.
- Hyperabduction of the hip joint is prevented by the:
- Pubofemoral ligament.
 - Iliofemoral ligament.
 - Ischiofemoral ligament.
 - Iliotibial tract.
 - Ligamentum teres.
- Hyperextension of the hip joint is prevented by the:
- Obturator internus tendon.
 - Iliofemoral ligament.
 - Tensor fascia lata muscle.
 - Iliotibial tract.
 - Ligamentum teres.

prevents dislocation of the femur backward at the knee joint:

16. The _____
- posterior cruciate ligament
 - anterior cruciate ligament
 - medial collateral ligament
 - lateral collateral ligament
 - tendon of the popliteus muscle

prevents abduction of the tibia at the knee joint.

17. The _____
- posterior cruciate ligament
 - anterior cruciate ligament
 - lateral (fibular) collateral ligament
 - lateral meniscus
 - medial (tibial) collateral ligament

is attached to the head of the fibula (styloid process):

18. The _____
- lateral meniscus
 - lateral collateral ligament
 - anterior cruciate ligament
 - posterior cruciate ligament
 - medial meniscus

19. The calcaneus participates in the formation of which arch(es) of the foot?

- Medial longitudinal arch only
- Medial and lateral longitudinal arches
- Transverse arch only
- Medial longitudinal and transverse arches
- Lateral longitudinal and transverse arches

20. The talus participates in the formation of which arch(es) of the foot?

- Transverse arch only
- Lateral longitudinal arch only
- Medial longitudinal arch only
- Medial and lateral longitudinal arches
- Transverse and medial longitudinal arches

21. The 3 cuneiform bones participate in the formation of which arch(es) of the foot?

- Medial longitudinal arch only
- Lateral longitudinal arch only
- Transverse arch only
- Medial longitudinal and transverse arches
- Lateral longitudinal and transverse arches

22. The cuboid bone participates in the formation of which arch(es) of the foot?

- Medial longitudinal arch only
- Lateral longitudinal arch only
- Transverse arch only
- Medial longitudinal and transverse arches
- Lateral longitudinal and transverse arches

23. The sesamoid bones below the head of the 1st metatarsal bone participate in the:

- Lateral longitudinal arch only
- Medial longitudinal arch only
- Medial longitudinal and transverse arches
- Transverse arch only
- Lateral longitudinal and transverse arches

24. Regarding the dorsalis pedis artery, one is incorrect:

- a. It is a continuation of the anterior tibial artery.
- b. It enters the sole of the foot by passing through the 1st interosseous space.
- c. It can be palpated between tibialis anterior & extensor hallucis longus tendons.
- d. It ends by joining the lateral plantar artery (plantar arch).
- e. It lies on the medial side of the deep peroneal nerve.

25. Which of the following structures does not bound the popliteal fossa?

- a. The semimembranosus muscle
- b. The plantaris
- c. The biceps femoris muscle
- d. The medial head of the gastrocnemius muscle
- e. The soleus

26. Which of the following structures does not pass through the greater sciatic foramen?

- a. The superior gluteal artery
- b. The sciatic nerve
- c. The obturator internus tendon
- d. The pudendal nerve
- e. The inferior gluteal vein

27. Which of the following structures does not bound the femoral ring?

- a. The femoral vein
- b. The lacunar ligament
- c. The superior ramus of the pubis
- d. The femoral artery
- e. The inguinal ligament

28. One of the following doesn't pass through the subsartorial (adductor) canal:

- a. The descending genicular artery.
- b. The nerve to vastus lateralis
- c. The femoral artery
- d. The saphenous nerve
- e. The femoral vein

29. Which of the following muscles is not present in the floor of the femoral triangle?

- a. The pectineus
- b. The adductor longus
- c. The iliacus
- d. The psoas
- e. The adductor brevis

30. Which of the following statements is incorrect regarding the ankle joint?

- a. It is strengthened by the deltoid (medial) ligament.
- b. It is a synovial joint.
- c. It is a hinge joint.
- d. It is formed by the articulation of the talus, lower ends of the tibia and fibula.
- e. It is most stable in the fully plantar-flexed position.

31. The foot is everted by the following muscle:

- a. The tibialis anterior
- b. The extensor hallucis longus
- c. The extensor digitorum longus
- d. The peroneus longus
- e. The tibialis posterior

32. Which group of muscles below is supplied by the obturator nerve?

- the chief flexors of the thigh
- the chief extensors of the thigh
- the chief adductors of the thigh
- the chief abductors of the thigh
- the chief lateral rotators of the thigh

33. Following a car accident and some healing time, a patient is not able to rise from a sitting position. This could be due to a lesion of which of the following nerves?

- Femoral nerve
- Obturator nerve
- Nerve to iliopsoas
- Superior gluteal nerve
- Inferior gluteal nerve

34. Loss of pain sensation from the skin between the big toe and 2nd toe indicates damage of which of the following nerves?

- Sural
- Saphenous
- Superficial peroneal
- Deep peroneal
- Lateral plantar

35. A man was hit in the leg by a bat producing a fracture of the head and neck of the fibula and damaging a nerve to the leg. Physical finding that would be found is:

- Inability to plantar flex his foot
- Inability to dorsiflex his foot (foot drop)
- The foot would be everted
- Flexion of the leg is lost
- Extension of the leg is lost

36. A misplaced intramuscular injection into the gluteal region infiltrated the sciatic nerve causing temporary paralysis of the muscles supplied by that nerve. Which of the following muscles spare the paralysis?

- Tibialis anterior
- Soleus
- Biceps femoris
- Gracilis
- Popliteus

37. When you stand on one foot, the opposite side of the pelvis has a tendency to drop. Which of the following prevents this from happening?

- Hamstring muscles
- Piriformis
- Iliacus
- Gluteus maximus
- Gluteus medius

38. The nerve that is responsible for the above is the:

- femoral
- obturator
- sciatic
- superior gluteal
- inferior gluteal

39. The superior and inferior gluteal arteries arise from the:

- a. common iliac artery
- b. external iliac artery
- c. internal iliac artery
- d. femoral artery
- e. median sacral artery

40. The femoral hernia descends:

- a. in the medial compartment of the femoral sheath.
- b. in the middle compartment of the femoral sheath
- c. in the lateral compartment of the femoral sheath
- d. lateral to the femoral vein.
- e. anterior to the inguinal ligament.

41. Paralysis of the quadriceps femoris results MAINLY in loss of:

- a. flexion of the thigh
- b. extension of the thigh
- c. flexion of the leg
- d. extension of the leg
- e. lateral rotation of the leg

42. If a patient lost sensations over his heel, which nerve is not functioning?

- a. Femoral nerve
- b. Medial plantar nerve
- c. Lateral plantar nerve
- d. Deep peroneal nerve
- e. Tibial nerve

43. Which one of the following is the principal invertor of the foot?

- a. Tibialis anterior
- b. Peroneus longus
- c. Popliteus
- d. Soleus
- e. Gastrocnemius

44. The medial border of the femoral triangle is formed by the:

- a. sartorius muscle
- b. adductor longus muscle
- c. gracilis muscle
- d. inguinal ligament
- e. pectineus muscle

45. At the ankle joint, the tibia and fibula articulate with the:

- a. Calcaneus
- b. Cuboid
- c. Cuneiform
- d. Talus
- e. Navicular

46. The abductor digiti minimi muscle is located in which layer of the sole of the foot?

- a. First
- b. Second
- c. Third
- d. Fourth
- e. Fifth

47. Anterior displacement of the tibia at the knee joint is prevented by which ligament:
- Tibial (medial) collateral ligament
 - Fibular (lateral) collateral ligament
 - Anterior cruciate ligament
 - Posterior cruciate ligament
 - Transverse ligament
48. The following is found superficially immediately anterior to the medial malleolus?
- Dorsalis pedis artery
 - Arcuate artery
 - Anterior tibial artery
 - Small saphenous vein
 - Greater saphenous vein
49. The dorsalis pedis artery is usually a direct continuation of which artery?
- Femoral
 - Popliteal
 - Peroneal
 - Anterior tibial
 - Posterior tibial
50. The safest place to administer an intramuscular injection in the gluteal region is the:
- Upper medial quadrant
 - Lower medial quadrant
 - Upper lateral quadrant
 - Lower lateral quadrant
 - None of the above
51. The medial plantar nerve is a branch of the:
- Femoral nerve
 - Deep peroneal nerve
 - Common peroneal nerve
 - Tibial nerve
 - Saphenous nerve
52. The ligament which prevents anterior displacement of the femur at the knee joint is:
- Anterior cruciate ligament
 - Posterior cruciate ligament
 - Transverse ligament
 - Tibial collateral ligament
 - Popliteal ligament
53. The most powerful (the main) extensor of the thigh is the:
- Gluteus maximus
 - Gluteus medius
 - Psoas major
 - Obturator externus
 - Piriformis
54. If the foot is permanently dorsiflexed and everted, which nerve is NOT functioning?
- Femoral nerve
 - Tibial nerve
 - Deep peroneal nerve
 - Common peroneal nerve
 - Superficial peroneal nerve

55. Nerve that can be compressed against the head and neck of the fibula is the:
- Saphenous nerve
 - Tibial nerve
 - Common peroneal nerve
 - Obturator nerve
 - Sural nerve
56. Which of the following is NOT a lateral rotator of the hip?
- Piriformis
 - Gluteus medius
 - Obturator internus
 - Obturator externus
 - Quadratus femoris
57. Which muscle is NOT a flexor of the thigh?
- Pectineus
 - Rectus femoris
 - Iliopsoas
 - Sartorius
 - Adductor magnus
58. The greater sciatic foramen is bounded by which of the following ligaments?
- Sacroiliac
 - Sacrotuberous
 - Inguinal
 - Uterosacral
 - None of the above
59. In the following list what is the main invertor of the foot?
- Peroneus brevis
 - Tibialis posterior
 - Flexor hallucis longus
 - Flexor digitorum longus
 - Peroneus longus
60. The nerve supply of the muscles inserted into the iliotibial tract is:
- Superior gluteal and sciatic nerves
 - Inferior gluteal and sciatic nerves
 - Superior and inferior gluteal nerves
 - Sciatic and obturator nerves
 - Obturator and femoral nerves
61. The Sciatic nerve:
- Is one of the branches of lumbar plexus
 - Enters the gluteal region by passing through lesser sciatic foramen
 - Lies superficial to gluteus maximus muscle
 - Divides into deep and superficial peroneal nerves
 - Supplies ischeal part of adductor magnus muscle
62. One of the following muscles is NOT inserted in the greater trochanter:
- Piriformis
 - Obturator internus
 - Obturator externus
 - Gluteus medius
 - Inferior gemellus

63. Which one of the following passes through both greater and lesser sciatic foramina?
- Nerve to quadratus femoris
 - Pudendal nerve
 - Tendon of obturator internus
 - Sciatic nerve
 - Inferior gluteal artery
64. Nerve to obturator internus supplies:
- Superior gemellus
 - Inferior gemellus
 - Obturator externus
 - Quadratus femoris
 - Tensor fascia lata
65. The small (short) saphenous vein terminates into:
- Femoral vein
 - Popliteal vein
 - Great saphenous vein
 - Dorsal venous arch
 - Vena comitantes of posterior tibial artery
66. The nerve supply of the muscles forming tendocalcaneus is:
- Deep peroneal nerve
 - Anterior tibial nerve
 - Superficial peroneal nerve
 - Common peroneal nerve
 - Tibial nerve
67. The plantar arch is formed mainly by:
- Dorsalis pedis artery
 - Medial plantar artery
 - Lateral plantar artery
 - Arcuate artery
 - 1st plantar metatarsal artery
68. Pain from medial side of the dorsum of the foot is carried by the:
- Saphenous nerve
 - Sural nerve
 - Medial plantar nerve
 - Lateral plantar nerve
 - Posterior tibial nerve
69. Pain from lateral side of the dorsum of the foot is carried by the:
- Saphenous nerve
 - Sural nerve
 - Superficial peroneal nerve
 - Deep peroneal nerve
 - Lateral plantar nerve
70. The spring ligament connects the following 2 bones:
- Calcaneus and cuboid
 - Calcaneus and talus
 - Calcaneus and navicular
 - Cuboid and navicular
 - Metatarsal bones

- 148
71. After a football injury, a physician noticed that the tibia could be moved anteriorly, especially when the knee was flexed. This disability should be diagnosed as a torn:
- Lateral collateral ligament
 - Medial collateral ligament
 - Anterior cruciate ligament
 - Posterior cruciate ligament
 - Oblique popliteal ligament
72. The muscle which flexes the hip joint and extends the knee joint is the:
- Quadratus femoris.
 - Rectus femoris.
 - Biceps femoris.
 - Psoas major.
 - Sartorius.
73. The muscle which extends the hip joint and flexes the knee joint is the:
- Gluteus maximus.
 - Psoas major.
 - Rectus femoris.
 - Biceps femoris.
 - Sartorius.
74. The dermatome of the upper part of the front of the thigh (inguinal region) is:
- L1.
 - L3.
 - L4.
 - L5.
 - S1.
75. The dermatome of the big toe is:
- L1.
 - L3.
 - L4.
 - L5.
 - S1.
76. The dermatome of the little toe is:
- L1.
 - L3.
 - L4.
 - L5.
 - S1.
77. The dermatome of the lower part of the front of the thigh (patellar region) is:
- L1.
 - L3.
 - L4.
 - L5.
 - S1.
78. Which one of the following joints is NOT a synovial joint?
- Hip joint
 - Knee joint
 - Superior tibiofibular joint
 - Inferior tibiofibular joint
 - Ankle joint

MCQ ANSWER KEY

1	B
2	C
3	D
4	E
5	A
6	C
7	D
8	E
9	C
10	A
11	C
12	D
13	D
14	A
15	B
16	B
17	E
18	B
19	B
20	C

21	D
22	E
23	B
24	C
25	E
26	C
27	D
28	B
29	E
30	E
31	D
32	C
33	E
34	D
35	B
36	D
37	E
38	D
39	C
40	A

41	D
42	E
43	A
44	B
45	D
46	A
47	C
48	E
49	D
50	C
51	D
52	B
53	A
54	B
55	C
56	B
57	E
58	B
59	B
60	C

61	E
62	E
63	B
64	A
65	B
66	E
67	C
68	A
69	B
70	C
71	C
72	B
73	D
74	A
75	D
76	E
77	B
78	D
GOOD LUCK	

1. B. The femoral nerve innervates the rectus femoris muscle.
 10. A. The lymphatic vessels follow the course of the great saphenous vein.
 13. D. The lymphatic vessels follow the course of the small saphenous vein.
 24. C. The dorsalis pedis pulse can be felt on the dorsum of the foot lateral to EHL tendon.
 30. E. Ankle joint is most stable in dorsiflexion.