

Lower limb

Anterior compartment of the thigh

1) The iliotibial tract:

- a. lies lateral to tensor fascia latae
- b. inserts onto the head of the fibula
- c. is the origin of 60% of gluteus maximus
- d. is a weak flexor of the knee via tensor fascia latae
- e. is attached superiorly to the iliac crest

2) Which of the following muscles is not found in the floor of the femoral triangle?

- a. iliacus
- b. psoas
- c. pectineus
- d. adductor magnus
- e. adductor longus

3) All the following are branches of the femoral nerve except:

- a. medial femoral cutaneous nerve
- b. saphenous nerve
- c. nerve to vastus medialis
- d. lateral femoral cutaneous nerve
- e. nerve to Sartorius

4) Which factor is most important in preventing lateral displacement of the patella during quadriceps contraction?

- a. the forward prominence of the lateral condyle of the femur
- b. the action of articularis genu
- c. the action of vastus medialis
- d. the action of rectus femoris
- e. the tension of the medial patellar retinaculum

5)Cutaneous nerve supply of the thigh involves all but which of the following:

- a. the intermediate femoral cutaneous nerve
- b. the obturator nerve
- c. the genital branch of the genitofemoral nerve
- d. the ilioinguinal nerve
- e. the posterior cutaneous nerves of the thigh

6)Regarding the femoral triangle:

- a. the lateral border of the adductor longus makes up its medial boundary
- b. it is bisected by the femoral nerve
- c. the lateral border of sartorius muscle makes up its lateral boundary
- d. the long saphenous vein joins the femoral vein within the triangle
- e. rectus femoris makes up part of the floor

7)The femoral nerve:

- a. is formed from the anterior divisions of the anterior primary rami of L2-4
- b. is formed from the same spinal segments as the obturator nerve
- c. divides after passing through the femoral triangle
- d. has no cutaneous branches other than the saphenous nerve
- e. enters the thigh in the femoral sheath

8)Which of the following is true of the saphenous nerve?

- a. is predominantly from L2

- b. supplies adductor magnus
 - c. pierces the deep fascia in the femoral triangle
 - d. is cutaneous only
 - e. terminates just below the knee
- 9) Structures closely related to long saphenous vein at ankle include which of the following?
- a. major perforating veins to deep venous system
 - b. a branch of the femoral nerve
 - c. medial superficial lymphatic trunks
 - d. a branch of the anterior tibial nerve
 - e. dorsalis pedis artery
- 10) The iliotibial tract is the conjoined aponeurosis of the tens fasciae lata and:
- a. gluteus minimus
 - b. gluteus medius
 - c. gluteus maximus
 - d. Camper's fascia
 - e. Scarpa's fascia
- 11) The base of femoral triangle is formed by:
- a. sartorius
 - b. adductor longus
 - c. inguinal ligament
 - d. pubic tubercle
 - e. none of the above
- 12) With regard to the femoral triangle
- a. It is bounded by adductor longus and magnus

- b. It has the fascia lata lining its floor
- c. The femoral artery lies between the femoral vein and nerve
- d. It is bounded superficially by the lacunar ligament
- e. The femoral sheath contains only the femoral nerve and artery

13) The femoral nerve

- a. Has superficial branches, one of which supplies the hip joint
- b. Supplies psoas major
- c. Lies within the femoral sheath
- d. Emerges from the medial side of psoas major
- e. Is formed from the posterior divisions of the ventral rami of L2,3,4

14) Contents of the femoral triangle include all except

- a. Genital branch of the genitofemoral nerve
- b. Great saphenous vein
- c. Pectineus
- d. Profunda femoris artery
- e. Femoral canal

15) The femoral nerve

- a. Supplies skin on the back of the thigh
- b. Runs medial to the femoral artery at the level of the femoral canal
- c. Has the sural nerve as the terminal branch
- d. Supplies the posterior half of adductor magnus
- e. Gives a muscular branch to the pectineus

16) The femoral triangle

- a. Has sartorius as its medial border
- b. Contains the posterior division of the obturator nerve
- c. Contains femoral nerve and vessels
- d. Has the lateral border of adductor longus laterally
- e. All of the above

17) The muscles supplied by the femoral nerve include all except

- a. Sartorius
- b. Iliacus
- c. Psoas major
- d. Pectineus
- e. Rectus femoris

18) Concerning the origins of the quadriceps femoris

- a. All 4 component muscles have an origin from the femur
- b. Vastus lateralis arises from the femur above intertrochanteric line and acetabulum
- c. Vastus medialis arises from femur and tendon of adductor longus
- d. Rectus femoris arises from 3 heads
- e. Vastus intermedius arises from shaft of femur and greater trochanter

19) The floor of the femoral triangle consists of all of the following except

- a. Psoas
- b. Iliacus
- c. Pectineus
- d. Adductor longus
- e. Adductor magnus

20) Concerning iliacus

- a. It inserts into the greater trochanter of the femur
- b. It occupies and is attached to the whole of the iliac fossa
- c. Nerve supply is femoral nerve
- d. It acts to flex and laterally rotate the hip
- e. It has the femoral vein as its intermediate anterior relation

1) Sartorius

- a. Arises from the anterior inferior iliac spine
- b. Is pierced by the lateral cutaneous nerve of the thigh
- c. Inserts into the styloid process of the fibula
- d. Is supplied by the obturator nerve
- e. Flexes and laterally rotates the thigh

2) Which of the following muscles is not in the floor of the femoral triangle

- a. Pectineus
- b. Adductor longus
- c. Sartorius
- d. Psoas
- e. Iliacus

) The femoral nerve

- a. Supplies the skin over the lateral thigh
- b. Lies between the pectineus and psoas in the femoral triangle
- c. Is formed by the posterior divisions of the anterior rami of lumbar nerves 2,3 and 4
- d. Breaks into superficial and deep branches, separated by the medial circumflex femoral artery
- e. Supplies the adductor muscles

24) Rectus femoris muscle:

- a. occupies an intermediate plane in the quadriceps muscle mass
- b. arises from the anterior superior iliac spine
- c. supplied by ilioinguinal nerve
- d. has two heads of origin
- e. none of the above

25) Attached to proximal tibial epiphysis:

- a. the whole of the medial collateral ligament of the knee
- b. sartorius
- c. ligamentum patellae
- d. popliteus
- e. semitendinosus

26) Regarding the branches of the femoral nerve:

- a. does NOT supply cutaneous sensation to skin over the femoral triangle
- b. femoral cutaneous nerves pass through the fascia lata via the saphenous canal
- c. saphenous nerve follows the whole course of the great saphenous vein
- d. branch to pectineus passes between the femoral artery and vein
- e. generally branch within the femoral sheath

27) Regarding superficial nerves of the anterior thigh

- a. The ilioinguinal nerve only supplies external genitalia
- b. The genitofemoral nerve supplies the skin over the femoral triangle
- c. The medial femoral cutaneous nerve arises from L3/4

- d. The intermediate femoral cutaneous nerve supplies a small area of skin over the inguinal ligament
- e. The lateral femoral cutaneous nerve runs lateral to the anterior superior iliac spine

28) Femoral nerve

- a. Gives the lateral, medial and intermediate cutaneous nerves of the thigh
- b. Its damage cause anaesthesia of the planter aspect of the foot
- c. Is contained within the femoral sheath
- d. Is formed by the anterior divisions of L2,3,4
- e. Gives a muscular branch to iliacus

29) Regarding the femoral nerve which statement is INCORRECT

- a. Is formed from the posterior divisions of the anterior primary rami of L2, 3, 4
- b. Is formed from the same spinal segments as the obturator nerve
- c. Divides within the femoral triangle
- d. Has no cutaneous branches other than the saphenous nerve
- e. Enters the thigh outside the femoral sheath

30) The skin over the femoral triangle is supplied by

- a. Ilioinguinal nerve
- b. Obturator nerve
- c. Medial femoral cutaneous nerve
- d. Lateral femoral cutaneous nerve
- e. Genitofemoral nerve

31) The deep fascia of the thigh:

- a. is attached to the inguinal ligament
- b. is mostly superficial to the long saphenous vein
- c. receives the insertion of the whole of the gluteus maximus
- d. receives the insertion of the whole of sartorius
- e. is less dense than the deep fascia of the upper arm

32) The muscle which can both flex the hip and extend the knee is:

- a. sartorius
- b. rectus femoris
- c. semimembranosus
- d. biceps femoris
- e. vastus lateralis

33) Difficulty extending the knee can result from damage to the

- a. femoral nerve
- b. inferior gluteal nerve
- c. common peroneal nerve
- d. superior gluteal nerve
- e. tibial nerve

34) The saphenous nerve

- a. is a motor branch of the femoral nerve
- b. accompanies the short saphenous vein in the leg
- c. is the only branch of the femoral nerve that extends considerably below the knee
- d. emerges through the saphenous hiatus
- e. is a sensory branch of the obturator nerve

35) Fascia lata

- a. is the investing fascia of the thigh

- b. Is fused with the inguinal ligament from the anterior superior iliac spine to the pubic tubercle
- c. is thickened laterally forming the iliotibial tract
- d. all of the above are correct
- e. only b and c are correct

36) The branch (es) of the femoral nerve that extend into the leg proper (ie lower leg) is (are):

- a. the motor nerve to the vastus lateralis
- b. the common peroneal nerve
- c. the obturator nerve
- d. the saphenous nerve, carrying only sensory fibers
- e. all of the above

37) What nerve has been damaged in a patient who can no longer extend his knee?

- a. sciatic
- b. common peroneal
- c. femoral
- d. obturator
- e. tibial

38) Which muscle can flex the hip joint and extend the knee joint?

- a. sartorius
- b. adductor longus
- c. biceps femoris
- d. quadriceps femoris
- e. gracilis

39) Which of the following muscles attaches to the anterior superior iliac spine?

- a. Pectineus

- b. sartorius
- c. rectus femoris
- d. vastus intermedius
- e. gracilis

40) The saphenous nerve is a sensory branch of which of the following nerves?

- a. sciatic
- b. femoral
- c. obturator
- d. common fibular (peroneal)
- e. tibial

41) Two muscles form the floor of the femoral triangle on the front of the thigh. Which of the following muscles forms the medial part of the floor?

- a. pectineus
- b. vastus medialis
- c. adductor longus
- d. rectus femoris
- e. iliopsoas

Medial compartment of the thigh

42) The actions of the gracilis muscle is of the thigh at the hip and of the leg at the knee.

- a. lateral rotation, lateral rotation
- b. medial rotation, extension
- c. abduction, flexion
- d. flexion, extension
- e. adduction, flexion

43) After an obturator nerve injury, some adduction of the thigh is

still possible because of double innervation to the:

- a. gracilis m.
- b. adductor magnus m.
- c. sartorius m.
- d. adductor longus m.
- e. adductor brevis m.

44) After passing through the obturator canal, divisions of the obturator nerve (an anterior branch and a posterior branch) pass on either side of the:

- a. pectineus m.
- b. adductor brevis m.
- c. gracilis m.
- d. adductor magnus m.
- e. adductor longus m.

45) Muscles inserting on the medial aspect of the proximal end of the tibia include:

- a. gracilis
- b. hamstring part of adductor magnus
- c. both
- d. neither

46) Which of the following muscles is not innervated by the obturator nerve?

- a. pectineus
- b. adductor brevis
- c. adductor longus
- d. rectus femoris
- e. gracilis

47) The adductor canal:

- a. contains the femoral artery and nerve
- b. ends distally in the adductor longus hiatus
- c. contains no muscular nerves
- d. has adductor longus forming the roof
- e. always has the femoral artery lying between the saphenous nerve and the femoral vein

48) Regarding the adductor compartment:

- a. adductor magnus lies between the anterior and posterior divisions of the obturator nerve
- b. adductor longus inserts into the upper two thirds of the linea aspera of the femur
- c. the hamstring part of adductor magnus is supplied by the tibial part of the sciatic nerve
- d. the medial intermuscular septum separates the adductor compartment from the posterior compartment
- e. obturator externus medially rotates the hip

49) The adductor canal is not:

- a. roofed by fascia containing the subsartorial plexus
- b. occupied by the sciatic nerve
- c. also known as Hunter's canal
- d. occupied by the femoral artery
- e. a gutter shaped groove between vastus medialis and the front of the adductor muscles

50) Which of the following is INCORRECT regarding the adductor canal?

- a. sartorius lies on the fascial roof
- b. in the distal part of the canal, the femoral vein is posterolateral to the artery

- c. at all levels, the artery lies between saphenous nerve and femoral vein
- d. the subsartorial plexus supplies overlying fascia lata and an area of skin above medial side of the knee
- e. the adductors in the floor of the canal are the adductor brevis above and the adductor longus below

51) In the medial compartment of the thigh:

- a. adductor magnus consists of adductor and hamstring muscle parts
- b. the contents are separated from the posterior compartment by the posterior intermuscular septum
- c. adductor longus is the most superficial muscle of the medial side of the thigh
- d. the anterior division of the obturator nerve passes through obturator externus
- e. the posterior division of the obturator nerve is the prime motor supply

52) Which of the following structures is NOT found in the adductor (subsartorial) canal?

- a. femoral artery
- b. femoral vein
- c. femoral nerve
- d. saphenous nerve
- e. nerve to vastus medialis

53) The obturator nerve partly supplies which of the following muscles:

- a. adductor magnus
- b. obturator internus
- c. quadratus femoris

- d. sartorius
- e. inferior gemellus

54) The obturator nerve is derived from the anterior rami of:

- a. T12, L1 and L2
- b. L1, L2 and L3
- c. L2, L3 and L4
- d. L3, L4 and S1
- e. none of the above

55) Which of the following is not an action of gracilis?

- a. adduction of thigh
- b. flexion of knee
- c. extension of thigh
- d. medial rotation of the flexed knee
- e. none of these

56) Obturator externus:

- a. is pierced by femoral circumflex artery
- b. lateral rotator of hip
- c. medial rotator of hip
- d. hip flexor
- e. supplied by S.1

57) Regarding the obturator nerve, all of the following are true EXCEPT

- a. It is formed from the anterior divisions of the anterior rami of L2/3/4
- b. Its anterior division supplies adductor longus, brevis, pectineus and gracilis
- c. Its posterior division supplies all of adductor magnus
- d. It gives a cutaneous branch which supplies the medial thigh

- e. It gives articular branches to the hip and knee joints

58) Which of the following are correctly paired?

- a. Adductor brevis – femoral nerve
- b. Adductor longus – sciatic nerve
- c. Adductor magnus – saphenous nerve
- d. Adductor longus – obturator nerve
- e. Adductor magnus – femoral nerve

59) The adductor canal

- a. Is bounded by adductor brevis
- b. Lies behind sartorius
- c. Contains the femoral nerve
- d. Commences just below the inguinal ligament
- e. Contains the femoral artery which lies infero-laterally to the femoral vein in the lower part

60) The obturator nerve

- a. Emerges on the lateral border of psoas
- b. Supplies obturator externus, adductor longus and the knee joint
- c. Supplies obturator internus, adductor magnus and the hip joint
- d. Runs inferior to the obturator artery in the obturator canal
- e. All of the above

61) Which one of the following muscles has a double nerve supply?

- a- rectus femoris
- b- sartorius
- c- pectineus

- d- adductor longus
- e- gracilis

62) The medial compartment of the thigh contains all of the following except

- a. The obturator nerve
- b. Obturator artery
- c. Gracilis muscle
- d. Obturator internus
- e. Obturator externus

63) The adductor (subsartorial) canal of Hunter is bounded laterally by:

- a. adductor longus
- b. adductor magnus
- c. vastus medialis
- d. sartorius
- e. femoral vein

Gluteal region:

64) The sciatic nerve in the gluteal region:

- a. rests directly on ischium
- b. is accompanied by posterior cutaneous nerve of thigh
- c. supplies obturator internus
- d. supplies quadratus femoris
- e. none of the above

65) Which of the following structures does NOT pass through the greater sciatic foramen?

- a. the sciatic nerve
- b. the pudendal nerve and vessels

- c. the posterior cutaneous nerve of the thigh
- d. the inferior gluteal nerve and vessels
- e. the nerve to obturator externus

66) Passing through the greater sciatic notch deep to the sciatic nerve is which of the following?

- a. internal pudendal vessels and nerve
- b. inferior gemellus muscle
- c. posterior cutaneous nerve of the thigh
- d. nerve to quadratus femoris
- e. none of the above

67) The main function of gluteus maximus is which of the following:

- a. a site for injections
- b. a cushion for sitting
- c. a flexor of the hip
- d. a lateral rotator of the hip
- e. an extensor to the hip

68) The superior gluteal nerve supplies:

- a. gluteus medius
- b. gluteus minimus
- c. tensor fascia latae
- d. none of the above
- e. all of the above

69) The most powerful extensor of the hip is:

- a. gluteus maximus
- b. psoas major
- c. iliacus
- d. obturator externus

- e. piriformis

70) Gluteus maximus

- a. Arises from the gluteal fossa of the ilium between the middle and superior gluteal lines
- b. is supplied by the superior gluteal nerve
- c. extends and medially rotates the hip
- d. mostly inserts into the iliotibial tract
- e. paralysis results in a waddling gait

71) The sciatic nerve

- a. Arises from L3/4/5/S1/2
- b. Emerges from below the lower border of piriformis
- c. Most commonly divides onto common peroneal and tibial branches in the middle of the popliteal fossa.
- d. Can be injured by intramuscular injections in the upper outer quadrant of the buttock
- e. Gives no branches in the gluteal compartment

72) All of the following are lateral rotators of the hip EXCEPT

- a. Gluteus maximus
- b. Gluteus medius
- c. Gemellus superior
- d. Quadratus femoris
- e. Piriformis

73) This structure forms the boundary between the greater and lesser sciatic foramina:

- a. ischial tuberosity
- b. posterior superior iliac spine
- c. sacrotuberous ligament
- d. sacrospinous ligament

e. piriformis muscle

74) This is the only gluteal muscle to originate from the posterior surface of the sacrum:

- a. quadratus femoris
- b. gluteus maximus
- c. gluteus medius
- d. piriformis
- e. biceps femoris

75) If the sciatic nerve were damaged by an intramuscular injection in the buttocks, the patient would have diminished cutaneous sensation on the:

- a. dorsum of the foot
- b. anteromedial side of the leg
- c. both
- d. neither

76) All the following pass through the greater sciatic foramen EXCEPT:

- a. piriformis muscle
- b. pudendal nerve
- c. sciatic nerve
- d. inferior gluteal vessels and nerve
- e. obturator internus tendon

77) The superior gluteal nerve:

- a. contains fibers from the second and third lumbar nerves
- b. leaves the pelvis just inferior to the piriformis muscle
- c. must be intact to resist pelvic tilt during gait

- d. innervates the piriformis muscle
- e. innervates the gemellus superior muscle

78) All of the following muscles are lateral rotators of the thigh the hip EXCEPT the:

- a. gluteus maximus
- b. gluteus minimus
- c. obturator internus
- d. obturator externus
- e. piriformis

79) What nerve has been damaged in a patient whose pelvis is too far toward the right when walking?

- a. left inferior gluteal nerve
- b. right femoral nerve
- c. right superior gluteal nerve
- d. left superior gluteal nerve
- e. right inferior gluteal nerve

80) Which nerve has been damaged in a patient whose pelvis is down to the left when he stands on his right foot?

- a. right inferior gluteal nerve
- b. right superior gluteal nerve
- c. left inferior gluteal nerve
- d. left superior gluteal nerve
- e. left obturator nerve

81) The gluteus maximus muscle:

- a. has an origin from the ischium
- b. inserts only onto the femur
- c. is a flexor of the thigh at the hip

- d. is innervated by the superior gluteal nerve
- e. none of the above

82) The inferior gluteal nerve:

- a. contains fibers from the second and third lumbar nerves
- b. leaves the pelvis just inferior to the piriformis muscle
- c. must be intact to resist pelvic tilt during gait
- d. innervates the tensor fasciae lata muscle
- e. innervates the gemellus inferior muscle

83) Which of the following muscles is not an abductor?

- a. gluteus maximus
- b. gluteus medius
- c. deltoid
- d. supraspinatus
- e. dorsal interosseous.

84) All of the following muscles are lateral rotators of the thigh at the hip joint except

- a. sartorius
- b. obturator externus
- c. obturator internus
- d. gracilis
- e. quadratus femoris

85) Tensor fascia lata is supplied by:

- a. anterior division of femoral nerve
- b. superior gluteal nerve
- c. nerve to vastus lateralis
- d. inferior gluteal nerve

- e. lateral femoral cutaneous nerve

86) Gluteus maximus:

- a. forms the gluteal fold
- b. has four bursae beneath it
- c. has blood supply solely from the inferior gluteal artery
- d. is the chief control of hip flexion
- e. has 50% of fibers inserting into the gluteal tuberosity

87) The sciatic nerve:

- a. lies deep to the posterior femoral cutaneous nerve
- b. passes down over obturator internus, quadratus, femoris and piriformis
- c. tibial and common peroneal components separate behind the hip joint
- d. in the buttock it lies midway between the greater trochanter and pubic tuberosity
- e. is derived from L3, 4, 5, S1, 2

88) Piriformis:

- a. is a lateral rotator of the hip joint
- b. passes through the lesser sciatic foramen
- c. is the preferred site of intramuscular gluteal injection
- d. passes anterior to the femoral neck
- e. lies deep to the sciatic nerve

89) Emerging from lesser sciatic foramen:

- a. piriformis
- b. superior gemellus
- c. obturator internus
- d. inferior gemellus
- e. obturator externus

90) Piriformis is the key to gluteal region. Which relation is not TRUE?

- a. in buttock, lower border lies alongside superior gemellus
- b. inserts on upper border of greater trochanter
- c. emerges through greater sciatic foramen with superior gluteal nerve and vessel above it
- d. emerges through the greater sciatic foramen with pudendal nerve and vessels above it
- e. emerges through greater sciatic foramen with sciatic nerve below it

Back of thigh:

91) With respect to the posterior compartment of the thigh:

- a. blood supply is mainly from the femoral artery
- b. the sciatic nerve lies lateral to the long head of biceps
- c. the long head of biceps is supplied by the common peroneal nerve
- d. the hamstrings form the apex of the popliteal fossa
- e. semimembranosus arises in common with the long head of biceps

92) The hamstring muscles originate from:

- a. body of the ischium
- b. ramus of the pubis
- c. iliac crest
- d. ischial tuberosity
- e. symphysis pubis

93) Which of the following is found in the popliteal fossa?

- a. sciatic nerve
- b. femoral vein
- c. common peroneal nerve

- d. femoral artery
 - e. saphenous nerve
- 94) The posterior cutaneous nerve of the thigh:
- a. consists only of S2, 3
 - b. travels most of its course superficial to the fascia lata
 - c. emerge below piriformis
 - d. emerges above piriformis
 - e. none of the above
- 95) Regarding the hamstring muscles, all of the following are true EXCEPT
- a. Most of the hamstrings arise from the innominate bone
 - b. All of the hamstring muscles are supplied by the sciatic nerve
 - c. The long head of biceps femoris and semitendinosus have a common origin
 - d. The adductor magnus is not part of the hamstring group
 - e. Biceps forms the medial border of the popliteal fossa
- 96) Contents of the popliteal fossa include all of the following EXCEPT
- a. Sural nerve
 - b. Superior medial genicular artery
 - c. Sural communicating nerve
 - d. Inferior medial genicular artery
 - e. Anterior tibial artery
- 97) Regarding the popliteal fossa:
- a. the common peroneal nerve passes through the lateral part of the fossa
 - b. it contains no lymph nodes
 - c. it has a roof pierced by the tibial nerve
 - d. the popliteal artery lies superficial to the popliteal vein throughout the fossa

- e. the small saphenous vein joins the popliteal vein before its entry into the fossa
- 98) Each of the following matchings of thigh muscles with points of origin is correct EXCEPT:
- sartorius—anterior superior iliac spine
 - long head of biceps femoris—ischial tuberosity
 - straight head of rectus femoris—anterior inferior iliac spine
 - pectineus—superior ramus of pubis
 - semitendinosus—linea aspera of femur
- 99) All of the following muscles receive motor innervation from the tibial division of the sciatic nerve except
- semitendinosus
 - semimembranosus
 - long head of biceps femoris
 - short head of biceps femoris
 - popliteus.
- 100) All of the following muscles are medial rotators of the leg EXCEPT:
- semimembranosus
 - semitendinosus
 - biceps femoris
 - gracilis
 - popliteus
- 101) The flexor compartment of the thigh contains which of the following muscles?
- gluteus maximus
 - adductor magnus
 - garacilis
 - semimembranosus
 - rectus femoris
- 102) Biceps femoris muscles:

- a. has a common origin with semitendinosus from the ischial tuberosity
 - b. has a short head attached to the femur medial to the attachment of adductor magnus
 - c. is entirely innervated by common peroneal nerve
 - d. is deep to the common peroneal nerve
 - e. is a lateral rotator of the leg on the thigh when the leg is fully extended at the knee joint
- 103) Which statement about the semimembranosus muscle is FALSE?
- a. it extends the hip joint
 - b. it attaches to the medial condyle of the tibia
 - c. it flexes at the knee joint
 - d. it forms the arcuate ligament
 - e. it is innervated by the tibial portion of the sciatic nerve
- 104) Muscles in the posterior thigh compartment are paralyzed because the sciatic nerve has been severed. Nevertheless the patient can still produce some flexion of the leg at the knee due to action of the:
- a. sartorius m.
 - b. semitendinosus m.
 - c. both A and B
 - d. short head of biceps femoris
 - e. gastrocnemius
- 105) The biceps femoris muscle
- a. lies lateral to the popliteal fossa
 - b. lies medial to the popliteal fossa
 - c. forms the floor of the popliteal fossa
 - d. gives rise to the oblique popliteal fossa
 - e. does not attach to the leg bone
- 106) Which of the following is TRUE?
- a. the popliteal fossa is bounded by the biceps medially

- b. the great saphenous vein runs through the popliteal fossa
- c. gluteus maximus inserts on the greater trochanter
- d. the sciatic nerve leaves the pelvis by passing through the lesser sciatic foramen
- e. the popliteal fossa contains the common peroneal nerve

The leg:

- 107) The muscles of the lateral compartment of the leg are:
- a. dorsiflexors of the foot
 - b. evertors of the foot
 - c. both
 - d. neither
- 108) If the foot is inverted due to paralysis of some muscles, one of the muscles that is paralyzed is the:
- a. tibialis posterior
 - b. tibialis anterior
 - c. peroneus longus
 - d. extensor hallucis longus
 - e. extensor digitorum longus
- 109) The lateral compartment of the leg
- a. usually contains no major artery
 - b. contains only two muscles
 - c. both a and b are correct
 - d. contains the peroneus tertius muscle
 - e. includes muscles innervated by the deep fibular

nerve

- 110) With respect to peroneus longus:
- a. it inserts into the base of the 5th metacarpal
 - b. it is supplied by the superficial peroneal nerve
 - c. it has a broad tendon that lies between the lateral malleolus and the tendon of peroneus brevis
 - d. it arises from the upper two thirds of the tibia
 - e. it is mainly supplied by L4 and L5 segments
- 111) What muscle cause dorsiflexion and inversion of the foot
- a. Tibialis anterior
 - b. Tibialis posterior
 - c. Extensor hallucis longus
 - d. Peroneus tertius
 - e. Peroneus longus
- 112) Tibialis anterior
- a. Dorsiflexes and everts the foot
 - b. Arises from the upper 2/3s of the fibula
 - c. Inserts into the medial cuneiform
 - d. Shares its site of insertion with peroneus tertius
 - e. Is supplied by L5, S1
- 113) Tibialis anterior inserts into
- a. Navicular
 - b. Medial cuneiform and base of first metatarsal
 - c. Lateral cuneiform and base of 5th metatarsal
 - d. Middle three metatarsals

- e. Medial cuneiform and base of 1st phalanx
- 114) Your patient is unable to stand on his toes. The damaged nerve is the:
- a. femoral
 - b. tibial
 - c. superficial peroneal
 - d. deep peroneal
 - e. sural
- 115) Each of the following statements concerning the tibial nerve is true EXCEPT:
- a. It is a branch of the sciatic nerve.
 - b. It is motor to all muscles in the superficial and deep compartments of the leg.
 - c. It passes down the leg between gastrocnemius and soleus
 - d. It passes posterior to the medial malleolus
 - e. Its terminal branches are the medial and lateral plantar nerves.
- 116) Severing the common peroneal nerve results in inability to:
- a. evert the foot
 - b. extend (dorsiflex) the foot
 - c. both
 - d. neither
- 117) The major cutaneous innervation of the anteromedial side of the leg is provided by which nerve?
- a. obturator
 - b. sural
 - c. superficial peroneal

- d. saphenous
- e. medial sural

118) The deep peroneal nerve supplies motor fibers to:

- a. peroneus longus
- b. peroneus brevis
- c. peroneus tertius
- d. flexor hallicis longus
- e. all of the above

119) The nerve supply to the muscles of the lateral fascial compartment of the leg is directly by this nerve:

- a. deep peroneal
- b. superficial peroneal
- c. anterior tibial
- d. posterior tibial
- e. common peroneal

120) What nerve is damaged in a patient whose foot is everted and dorsiflexed and who cannot flex his/her toes?

- a. tibial
- b. superficial peroneal
- c. deep peroneal
- d. common peroneal
- e. medial peroneal

121) A tight plaster cast that exerted pressure on the head and neck of the fibula might result in loss of:

- a. eversion of the foot
- b. foot drop
- c. both
- d. neither

122) A 25 year old man is unable to plantar flex his foot. The most likely cause is damage to:

- a. the superficial peroneal nerve
- b. soleus
- c. gastrocnemius
- d. the L5 nerve root
- e. the tibial nerve

123) If the common peroneal nerve is divided the following are lost:

- a. dorsiflexion of foot
- b. plantar flexion of foot
- c. there is no loss of cutaneous sensation
- d. plantar flexion of toes
- e. inversion of foot

124) The deep peroneal nerve supplies:

- a. skin between the first and second toes
- b. skin between the second and third toes
- c. skin on the medial side of the shin
- d. skin on the dorsolateral side of the foot
- e. the great toe only

125) A 25 years old man is unable to plantarflex his foot. The most likely cause is damage to

- a. The superficial peroneal nerve
- b. L5 nerve root
- c. Tibial nerve
- d. Gastrocnemius

e. Soleus

126) Paralysis of which nerve would result in "foot drop"?
a. femoral
b. tibial
c. superficial peroneal
d. deep peroneal
e. obturator

127) Deep peroneal nerve supplies

- a. Tibialis anterior
- b. tibialis posterior
- c. Peroneus longus
- d. Peroneus brevis
- e. Gastrocnemius

128) On the front of the ankle joint, the tendon of extensor hallucis longus:

- a. is medial to tibialis anterior
- b. is medial to deep peroneal nerve
- c. is lateral to extensor digitorum longus
- d. possesses no synovial sheath
- e. is lateral to extensor hallucis brevis

129) Regarding lower leg structures

- a. The anterior tibial artery lies lateral to tibialis anterior
- b. The deep peroneal nerve originates in the popliteal fossa
- c. Peroneus longus is supplied by the deep peroneal nerve
- d. Extensor digitorum longus lies medial to tibialis

- anterior
- e. The tibia receives a nutrient vessel from the anterior tibial

Dorsum of foot:

130) The most medial structure passing under the inferior extensor retinaculum of the foot is:

- a. deep peroneal nerve
- b. tibialis anterior
- c. extensor hallucis longus
- d. anterior tibial artery
- e. peroneus brevis

131) All of the following structures pass deep into the superior extensor retinaculum at the ankle except

- a. Extensor digitorum longus
- b. Deep peroneal nerve
- c. Anterior tibial artery
- d. Superficial peroneal nerve
- e. Peroneus tertius

132) Under the extensor retinaculum of the foot the most lateral structure is

- a. Sural nerve
- b. Dorsalis pedis artery
- c. Peroneus tertius
- d. Extensor digitorum longus
- e. Extensor hallucis longus

Posterior compartment of leg:

133) Tibialis anterior and tibialis posterior BOTH:

- a. invert the foot

- b. insert on the tuberosity of the navicular bone
 - c. both
 - d. neither
- 134) The flexor retinaculum of the foot attaches to which two bony structures?
- a. distal ends of tibia and fibula
 - b. calcaneus and medial malleolus
 - c. lateral malleolus and calcaneus
 - d. medial malleolus and talus
 - e. talus and navicular
- 135) Popliteus
- a. Arises from the tibia above the condyles
 - b. Slopes upwards and medially
 - c. Inserts into the lateral meniscus
 - d. Is innervated by a branch of the common peroneal nerve
 - e. Acts to "lock" the knee in full extension
- 136) Tibialis posterior inserts into
- a. The base of the first metatarsal
 - b. The navicular
 - c. Base of 2nd and third metatarsals
 - d. The cuboid
 - e. None of the above
- 137) Which of the following structures does not pass anterior to the medial malleolus
- a. Tibialis anterior
 - b. Extensor hallucis longus
 - c. Flexor hallucis longus

- d. Anterior tibial artery
 - e. Peroneus tertius
- 138) popliteus:
- a. attaches to the medial femoral condyle
 - b. "locks" the knee in extension
 - c. supplied by a branch of the femoral nerve
 - d. lies immediately deep to the popliteal artery
 - e. may attach to the medial meniscus
- 139) Flexor digitorum longus:
- a. crosses deep to tibialis posterior in calf
 - b. crosses superficial to flexor longus hallucis in sole
 - c. is an evor of the foot
 - d. supplied by musculocutaneous nerve
 - e. supplied by L5 nerve root
- 140) Features of the fibula include which of the following?
- a. it is on the medial side of the tibia
 - b. its medial surface is grooved for the origin of tibialis posterior
 - c. it is ossified from five centers
 - d. it does not provide origin for flexor digitorum longus
 - e. its lower third is rough for the origin of soleus
- 141) A ruptured tendo calcaneus ("Achilles tendon") would result in severe impairment or loss of which of the following movements?
- a. plantarflexion
 - b. dorsiflexion
 - c. inversion
 - d. eversion
 - e. rotation

- 142) The gastrocnemius muscle can produce:
- a. plantar flexion of the foot
 - b. flexion of the leg at the knee
 - c. both
 - d. neither
- 143) The soleus muscle:
- a. is innervated by the tibial nerve
 - b. is a plantar flexor of the foot
 - c. both
 - d. neither
- 144) Which muscle is essential to lift the heel off the ground in walking?
- a. peroneus longus
 - b. tibialis anterior
 - c. flexor hallucis longus
 - d. flexor hallucis brevis
 - e. gastrocnemius
- 145) The muscle that contracts to unlock the extended knee joint is the:
- a. popliteus
 - b. plantaris
 - c. medial head of the gastrocnemius
 - d. lateral head of gastrocnemius
 - e. soleus
- 146) Following an injury to the leg, a patient is unable to dorsiflex his foot. Which nerve is most likely to be damaged?
- a. the deep peroneal nerve
 - b. the sural nerve
 - c. the superficial peroneal nerve

- d. the saphenous nerve
- e. none of the above

Sole of foot:

- 147) In the sole of the foot, layer 2 consists of intrinsic muscles and tendons of extrinsic muscles. Therefore, all of the following muscles are considered part of layer 2 except
- a. flexor digitorum longus
 - b. flexor digitorum brevis
 - c. flexor hallucis longus
 - d. quadratus plantae
- 148) Which is not a component of the second layer of the sole of the foot?
- a. Tendon of flexor hallucis longus
 - b. Abductor hallucis
 - c. Flexor accessorius
 - d. The lumbrical muscles
 - e. Tendon of flexor digitorum longus
- 149) Regarding the medial longitudinal arch
- a. Its stability is due to its bony structures
 - b. Flexor hallucis brevis acts as a bowstring
 - c. The plantar aponeurosis is of minimal importance
 - d. Peroneus longus supports the posterior portion
 - e. None of the above
- 150) The third layers of muscles of the sole of the foot include:
- a. four lumbricals and quadratus plantae
 - b. plantar and dorsal interossei
 - c. extensor hallucis longus and extensor digitorum longus

- d. flexor digiti minimi brevis, adductor hallucis transversus, adductor hallucis obliquus and flexor hallucis brevis
- e. abductor hallucis, abductor digiti minimi and flexor digitorum brevis

151) The transverse arch of the foot is formed by:

- a. calcaneus, talus, navicular, cuneiforms and first three metatarsals
- b. Cuneiforms, cuboid, all five metatarsals
- c. calcaneus, cuboid, lateral two metatarsals
- d. calcaneus, navicular, lateral two metatarsals
- e. sustentaculum tali and cuneiforms

152) In the sole:

- a. plantar arteries and nerves lie between second and third layers
- b. the myotome is essentially S1
- c. the medial plantar nerve supplies the medial 4. toes
- d. tendons for peroneus longus and tibialis posterior lie in the third layer
- e. abductor digiti minimi is in the first layer

153) Which of the following is NOT a component of the second layer of the sole of the foot?

- a. tendon of flexor hallucis longus
- b. abductor hallucis
- c. flexor accessorius
- d. the lumbrical muscles
- e. tendon of flexor digitorum longus

Nerves of leg and foot:

- 154) With regard to the cutaneous innervation of the lower limb
- a. Superficial peroneal nerve supplies the first inter-digital cleft
 - b. Sural nerve supplies the medial malleolus
 - c. Deep peroneal nerve supplies the third inter-digital cleft
 - d. The medial plantar nerve supplies a greater area than the lateral
 - e. Branches of the tibial nerve supply much of the dorsum of the foot
- 155) Injury to tibial nerve in the popliteal fossa might result in:
- a. loss of eversion
 - b. (diminished sensation) on dorsal surface of foot
 - c. inability to stand on one's toes
 - d. drop foot
 - e. loss of sensation between the great and second toe
- 156) The only innervation of the femoral nerve below the level of the knee is:
- a. the gracilis muscle
 - b. the tibialis anterior muscle
 - c. the skin of the lateral foot
 - d. the adductor longus
 - e. the skin of the medial leg
- 57) The following nerves which damaged may result in foot drop?
- a. femoral n.
 - b. tibial n.
 - c. common peroneal n.
 - d. obturator n.
 - e. pudendal n.

- 158) Deep peroneal n. supplies
- Tibialis anterior
 - tibialis posterior
 - Peroneus longus
 - Peroneus brevis
 - Gastrocnemius
- 159) The sural nerve is located most closely to the:
- inferior border of the gluteus maximus
 - lateral malleolus
 - great saphenous vein
 - neck of the fibula
 - medial side of the patella
- 160) The best diagnostic test for the anterior compartment syndrome of the leg is to check for loss of cutaneous sensation on the:
- medial part of the plantar surface of the foot
 - lateral part of the plantar surface of the foot
 - dorsum of the foot, between the first and second toes
 - dorsolateral side of the foot and little (5th) toe
 - anterolateral side of the leg
- 161) A sharp blow to the neck of the fibula may result in damage to which structure?
- sciatic nerve
 - common peroneal nerve
 - tibial nerve
 - popliteal nerve
 - none of the above

Joints of lower limb:

Hip joint

- 162) Which statement is false concerning the hip joint?
- a. it is a ball and socket joint
 - b. the entire acetabulum articulates with the femoral head
 - c. the acetabulum is deepened by the acetabular labrum
 - d. the ligamentum teres is attached to the fovea on the femoral head
 - e. the iliofemoral ligament attaches to the intertrochanteric line
- 163) Regarding the hip joint all of the following statements are true EXCEPT:
- a. the anterior fibers of the gluteus medius and minimus act as medial rotator of the hip joint
 - b. the gluteus maximus is the most powerful lateral rotator of the hip joint
 - c. gluteus medius and minimus are supplied by the inferior gluteal nerve (L5, S1, S2)
 - d. the superior gluteal nerve (L4, L5, S1) emerges from the greater sciatic notch
 - e. the iliofemoral ligament limits extension at the hip joint
- 164) Regarding movements of the hip joint
- a. Stability is reduced by the presence of the long femoral neck
 - b. Medial rotation is achieved by obturator externus
 - c. Sartorius is a lateral rotator
 - d. Medial rotation is resisted by the iliotibial tract
 - e. Abduction is not usually required in normal walking

165) The hip joint

- a. Derives its stability largely from the shape of its articular surfaces
- b. Has the ischiofemoral as its strongest ligament
- c. Is only supplied by the obturator and sciatic nerves
- d. Is limited in full extension by the pubofemoral ligament
- e. Is flexed largely by sartorius and rectus femoris

166) Stability for the hip joint is provided chiefly by the

- a. Ligamentum teres
- b. Ischiofemoral ligament
- c. Pubofemoral ligament
- d. Iliofemoral ligament
- e. Transverse ligament

167) Ilio-femoral ligament:

- a. limits hip flexion
- b. limits hip extension
- c. limits hip adduction
- d. limits hip internal rotation
- e. none of the above

168) Concerning the hip joint, the right description is:

- a. is formed by the head of femur and the acetabulum.
- b. is the biaxial joint
- c. the iliofemoral ligament prevents hyperflexion of the thigh
- d. the fracture of femoral neck must be the intracapsula type.
- e. the ischiofemoral ligament block the acetabular notch.

169) Which joint has the articular labrum

- a. inferior tibiofibular joint
- b. superior tibiofibular joint
- c. knee joint
- d. hip joint
- e. ankle joint

170) True regarding the hip joint is:

- a. Medial rotation and abduction is caused by gluteus medius and gluteus minimus
- b. Medial rotation and adduction is caused by gluteus medius and gluteus minimus
- c. Lateral rotators of thigh are supplied by femoral nerve
- d. Hyperextension of hip is prevented by capsular thickening

Knee joint

171) Excessive anterior movement of the tibia when pulling forward on the leg with the knee flexed would indicate damage to this ligament of the knee:

- a. medial collateral
- b. lateral collateral
- c. anterior cruciate
- d. posterior cruciate
- e. oblique popliteal

172) The suprapatellar bursa:

- a. communicates directly with the knee joint cavity
- b. is located superficial to the quadriceps femoris

- muscle
- c. both
- d. neither

173) Anterior cruciate ligament prevents:

- a. Anterior dislocation of tibia
- b. Posterior dislocation of tibia
- c. Anterior dislocation of femur
- d. Posterior dislocation of femur

174) The fibular collateral ligament is:

- a. tested by the application of valgus stress, rather than varus stress, to the knee joint
- b. located superficial to the tendon of the popliteus
- c. attached to the lateral meniscus
- d. located superficial to the biceps femoris tendon
- e. located deep to the iliotibial tract

175) Which joint has the articular disc?

- a. inferior tibufibular joint
- b. superior tibufibular joint
- c. knee joint
- d. hip joint
- e. ankle joint

176) All of the following statements about the knee joint are correct except

- a. the principal movements of the knee joint are flexion and extension
- b. when the knee is fully extended, the anterior cruciate ligament prevents posterior displacement of femur on tibia and hyperextension of the knee

- c. when the knee is fully flexed, the posterior cruciate ligament prevents anterior displacement of the femur on the tibia and stabilizes the knee when walking downstairs
- d. when the knee is fully extended, it "locks" because of lateral rotation of the femur on the tibia
- e. the popliteus muscle contracts to "unlock" the knee so that flexion of the knee can occur

177) The nerve supply to the knee joint comes from:

- a. sciatic
- b. femoral
- c. obturator
- d. all of these
- e. none of these

178) The suprapatellar bursa

- a. Does not communicate with the knee joint
- b. Lies in front of the vastus intermedius muscle
- c. Extends 5cm or more above the patella
- d. Lies deep to the patella retinacula
- e. None of the above

179) When standing, the knee joint is locked in extension by:

- a. lateral rotation of the tibia
- b. medial rotation of the femur
- c. tightening of the medial ligament
- d. tightening of the lateral ligament
- e. tension in the oblique popliteal ligament

180) Which one of the following has an intracapsular tendon?

- a- plantaris

- b- popliteus
- c- rectus femoris
- d- psoas major
- e- peroneal longus

- 181) The anterior cruciate ligament:
- a. lies within the synovial membrane of the knee joint
 - b. attaches from the anterior tibial plateau to the medial condyle of the femur
 - c. prevents forward displacement of the femur on the tibial plateau
 - d. produces lateral rotation of the femur in the 'screw home' position of full extension
 - e. none of the above

- 182) Important stabilizing structures on lateral side of knee are:
- a. tibial collateral ligament; biceps tendon; and iliotibial tract
 - b. fibular collateral ligament; biceps tendon; and iliotibial tract
 - c. fibular collateral ligament; cruciate ligament; and pes anserinus
 - d. tibial collateral ligament; popliteus tendon; and patellar ligament
 - e. fibular collateral ligament; coronary ligament; and transverse ligament

- 183) Which structure is intrasynovial at the knee joint:
- a. oblique popliteal ligament
 - b. tendon of popliteus
 - c. medial and lateral menisci
 - d. anterior cruciate ligament

- e. none of the above

184)

With respect to the ligaments around the knee joint:

- a. the fibular collateral ligament blends with the capsule and is attached to the lateral meniscus
- b. the tibial collateral ligament blends posteriorly with the capsule and is attached to the medial meniscus
- c. the oblique popliteal ligament is an expansion from the tendon of semitendinosus that blends with the capsule posteriorly
- d. the posterior cruciate ligament runs from anterior tibial plateau to the posteromedial aspect of the lateral femoral condyle
- e. the transverse ligament runs posteriorly between menisci

185)

Which of the following structures lies within the knee joint?

- a. patellar ligament
- b. tibial collateral ligament
- c. fibular collateral ligament
- d. tendon of popliteus
- e. patellar retinacula

186)

Regarding the movements at the knee joint:

- a. popliteus 'unlocks' the extended knee by producing medial rotation of the femur
- b. there is no active rotation of the extended knee
- c. passive extension of the knee does not result in 'locking' of the joint
- d. the posterior cruciate ligament prevents backward

- displacement of the femur on the tibial plateau
- e. the major role of the menisci is in flexion/extension of the knee

187) Attached to lateral meniscus is which of the following:

- a. posterior cruciate ligament
- b. lateral ligament of the knee
- c. mucous fold
- d. popliteus muscle
- e. oblique popliteal ligament

188) The knee:

- a- the cruciate ligaments are intra-synovial
- b- the posterior cruciate ligament stabilizes the flexed weight-bearing knee
- c- the posterior horn of the medial meniscus is attached to the intercondylar area of the tibia behind the posterior cruciate ligament
- d- the knee joint normally contains 5.0ml of synovial fluid
- e- active rotation may occur in the extended knee

189) Which of the following bursae is most likely to communicate with the knee joint?

- a. deep infrapatellar bursa
- b. superficial infrapatellar bursa
- c. prepatellar bursa
- d. semimembranosis bursa
- e. suprapatellar bursa

190) Regarding the menisci of the knee

- a. They are attached to the femoral condyles

- b. They have a rich blood supply
- c. The medial meniscus is firmly attached to the capsule of the joint
- d. The medial meniscus gives origin to the menisofemoral ligaments
- e. They are composed of mainly elastic fibers

91) Regarding movements of the knee joint which statement is **INCORRECT**:

- a. Sartorius assists with flexion
- b. The main action of popliteus is unlocking of the knee
- c. Extension occurs by the quadriceps muscle
- d. Locking of the knee is an active process
- e. The hamstrings are flexors for the knee

92) Regarding innervation of flexion/extension of the knee

- a. All muscles of the quadriceps femoris are supplied by L3/4
- b. Vastus medialis and intermedius are supplied by the same branch of the femoral nerve
- c. Flexion of the knee by hamstrings is supplied solely by tibial component of sciatic nerve
- d. Flexor innervation of knee can be tested by knee jerk L3
- e. Traumatic fracture dislocation of hip affects knee extension greater than flexion

93) With regards to the knee joint

- a. The medial collateral ligament extends 8cm below the joint margin
- b. The medial collateral ligament is extra-articular
- c. The tendon of popliteus is intra-articular
- d. The lateral meniscus is more C-shaped
- e. All of the above

194) True statements concerning the anterior cruciate ligament include:

- a. its attachment to the tibial plateau (top of tibia) is anterior to the attachment of the anterior horn of the medial meniscus
- b. it prevents posterior displacement of the femur on the tibia
- c. both
- d. neither

Ankle joint

195) The deltoid ligament :

- a. is attached to the talus, navicular and calcaneus
- b. is torn less commonly than lateral ligament of ankle
- c. both
- d. neither

196) The "spring" ligament attaches to what two bones?

- a. fibula and calcaneus
- b. tibia and calcaneus
- c. tibia and talus
- d. talus and navicular
- e. calcaneus and navicular

197) The ligament usually injured in a hyper-inversion of the foot (sprained ankle) is the:

- a. Anterior tibiotalar
- b. Posterior tibiotalar
- c. Medial collateral
- d. Lateral collateral

198) At the ankle joint:



- a. the distal fibular is not part of the joint
 - b. the medial ligament is made up of three separate bands
 - c. the posterior tibiofibular ligament is also known as posterior transverse ligament
 - d. the peroneus longus and brevis muscles assist in plantar flexion
 - e. nerve supply is from the anterior and posterior tibial nerves
- 199) The deltoid ligament belongs to the:
- a. ankle joint
 - b. hip joint
 - c. knee joint
 - d. talocalcaneonavicular joint
 - e. calcaneocuboid joint
- 200) Concerning the ankle:
- a. movements of dorsi and plantar flexion only are possible
 - b. communicates with inferior tibiofibular joint
 - c. communicates with posterior subtalar joint
 - d. dorsiflexion accompanied by some eversion
 - e. can be plantarflexed by flexor longus hallucis
- 201) The anterior part of the talus articulates with the:
- a. cuboid
 - b. first, second and third cuneiform bones
 - c. fourth and fifth cuneiform bones
 - d. calcaneus
 - e. navicular

Movement of eversion and inversion

- 202) Which muscle can evert the ankle joint?
- a. peroneus longus
 - b. the tibialis anterior
 - c. the tibialis posterior
 - d. the soleus
 - e. flexor digitorum brevis
- 203) What movement occurs at the subtalar joint
- a. Inversion
 - b. Eversion
 - c. dorsiflexion
 - d. Plantarflexion
 - e. Both a and b
- 204) Inversion and eversion of the foot take place MAINLY at
- a. ankle joint
 - b. joints between the talus and calcaneus
 - c. joint between calcaneus and navicular bone
 - d. calcanocuboid joint
 - e. inferior tibiofibular joint
- 205) Inversion of the foot is by which pair of muscles?
- a. peroneus longus and peroneus brevis
 - b. peroneus longus and tibialis posterior
 - c. tibialis anterior and tibialis posterior
 - d. peroneus brevis and plantaris
 - e. none of the above

Answers of lower limb MCQ

1	e	31	a	61	c	91	d	121	c
2	d	32	b	62	d	92	d	122	e
3	d	33	a	63	c	93	c	123	a
4	c	34	c	64	a	94	c	124	a
5	c	35	d	65	e	95	e	125	c
6	d	36	d	66	d	96	e	126	d
7	b	37	c	67	e	97	a	127	a
8	d	38	d	68	e	98	e	128	b
9	b	39	b	69	a	99	d	129	a
10	c	40	b	70	d	100	c	130	b
11	c	41	c	71	b	101	d	131	d
12	c	42	e	72	b	102	a	132	c
13	e	43	b	73	d	103	d	133	a
14	a	44	b	74	b	104	a	134	b
15	e	45	a	75	a	105	a	135	c
16	c	46	d	76	e	106	e	136	b
17	c	47	e	77	c	107	b	137	c
18	e	48	c	78	d	108	c	138	d
19	a	49	b	79	b	109	c	139	b
20	e	50	e	80	e	110	b	140	b
21	e	51	a	81	e	111	a	141	a
22	c	52	c	82	b	112	c	142	c
23	c	53	a	83	a	113	b	143	c
24	d	54	c	84	d	114	b	144	e
25	c	55	c	85	b	115	c	145	a
26	a	56	b	86	d	116	c	146	a
27	b	57	c	87	a	117	d	147	b
28	e	58	d	88	a	118	c	148	b
29	d	59	b	89	c	119	b	149	e
30	e	60	b	90	d	120	a	150	d

171	c	181	e	191	d	201	e
172	a	182	b	192	a	202	a
173	a	183	e	193	e	203	e
174	b	184	b	194	b	204	b
175	e	185	d	195	c	205	c
176	d	186	b	196	e		
177	d	187	d	197	d		
178	c	188	b	198	d		
179	b	189	e	199	a		
180	b	190	c	200	e		

171	c	181	e	191	d	201	e
172	a	182	b	192	a	202	a
173	a	183	e	193	e	203	e
174	b	184	b	194	b	204	b
175	e	185	d	195	c	205	c
176	d	186	b	196	e		
177	d	187	d	197	d		
178	c	188	b	198	d		
179	b	189	e	199	a		
180	b	190	c	200	e		