

Musculoskeletal Anatomy

M.C.Q



By

Mohey Hulail

***Professor of Anatomy & Embryology
Zagazig University***

Musculoskeletal Anatomy

MCQ



By
Mohey Hulail
Professor of anatomy
& embryology
Zagazig University

Contents

- Upper limb	1
- Pectoral region	1
- Axilla & brachial plexus	3
- Shoulder region	9
- Arm	13
- Forearm	26
- Hand, joints & nerve injury.....	31
- Answers of upper limb MCQ	39
- Lower limb	40
- Anterior compartment of the thigh	40
- Gluteal region	56
- Back of thigh	63
- The leg	67
- Dorsum of foot	73
- Sole of foot	77
- Hip joint	81
- Answers of lower limb MCQ	94
- Head & Neck	95
- Scalp	95
- The Face	97
- Triangles.....	99
- Temporal and Infratemporal fossa	107
- Answers of Head and neck MCQ	111
- Lymphatic system	112
- Answers of lymphatic system MCQ	122

Upper limb
Pectoral region

- 1) These muscles are attached to the medial two thirds of the clavicle EXCEPT:
 - a. Sternomastoid.
 - b. Deltoid.
 - c. Pectoralis major.
 - d. Subclavius.
 - e. Sternohyoid.
- 2) The deltopectoral groove contains:
 - a. Deltoid branch of the lateral thoracic artery.
 - b. Cephalic vein.
 - c. Infraclavicular lymph nodes.
 - d. All of the above.
 - e. None of the above.
- 3) What structure does NOT pierce the clavipectoral fascia?
 - a. cephalic vein
 - b. lymphatics
 - c. lateral thoracic artery
 - d. lateral pectoral nerve
 - e. thoracoacromial artery
- 4) About Pectoralis major which is false:
 - a. is innervated by the medial and lateral pectoral nerves
 - b. inserts into the medial lip of the intertubercular sulcus
 - c. can act as an accessory muscle of inspiration
 - d. acts with latissimus dorsi to draw the body upwards
 - e. shares in formation of anterior wall of axilla
- 5) The subclavius muscle:

- a. arises from the costochondral junction of the second rib
 - b. passes in the subclavian groove on the inferior aspect of the clavicle to attach to the acromion
 - c. is enclosed by the upper attachment of the clavipectoral fascia
 - d. is supplied by C7 and C8
 - e. assists shoulder adduction
- 6) The nerve supply of pectoralis minor is from the:
- a. Medial pectoral nerve
 - b. thoracodorsal nerve
 - c. axillary nerve
 - d. long thoracic nerve
 - e. dorsal scapular nerve
- 7) Pectoralis major arises from all EXCEPT the:
- a. lateral anterior half of manubrium
 - b. body of sternum
 - c. aponeurosis of external oblique
 - d. upper six ribs
 - e. medial half of anterior clavicle
- 8) Paralysis of the serratus anterior muscle causes:
- a. Winging of the scapula.
 - b. Claw hand.
 - c. Ape hand.
 - d. Wrist drop.
 - e. Policeman's tip position.
- 9) Which nerve supplies serratus anterior?
- a. axillary nerve (C5, C6)
 - b. long thoracic nerve (C5, C6, C7)

- c. musculocutaneous nerve (C5, C6)
- d. thoracodorsal nerve (C6, C7, C8)
- e. suprascapular nerve (C5, C6)

Axilla & brachial plexus

10) The axilla contains all of the following EXCEPT:

- a. Axillary artery.
- b. Axillary vein.
- c. Trunks of the brachial plexus.
- d. Fat.
- e. Tail of the breast.

11) The brachial plexus has:

- a. 8 rami (roots).
- b. 5 trunks.
- c. 6 divisions.
- d. 4 cords.

12) The usual number of branches of the medial cord of the brachial plexus are:

- a. 1.
- b. 2.
- c. 3.
- d. 4.
- e. 5.

13) The anterior wall of the axilla is formed by all of the following EXCEPT:

- a. Pectoralis major.
- b. Teres major.
- c. Pectoralis minor.
- d. Subclavius.

- e. Clavipectoral fascia.
- 14) In Erb's paralysis, the deformity is called:
- a. Winging of the scapula.
 - b. Claw hand.
 - c. Ape hand.
 - d. Wrist drop.
 - e. Policeman's tip position.
- 15) Paralysis of the serratus anterior muscle causes:
- a. Winging of the scapula.
 - b. Claw hand.
 - c. Ape hand.
 - d. Wrist drop.
 - e. Policeman's tip position.
- 16) The suprascapular nerve is a branch from the:
- a. Roots of the brachial plexus.
 - b. Upper trunk.
 - c. Middle trunk.
 - d. Posterior cord.
 - e. Lateral cord.
- 17) The root value of the long thoracic nerve is:
- a. C5.
 - b. C6.
 - c. C7.
 - d. C5, 6 & 7
 - e. C5, 6, 7 & 8
- 18) Regarding the brachial plexus, which is INCORRECT?
- a. it is derived from the anterior rami of C5-T1

- b. the roots give off the long thoracic nerve to serratus anterior
 - c. the trunks give off one branch
 - d. the cords are divided into anterior and posterior divisions
 - e. the lateral cord supplies part of the median nerve
- 19) As it emerges from the axilla, the median nerve lies where with regards to the brachial artery?
- a. lateral
 - b. anterior
 - c. medial
 - d. posterior
- 20) Regarding the brachial plexus, which is FALSE?
- a. the nerve to serratus anterior comes from the anterior rami of C8-T1
 - b. the suprascapular nerve arises from C5 anterior ramus
 - c. the divisions are at the level of the clavicle
 - d. the thoracodorsal nerve arises from the posterior cord
 - e. the radial nerve is the largest branch of the posterior cord
- 21) Regarding the brachial plexus:
- a. the five roots lie in cervicoaxillary canal
 - b. the anterior division of the lower trunk runs on as the medial cord
 - c. the suprascapular nerve is the only branch from the trunks
 - d. the radial nerve is a branch of the lateral cord
 - e. the ulnar nerve is a branch of the posterior cord.
- 22) Regarding the brachial plexus:
- a. the dorsal scapular nerve is the only branch from the

- trunks
- b. the axillary nerve is derived from C5, 6, 7, 8, T1
 - c. the ulnar nerve is the largest branch of the whole plexus
 - d. the axillary nerve passes below teres major
 - e. the thoracodorsal nerve supplies latissimus dorsi

2

23) With respect to the brachial plexus:

- a. it consists of the entire anterior rami of C5 → T1
- b. there are five roots, three trunks, five divisions and three cords
- c. there are two branches from the divisions
- d. there are a total of 13 branches from the cords
- e. there are two branches from the roots

24) With respect to the brachial plexus:

- a. the lateral cord lies in the axilla
- b. the median nerve is the largest branch of the whole plexus
- c. the cords lie within the posterior triangle of the neck
- d. the ulnar nerve is a branch of the lateral cord
- e. the musculocutaneous nerve is a branch of the medial cord

25) Regarding the brachial plexus:

- a. the medial cord has three branches
- b. the lateral cord has fibres from C7, C8 and T1
- c. the musculocutaneous nerve is a branch of the medial cord
- d. the five roots are found from the posterior rami and C5 – T1 nerves
- e. none of the above

26) The nerve supply of serratus anterior is:

- a. axillary nerve
- b. C4
- c. dorsal scapular nerve
- d. thoracodorsal nerve
- e. long thoracic nerve

27) Which of the following statements about the brachial plexus is INCORRECT?

- a. long thoracic nerve arises from roots
- b. Erb's palsy occurs following traction to upper roots and trunk
- c. posterior cord has five branches
- d. suprascapular nerve arises from trunks
- e. trunks lie behind clavicle on first rib

28) Regarding the axilla:

- a. the posterior wall consists of subscapularis and teres minor
- b. the medial wall is formed by serratus anterior down to the 4th intercostal space
- c. the apex communicates directly with the posterior triangle of the neck
- d. coracobrachialis forms part of the anterior wall
- e. the axillary artery is generally described as having eight branches

29) Which nerve is a branch of the lateral cord of the brachial plexus?

- a- upper subscapular nerve
- b- thoracodorsal nerve
- c- musculocutaneous nerve
- d- ulnar nerve
- e- lower subscapular nerve

30) A nerve arising directly from the upper trunk of the brachial plexus is the:

- a. dorsal scapular
- b. thoracodorsal
- c. suprascapular
- d. upper subscapular
- e. lower subscapular

c.

d.

e.

31) A patient has been thrown from a motorcycle, landing on the shoulder such that the neck and shoulder are widely separated. You suspect an upper brachial plexus injury. What signs do you expect?

34) With

wro

a

t

- a. "claw hand"
- b. paralysis of flexor carpi ulnaris, flexor digitorum superficialis, and flexor digitorum profundus
- c. adducted shoulder, medially rotated arm, and extended elbow
- d. loss of sensation in the medial forearm
- e. "wrist-drop"

35) V

32) A patient exhibits "claw hand." What might have happened?

- a. upper brachial plexus injury
- b. acute brachial plexus neuritis
- c. compression of the cords of the brachial plexus
- d. lower brachial plexus injury
- e. damage to dorsal rami that send fibers to the brachial plexus

33) Brachial plexus:

- a. the thoracodorsal nerve is a branch of the lateral cord
- b. the anterior division of the lower trunk runs on as the medial cord

36

- c. the suprascapular nerve is the only branch from the trunks
- d. the radial nerve is a branch of the lateral cord
- e. the ulnar nerve is a branch of the posterior cord.

34) With respect to the brachial plexus, which of the following is wrong?

- a. it consists of the ventral rami of C5 → T1
- b. there are five roots, three trunks, six divisions and three cords
- c. there are two branches from the divisions
- d. there are a total of 13 branches from the cords
- e. there are two branches from the roots

35) With respect to the brachial plexus:

- a. the lateral cord lies in the cervicoaxillary canal
- b. the median nerve is the largest branch of the whole plexus
- c. the trunks lie within the posterior triangle of the neck
- d. the ulnar nerve is a branch of the lateral cord
- e. the musculocutaneous nerve is a branch of the medial cord

Shoulder region

36) The glenoid cavity articulates with the head of the humerus by a:

- a. Fibrous articulation.
- b. Cartilaginous articulation.
- c. Plane synovial articulation.
- d. Ball and socket synovial articulation.
- e. Hinge synovial articulation.

37) One of the following muscles is not a rotator cuff muscle:

- a. Supraspinatus.
- b. Infraspinatus.
- c. Teres minor.
- d. Teres major.
- e. Subscapularis.

38) Paralysis of the deltoid muscle causes:

- a. Loss of abduction of the arm from zero to 15° .
- b. Loss of abduction of the arm from $15^\circ - 90^\circ$.
- c. Loss of abduction of the arm from zero to 90° .
- d. Loss of abduction of the arm from $90^\circ - 180^\circ$.
- e. None of the above.

39) The axillary nerve supplies the:

- a. Muscle that abducts the arm from zero to 15° .
- b. Muscle that abducts the arm from $15^\circ - 90^\circ$.
- c. Muscle that forms the rounded contour of the shoulder.
- d. a and c.
- e. b and c.

40) Lesion of the axillary nerve causes:

- a. Winging of the scapula.
- b. Flat shoulder.
- c. Claw hand.
- d. Ape hand.
- e. Wrist drop.

41) Regarding the shoulder joint, which is INCORRECT?

- a. full abduction requires medial rotation of the scapula
- b. the long head of biceps is intracapsular
- c. subscapularis muscle rotates the shoulder medially

- d. flexion involves pectoralis major, deltoid, coracobrachialis and biceps
- e. supraspinatus initiates abduction

42) Teres major:

- a. receives its nerve supply from the lateral cord of the brachial plexus
- b. is part of the medial wall of the axilla
- c. forms the medial border of the triangular space
- d. is separated from teres minor by the long head of triceps
- e. has the same nerve supply as teres minor

43) Teres major:

- a. is separated from subscapularis by the quadrangular space in the posterior wall of the axilla
- b. is supplied by the lateral subscapular nerve
- c. inserts into the lateral lip of the intertubercular groove of the humerus
- d. acting alone, it abducts and medially rotates the humerus at the shoulder joint
- e. arises from the ventral aspect of lateral border of the scapula

44) The shoulder joint:

- a. subscapularis, supraspinatus, infraspinatus and teres minor form the rotator cuff of the shoulder
- b. synovial joint of ellipsoid variety
- c. the joint capsule is attached to the surgical neck of the humerus inferiorly
- d. the articular surfaces are the head of the humerus and subscapular fossa
- e. anterior dislocation is very common

45) The nerve supply to deltoid is from the:

- a. pectoral nerves
- b. thoracodorsal nerve
- c. axillary nerve
- d. long thoracic nerve
- e. dorsal scapular nerve

46) The nerve supply to supraspinatus is from the:

- a. lower subscapular nerve
- b. dorsal scapular nerve
- c. suprascapular nerve
- d. upper subscapular nerve
- e. thoracodorsal nerve

47) The nerve supply to teres major is from the:

- a. Lower subscapular nerve
- b. Dorsal scapular nerve
- c. Suprascapular nerve
- d. Upper subscapular nerve
- e. Thoracodorsal nerve

48) Deltoid:

- a. anterior fibres arise from lateral two thirds of clavicle
 - b. posterior fibres arise from superior lip of crest of scapular spine
 - c. middle fibres abducts the shoulder
 - d. supplied by suprascapular nerve
 - e. the only muscle that abducts the shoulder
- 49) A 45-year-old man is unable to initiate abduction of the arm following reduction of a dislocated shoulder. Damage to which nerve is most likely to be responsible for that:

- a. the suprascapular nerve
- b. the long thoracic nerve
- c. the radial nerve
- d. the infrascapular nerve
- e. the axillary nerve

50) The nerve supply of infraspinatus is the:

- a. axillary nerve
- b. long thoracic nerve
- c. infrascapular nerve
- d. **suprascapular nerve**
- e. subscapular nerve

51) Rotator cuff muscles include:

- a. subclavius
- b. **teres minor**
- c. pectoralis major
- d. teres major
- e. deltoid

52) The quadrangular space between teres major and subscapularis transmits:

- a. lower subscapular nerve
- b. axillary artery
- c. profunda brachii vessels
- d. radial nerve
- e. **post circumflex humeral artery**

53) Which of the following statements about the glenohumeral (shoulder joint) is INCORRECT?

- a. long tendon of biceps is extrasynovial
- b. **innervated by medial pectoral nerve**

- c. branches of third part of axillary artery provide majority of blood supply
- d. Gleno-humeral ligament are three weak bands in front of the capsule
- e. The head of humerus is more than half sphere

54) "Winging" of the scapula is most likely caused by which of the following?

- a. **a lesion to the long thoracic nerve**
- b. a lesion to the thoracodorsal nerve
- c. injury to the suprascapular nerve
- d. damage to the dorsal scapular nerve
- e. damage to the upper and lower subscapular nerves

55) The axillary nerve is damaged. What is the likely result?

- a. The teres major atrophies.
- b. **The rounded contour of the shoulder disappears.**
- c. A loss of sensation may occur in the lateral forearm.
- d. The patient may lose the ability to adduct the arm.
- e. The patient may exhibit "wrist-drop."

56) The quadrangular space is bounded by all of the following EXCEPT:

- a. Teres minor.
- b. Teres major.
- c. **Long head of biceps.**
- d. Long head of triceps.
- e. Surgical neck of the humerus.

57) The lesser tuberosity of the humerus gives attachment to the:

- a. Supraspinatus.
- b. **Infraspinatus.**

- c. Teres minor.
- d. Teres major.
- e. **Subscapularis.**

Muscles of the back

58) One of the following is not attached to the medial border of the scapula:

- a. Levator scapulae.
- b. **Teres minor.**
- c. Serratus anterior.
- d. Rhombideus minor.
- e. Rhomboideus major.

59) The trapezius muscle is supplied by the:

- a. Cranial accessory nerve.
- b. **Spinal accessory nerve.**
- c. Long thoracic nerve.
- d. Dorsal scapular nerve.
- e. Thoracodorsal nerve.

60) Actions of the latissimus dorsi:

- a. Flexes the arm.
- b. Abducts the arm.
- c. Laterally rotates the arm.
- d. All of the above.
- e.. **None of the above**

61) Identify the incorrect innervation:

- a. subclavius – own nerve from the brachial plexus
- b. serratus anterior – long thoracic nerve
- c. clavicular head of pectoralis major – medial pectoral nerve
- d. **latissimus dorsi – dorsal scapular nerve**

- e. trapezius – accessory nerve
- 62) The nerve supply of latissimus dorsi is from the:
- a) pectoral nerves
 - b) thoracodorsal nerve**
 - c) axillary nerve
 - d) long thoracic nerve
 - e) dorsal scapular nerve
- 63) The nerve supply of rhomboid major and rhomboid minor is from the:
- a. pectoral nerves
 - b. thoracodorsal nerve**
 - c. axillary nerve
 - d. long thoracic nerve
 - e. dorsal scapular nerve**
- 64) The insertion of latissimus dorsi in the arm is into the:
- a. lateral lip of the intertubercular groove
 - b. medial lip of the intertubercular groove
 - c. posterior border of the proximal humerus
 - d. floor of the intertubercular groove**
 - e. superior facet of the greater trochanter
- 65) A patient cannot raise the trunk (as in climbing). What is most likely the problem?
- a. damage to the ventral rami of C5–C6–C7
 - b. paralysis of the latissimus dorsi**
 - c. injury to the dorsal scapular nerve
 - d. damage to the dorsal rami of C8–T1
 - e. injury to the axillary nerve

Arm

66) The surgical neck of the humerus is related to the:

- a. Radial nerve.
- b. Axillary nerve.
- c. Ulnar nerve.
- d. Median nerve.
- e. None of the above.

67) The back of the medial epicondyle is related to the:

- a. Radial nerve.
- b. Axillary nerve.
- c. Ulnar nerve.
- d. Median nerve.
- e. None of the above.

68) The bicipital groove of the humerus lodges (contains) the:

- a. Axillary artery.
- b. Long head of the biceps.
- c. Long head of the triceps.
- d. Coracobrachialis.

69) All the muscles attached to the bicipital (intertubercular) groove:

- a. Abduct the arm (shoulder joint).
- b. Adduct the arm.
- c. Medial rotation of the arm.
- d. a and c.
- e. b and c.

70) Regarding the biceps brachii, one is INCORRECT:

- a. Has long head attached to the supraglenoid tubercle.
- b. Has short head attached to the tip of the coracoid process.

- c. Has a tendon inserted into the radial tuberosity.
- d. It flexes the elbow joint.
- e. It is powerful supinator of the extended elbow.

71) The bicipital aponeurosis passes obliquely deep to the:

- a. Brachial artery.
- b. Median cubital vein.
- c. Median nerve.
- d. Radial nerve.
- e. None of the above.

72) The anterior compartment of the arm contains all of the following EXCEPT:

- a. Biceps.
- b. Triceps.
- c. Brachialis.
- d. Coracobrachialis.
- e. Brachial artery.

73) Muscles innervated by the musculocutaneous nerve:

- a. Brachialis.
- b. Coracobrachialis.
- c. Biceps.
- d. All of the above.
- e. Triceps.

74) The constriction below the greater and lesser tuberosities of the humerus is called:

- a. Anatomical neck.
- b. Surgical neck.
- c. Humeral shaft.
- d. Humeral ring.

e. None of the above.

75) The lesser tuberosity of the humerus gives attachment to:

- a. Supraspinatus.
- b. Infraspinatus.
- c. Teres minor.
- d. Teres major.
- e. Subscapularis.

76) Which muscle does not insert in or next to the intertubercular groove of the upper humerus?

- a. pectoralis major
- b. pectoralis minor
- c. latissimus dorsi
- d. teres major

77) Which of the following has some nerve supply from the radial nerve?

- a. long head of biceps
- b. coracobrachialis
- c. short head of biceps
- d. brachialis

78) Regarding triceps:

- a. lateral head arises from humerus below radial groove
- b. midshaft humerus fractures may paralyse triceps
- c. has an insertion into the elbow joint capsule
- d. aids in adduction of shoulder
- e. is pierced by ulnar nerve

79) The musculocutaneous nerve passes through which muscle?

- a. coracobrachialis

- b. short head of biceps
- c. brachialis
- d. medial head of triceps
- e. long head of biceps

80) Regarding the nerves and vessels of the arm:

- a. the circumflex vessels and the axillary nerve enter the extensor compartment in the axilla through the triangular space
- b. the ulnar nerve pierces the lateral intermuscular septum in the lower third of the arm
- c. the median nerve crosses obliquely in front of the radial artery in the arm
- d. supratrochlear lymph nodes lie in the subcutaneous fat just above the medial epicondyle
- e. the basilic vein is lateral to the cephalic vein

81) The radial nerve gives rise to all the following except the:

- a. posterior cutaneous nerve of the arm
- b. lower lateral cutaneous nerve of the arm
- c. posterior cutaneous nerve of the forearm
- d. posterior interosseous nerve
- e. medial cutaneous nerve of the arm

82) Which of the following is INCORRECT? The biceps:

- a) is a powerful flexor of the elbow
- b) supinates the forearm
- c) is paralysed by an injury to the posterior cord of the brachial plexus
- d) is a weak flexor of the shoulder
- e) inserts into the bicipital aponeurosis

83) Regarding the humerus:

- a- the head forms half a sphere
- b- the lateral epicondyle is more prominent
- c- the axillary nerve winds around the anatomical neck
- d- the greater tuberosity continues distally as the medial lip of the intertubercular groove
- e- the capsule of the shoulder joint extends down the medial side of the humeral shaft

84) Brachialis is supplied by:

- a. median nerve
- b. musculocutaneous nerve
- c. radial nerve
- d. median and musculocutaneous nn (?nerves)
- e. musculocutaneous and radial nn (?nerves)

85) Long head of triceps:

- a. lies between teres minor and teres major
- b. is supplied by branch of radial nerve at the humeral groove
- c. arises from the humerus above the radial groove
- d. converges with the medial head
- e. arises from the supraglenoid rim

86) The medial cutaneous nerve of the arm:

- a. has fibres from C8 and T1
- b. arises from the dorsal cord of the brachial plexus
- c. pierces the deep fascia adjacent to the tendon of biceps
- d. has communications with the circumflex nerve
- e. none of the above

87) Paralysis of which nerve results in inability to initiate abduction

of the arm?

- a. the axillary nerve
- b. the suprascapular nerve
- c. the subscapular nerve
- d. the dorsal scapular nerve
- e. the thoracodorsal nerve

88) Brachialis muscle:

- a. is innervated by the median nerve
- b. arises from the lower half of the humerus
- c. is both a flexor and extensor of the elbow joint
- d. attaches to the bicipital tuberosity of the radius
- e. has no relation to the medial intermuscular septum

89) The musculocutaneous nerve:

- a. supplies brachioradialis
- b. terminates as the posterior interosseous nerve
- c. supplies all of the brachialis muscle
- d. contains fibres from C5, C6, C7 and C8
- e. arises from the lateral cord of the brachial plexus

90) Biceps:

- a. long head arises from infraglenoid tubercle
- b. is supplied by a branch of the medial cord of the brachial plexus
- c. the two bellies do not merge
- d. is a pronator of the forearm
- e. short head arises from coracoid process, lateral to coracobrachialis

91) Which of the following pierce the medial intermuscular septum of the arm?

- a. medial cutaneous nerve of the arm
- b. ulnar nerve
- c. median nerve
- d. basilic nerve
- e. profunda brachii artery

92) Which of the following parts of the humerus is matched correctly with the nerve with which it is in direct contact?

- a- distal end of humerus . . . radial nerve
- b- surgical neck . . . musculocutaneous nerve
- c- radial groove . . . musculocutaneous nerve
- d- medial epicondyle . . . ulnar nerve

93) Which of the following is true in respect to ulnar nerve injuries?

- a. The injury often occurs where the nerve passes posterior to the medial epicondyle of the humerus.
- b. The patient experiences numbness and tingling on the lateral part of the palm and the thumb.
- c. The patient may exhibit "waiter's tip hand."
- d. Patients have difficulty because they cannot flex their first, second, and third digits at the DIP joints.
- e. Power of abduction is impaired, and when the patient attempts to flex the wrist, the flexor carpi ulnaris brings the hand to the medial side.

94)

in movements of the arm, the coracobrachialis assists in:

- a. Flexion.
- b. Extension.
- c. Abduction.

- d. Rotation.
- e. Supination

- 95) Regarding the bicipital aponeurosis, one is incorrect:
- a. It is an extension from the tendon of the biceps.
 - b. It blends with the deep fascia covering the extensor muscles of the forearm.
 - c. Passes over the brachial artery.
 - d. Crosses the front of the median nerve.
 - e. Separates the median nerve from the median cubital vein

Cubital fossa

- 96) The cubital fossa is bounded by the:

- a. Pronator quadratus.
- b. Brachioradialis.
- c. Coracobrachialis.
- d. Biceps tendon.
- e. a and b only.

- 97) The cubital fossa contains all of the following EXCEPT:

- a. Brachial artery.
- b. Radial artery.
- c. Radial nerve.
- d. Ulnar nerve.
- e. Median nerve.

- 98) At the cubital fossa:

- a. the median nerve is medial to the brachial artery
- b. the biceps tendon separates the brachial artery from the median nerve
- c. the radial artery leaves the brachial artery at right angles

- d. the medial cutaneous nerve of the forearm runs deep to the bicipital aponeurosis
 - e. pronator teres forms the floor of the fossa
- 99) In the cubital fossa:
- a. the floor is formed mainly by the supinator
 - b. the medial wall is formed by pronator teres
 - c. the ulnar nerve lies medially
 - d. the radial nerve lies medial to the biceps tendon
 - e. none of the above
- 100) The cubital fossa:
- a- is bounded by pronator teres, brachioradialis and the tendon of biceps
 - b- the brachial artery lies lateral to the radial nerve
 - c- has a floor made up of brachialis and supinator
 - d- contains the median basilic vein
 - e- the radial nerve lies medial to the posterior interosseous nerve
- 101) which of the following DOES NOT form the boundaries of the cubital fossa:
- a) anconeus
 - b) brachioradialis
 - c) pronator teres
 - d) deep fascia of the forearm
 - e) bicipital aponeurosis
- 102) In the antecubital fossa:
- a. the ulnar nerve is on the medial side
 - b. the median nerve is lateral to the brachial artery
 - c. the radial nerve is on the lateral side
 - d. all the superficial veins are deep to the cutaneous nerves

- e. the brachial artery is lateral to the tendon of biceps
- 103) Which of the following structures is NOT one of the contents of the cubital fossa?
- a. brachial artery
 - b. tendon of biceps brachii
 - c. radial nerve
 - d. anterior interosseous nerve
 - e. median nerve

- 104) Which of the following DOES NOT form one of the boundaries of the cubital fossa?
- a. brachialis
 - b. supinator
 - c. line connecting the two humeral epicondyles
 - d. bicipital aponeurosis
 - e. medial margin of pronator teres

Forearm

- 105) The lateral cutaneous nerve of the forearm is derived from the:
- a. C5.
 - b. C6.
 - c. C7.
 - d. C5, 6, 7
 - e. C6, 7, 8
- 106) The median nerve supplies all of the following muscles EXCEPT:
- a. Flexor carpi ulnaris.
 - b. Flexor carpi radialis.
 - c. Flexor digitorum superficialis.

- d. Palmaris longus.
 - e. Pronator teres.
- 107) The ulnar nerve supplies all of the following muscles EXCEPT:
- a. Adductor pollicis.
 - b. Flexor pollicis longus.
 - c. Interossei muscles.
 - d. Hypothenar muscles.
 - e. Medial two lumbricals.
- 108) The ulnar nerve:
- a. is a branch from the lateral cord of the brachial plexus.
 - b. Carries fibers from C8 and T1.
 - c. Supplies all the superficial muscles of the front of the forearm except flexor carpi ulnaris.
 - d. All of the above.
 - e. None of the above.
- 109) The following muscles are innervated by the median nerve EXCEPT:
- a. Pronator teres.
 - b. Pronator quadratus.
 - c. Flexor pollicis longus.
 - d. Opponens pollicis.
 - e. Adductor pollicis.
- 110) Muscles taking origin from the radius:
- a. Biceps.
 - b. Supinator.
 - c. Pronator quadratus.
 - d. Flexor pollicis longus.
 - e. None of the above.

- 111) Which is NOT an origin of supinator
- a. radial tuberosity
 - b. radial collateral ligament
 - c. lateral epicondyle of humerus
 - d. supinator crest of ulnar
 - e. aponeurosis overlying supinator muscle
- 112) Which muscle does not arise from the common extensor origin?
- a. extensor carpi radialis brevis
 - b. extensor digitorum
 - c. extensor digiti minimi
 - d. extensor carpi radialis longus
 - e. extensor carpi ulnaris
- 113) In the forearm:
- a. the brachial artery lies medial to the median nerve in the cubital fossa
 - b. the radial artery lies lateral to the radial nerve in the middle third
 - c. the anterior interosseous nerve arises from the radial nerve
 - d. the median nerve passes between the two heads of flexor carpi ulnaris
 - e. the posterior interosseous nerve passes between the two layers of the supinator
- 114) Which of the following IS NOT true about the flexor carpi ulnaris?
- a. acts as a synergist during wrist extension

- b. ulnar artery passes deep to its deep head
- c. inserts into base of fifth metacarpal
- d. most medial muscles arising from the common flexor origin
- e. ulnar nerve passes between its two heads

115) Radial nerve:

- a. is the largest branch of the brachial plexus
- b. occupies the length of the spiral groove
- c. damage causes inability to extend the wrist and the interphalangeal joints
- d. damage causes sensory loss to the palm
- e. arises from the lateral cord of the brachial plexus

116) Which of the following is CORRECT concerning flexor digitorum profundus?

- a. it arises from both radius and ulna
- b. is innervated by both median and ulnar nerves
- c. the origins of the lumbricals arise from its tendons
- d. it flexes the terminal phalangeal joints

117) Regarding flexor digitorum profundus, which is NOT TRUE?

- a. is the most powerful and bulky muscle of the forearm
- b. arises from the common flexor origin
- c. the tendon for the index separates in the forearm from the three other tendons
- d. it gives rise to origin of four lumbricals
- e. it is supplied by the anterior interosseous nerve and the ulnar nerve in a variable distribution

118) Which of the following DOES NOT contribute to the innervation of the elbow joint?

- a. radial nerve
- b. ulnar nerve
- c. median nerve
- d. musculocutaneous nerve
- e. medial cutaneous nerve of the arm

119) Which of the following muscles arises, in part, from the radial collateral ligament of the elbow joint?

- a. extensor carpi radialis brevis
- b. supinator
- c. extensor digiti minimi
- d. extensor carpi ulnaris
- e. extensor digitorum

120) In the forearm:

- a. the median nerve passes between the two heads of pronator teres
- b. the ulnar nerve lies deep to flexor digitorum profundus
- c. superficial fibres of flexor digitorum superficialis gives rise to the tendons for index and middle fingers
- d. the radial artery is on the lateral side of the radial nerve
- e. the fibres of the interosseous membrane run obliquely down from ulna to radius

121) What structure does NOT lie in the anatomical snuff box?

- a. cephalic vein
- b. radial artery
- c. radial styloid
- d. first metacarpal bone
- e. extensor pollicis longus

Hand, joints & nerve injury

- 122) The proximal row of the carpal bones consists of all of the following EXCEPT:
- Scaphoid.
 - Lunate.
 - Trapezium.
 - Pisiform.
 - Triquetral.
- 123) Regarding the ulnar nerve, all are correct EXCEPT:
- Is the largest branch of the medial cord of the brachial plexus.
 - Carries fibers from C6, C7 and C8.
 - Have no branches in the arm.
 - It enters the hand superficial to the flexor retinaculum.
 - Injury of the nerve causes partial claw hand.
- 124) Regarding the median nerve, all are correct EXCEPT:
- Arises from both the medial and lateral cords of the brachial plexus.
 - It crosses the brachial artery at the insertion of the coracobrachialis.
 - In the cubital fossa, it lies lateral to the brachial artery.
 - It enters the hand in the carpal tunnel.
 - Injury of the nerve causes ape-like hand.
- 125) The distal row of the carpal bones consists of:
- Scaphoid.
 - Lunate.
 - Triquetral.
 - Pisiform.
 - None of the above.

- 126) Structure crossing superficial to the flexor retinaculum:
- Median nerve.
 - Ulnar nerve.
 - Flexor digitorum superficialis.
 - Flexor pollicis longus.
 - None of the above.
- 127) Carpal tunnel syndrome causes:
- Claw hand.
 - Wrist drop.
 - Paralysis of all muscles of the hand.
 - Ape-like hand.
 - Waiter's tip position.
- 128) Relations of the median nerve at the wrist:
- Lies in the carpal tunnel.
 - 2 inches above the wrist, it becomes subcutaneous.
 - Lies between the tendons of the palmaris longus and flexor carpi radialis.
 - All of the above.
 - A and B only.
- 129) As a result of injury of the ulnar nerve at the wrist:
- Marked wasting (atrophy) of the thenar eminence.
 - The thumb cannot be abducted.
 - The thumb cannot be adducted.
 - Loss of sensation over the thumb.
 - A and B are correct.
- 130) At the wrist / carpal tunnel:
- flexor retinaculum attaches to scaphoid and trapezoid laterally
 - all superficial and deep flexors, except flexor pollicis

- longus, share a common flexor sheath
- c. the ulnar nerve lies medial to pisiform
- d. tendons of flexor digitorum remain attached until they reach the palm
- e. median nerve lies beneath flexor retinaculum between flexor digitorum superficialis and flexor pollicis longus

131) The elbow joint:

- a. is supplied exclusively by the radial nerve
- b. permits flexion/extension and pronation/supination
- c. has a capsule which attaches in part to the radius
- d. communicates with the superior radioulnar joint
- e. none of the above

132) The carpus:

- a. there are five extensor tunnel compartments
- b. the extensor expansion attaches to the anterolateral border of the radius above the styloid process, and distally to the pisiform and trapezoid bones
- c. the radiocarpal joint is an uniaxial synovial joint
- d. there are no muscular attachments to the extensor surface of the carpus
- e. the bones palpable in the anatomical snuff box are the radial styloid, triquetral, scaphoid and base of first metacarpal

133) What structure pierces the flexor retinaculum?

- a. ulnar nerve
- b. flexor carpi radialis tendon
- c. palmar branch of median nerve
- d. flexor pollicis longus tendon
- e. ulnar artery

- 134) The lumbrical muscles:
- arise from the four superficial tendons
 - are all supplied by median nerve
 - flex the metacarpophalangeal joint
 - attach directly to the first phalanx
- 135) Regarding the extensor retinaculum:
- septa divide the extensor region into four compartments
 - it attaches to the ulnar styloid
 - it is a thickening of the deep fascia on the back of the wrist joint
- 136) The median nerve goes through the heads of which muscle?
- biceps brachii
 - supinator
 - pronator teres
 - pronator quadratus
 - flexor digitorum superficialis
- 137) A high ulnar nerve injury might produce:
- weakness of elbow flexion
 - a 'claw hand'
 - weak abduction of the index finger
 - triceps paralysis
 - sensory loss over the radial three fingers
- 138) Regarding the anatomical snuffbox:
- the cephalic vein begins in the roof
 - the extensor pollicis brevis tendon forms the ulnar side
 - the trapezoid is palpable in the base
 - the cutaneous branches of the radial nerve run deep to

- e. the tendons
- e. the flexor pollicis longus tendon forms the radial side

139) With respect to the flexor retinaculum:

- a. it attaches to the scaphoid, hamate, pisiform and trapezium
- b. the tendon of palmaris longus passes deep to it
- c. the median nerve passes superficial to it
- d. the ulnar artery passes deep to it
- e. it is pierced by the tendon of flexor carpi ulnaris

140) The palmar interossei:

- a. abduct the fingers
- b. are usually supplied by a branch of the ulnar nerve
- c. have two heads
- d. arise from the medial side of the metacarpal bone of the index, ring and little fingers
- e. have no role in extension of the terminal phalanges

141) Compression of the median nerve in the carpal tunnel produces:

- a. anaesthesia over the thenar eminence
- b. wasting of the hypothenar eminence
- c. loss of power of opposition of the thumb
- d. anaesthesia of the little finger
- e. loss of power of flexion of the thumb

142) Which one of the following structures pass beneath the flexor retinaculum at the wrist?

- a. palmaris longus
- b. ulnar nerve
- c. radial artery

- d. ulnar artery
e. median nerve
- 143) The skin of the top of the index finger is supplied by:
a. radial nerve only
b. median nerve only
c. ulnar nerve only
d. radial and ulnar nerve
e. radial and median nerve
- 144) The extensor retinaculum:
a. is attached to the radius and the ulna
b. is attached to the pisiform
c. is attached to the scaphoid
d. is attached to the trapezium
- 145) The anatomical snuff box:
a. is crossed by cutaneous branches of the median nerve
b. has the anterior interosseous artery on its floor
c. contains the origin of the basilic vein
d. has extensor pollicis longus on the radial side
e. is crossed by cutaneous branches of the radial nerve
- 146) Compression of structures in the carpal tunnel usually result in:
a. weakness of the forearm flexors
b. weakness of flexion of the terminal phalanx of the thumb
c. weakness of the interossei muscles
d. normal sensation on the palmar surface of the little finger
e. weakness of the lumbrical muscles

- 147) The anatomical snuff box:
- a. has the tendon of extensor pollicis brevis on its ulnar side
 - b. has the radial artery lying in its floor
 - c. has the radial styloid process as the only palpable bony point
 - d. has the tendon of abductor pollicis brevis on its radial side
 - e. contains the origin of the basilic vein
- 148) The superficial branch of the ulnar nerve supplies:
- a. palmaris brevis
 - b. flexor digitorum ulnaris
 - c. palmar aspect index finger
 - d. opponens pollicis
 - e. thenar eminence
- 149) Damage to the median nerve may be suggested clinically by:
- a. a wasting of the interossei
 - b. pointing index finger
 - c. 'claw hand'
 - d. sensory loss over ulnar side of hand
 - e. inability to abduct the index finger
- 150) Which of the following structures IS NOT one of the contents of the carpal tunnel?
- a. flexor digitorum superficialis
 - b. ulnar artery
 - c. flexor digitorum profundus
 - d. median nerve
 - e. flexor pollicis longus

151) The only carpal bone to give attached to both flexor and extensor retinacula is:

- a. scaphoid
- b. trapezoid
- c. hamate
- d. triquetral
- e. pisiform

152) Structures in the anatomical snuff box include all the following EXCEPT:

- a. the radial tubercle
- b. the scaphoid
- c. the base of the thumb metacarpal
- d. the trapezium
- e. the radial artery

153) Which joint is the odd one out with regard to degrees of freedom?

- a. 1st carpometacarpal joint
- b. radiohumeral joint
- c. sternoclavicular joint
- d. acromioclavicular joint
- e. radiocarpal joint.

”صل على النبي“



Answers of upper limb MCQ									
1	b	31	c	61	d	91	b	121	d
2	d	32	d	62	b	92	d	122	c
3	c	33	b	63	e	93	a	123	b
4	b	34	c	64	d	94	a	124	c
5	c	35	c	65	b	95	b	125	e
6	a	36	d	66	b	96	e	126	b
7	c	37	d	67	c	97	d	127	d
8	a	38	b	68	b	98	a	128	d
9	b	39	e	69	e	99	b	129	c
10	c	40	b	70	e	100	c	130	a
11	c	41	a	71	b	101	a	131	d
12	e	42	d	72	b	102	c	132	d
13	b	43	a	73	d	103	d	133	b
14	e	44	a	74	b	104	e	134	c
15	a	45	c	75	e	105	d	135	c
16	b	46	c	76	b	106	a	136	c
17	d	47	a	77	d	107	b	137	b
18	c	48	c	78	b	108	b	138	a
19	a	49	a	79	a	109	e	139	a
20	a	50	d	80	d	110	d	140	b
21	b	51	b	81	e	111	a	141	c
22	e	52	e	82	c	112	d	142	e
23	a	53	b	83	a	113	e	143	b
24	a	54	a	84	e	114	a	144	b
25	e	55	b	85	a	115	b	145	e
26	e	56	c	86	a	116	a	146	d
27	e	57	e	87	b	117	b	147	b
28	b	58	b	88	b	118	e	148	a
29	c	59	b	89	e	119	b	149	b
30	c	60	e	90	e	120	a	150	b
151	e	152	c	153	a				