



**INSTITUTE OF HEALTH
DEPARTMENT OF BIOMEDICAL SCIENCE
ANATOMY COURSE UNIT**

CLINICAL ANATOMY OF THE UPPER LIMB.

**BY REBUMA S.
JUNE, 2022**

outline

-  Objectives
-  Introduction
-  Fracture
-  Common nerve injury
-  Occlusions of blood vessels
-  Conclusions
-  References

Objectives

At the end of this session you will be able to:

- ❖ Identify the common fracture of upper limb
- ❖ Describe common nerve injury to Upper limb
- ❖ Atrophy of related muscles and function lost
- ❖ Explain the Occlusions of Blood vessels in UL
- ❖ Identify the Clinical significance of arterial anastomosis in UL

Introduction

- ❑ **Clinical Anatomy:** Practical applications of anatomical knowledge to diagnosis and treatment of disease.
- ❑ Upper limb is a region consists of Shoulder, Axilla, Arm, Forearm & Hand with their entire neurovascular, Musculoskeletal components.

Introduction cont'd


- All structures of the upper limbs are integrated and related.
- Disease of one structure may affect the overall activity of the entire limb.
- Example: Fracture of humerus → Nerve injury → Paralysis.

Fracture

- Fracture is when the continuity of the bone is broken.
- It is common; the average person has two during a life time.
- Fracture can be simple/closed or compound/open.

Fracture of the Clavicle

- ❖ Most commonly fractured bone in the body.
- ❖ Weakest part medial and lateral third mostly involved.
- ❖ Lateral fragment is depressed by weight of the arm.
- ❖ Pulled medially and forward by the adductor muscle of the upper limb.



Fracture of the Clavicle cont'd

- ❖ Sternocleidomastoid elevates the medial fragments of the bone.
- ❖ Supraclavicular nerve involved callus formation due to close r/n ship with clavicle.
- ❖ Compression of Brachial plexus and Blood vessels.

Fractures of the Scapula

- ❖ Usually occur in a runover accident victims or occupants of automobiles involved in crashes.
- ❖ Associated with fractured ribs.
- ❖ Mostly require little treatment.

Humerus Fracture

1. Humeral Head Fracture

- ❖ Common in elder people with Osteoporosis.
- ❖ Fracture produced by fibrocartilaginous glenoid labrum of the scapula.
- ❖ Labrum become stucked and unable to move in the defect shoulder joint difficulty.

Humerus Fracture cont'd

2. Greater tuberosity fracture

- ❖ Direct trauma or displaced by the glenoid labrum in shoulder joint dislocation.
- ❖ Dislocation severing tearing of the rotator cuff.
- ❖ Open reduction of the fracture attaches the rotator cuff to its place.

Humerus Fracture cont'd


3. Lesser tuberosity Fracture

- Can be caused by Posterior dislocation of the shoulder joint.
- Fragment receive the insertion of subscapularis tendon.
- Axillary nerve may involved.

Humerus fracture cont'd

4. Surgical neck fracture


- **Directly:** blow on the lateral aspect of the shoulder.
- **Indirectly:** falling on stretched hand.
- Axillary artery and posterior humeral circumflex artery damaged
- Deltoid atrophy but not teres minor



Humerus Fracture cont'd

5. Fracture of the humerus Shaft

- Radial nerve and deep arm artery are damaged.
- Posterior brachium and antebrachium muscle are paralyzed.
- Triceps not fully paralyzed but become weak.



Humerus Fracture cont'd

6. Fracture at distal end of humerus

- ✓ Supracondylar common in children.
- ✓ Injury to radial and median nerve common but, ulnar nerve injury is not common.
- ✓ Damaged to brachial artery occur at the time of fracture.

Humerus Fracture cont'd

6. Fracture at distal end....

- It can be due to Fall on stretched hand and radial head fracture.
- Radial neck in children.
- Displacement of the fragment is usually considered.

Humerus Fracture cont'd

In general because nerves are in contact with the humerus, there may be injury when the associated part of the humerus is fractured.

- ❖ Surgical neck – axillary nerve
- ❖ Radial groove – radial nerve
- ❖ Distal humerus – median nerve
- ❖ Medial epicondyle – ulnar nerve

Fractures of the Radius and Ulna

- ❑ Fracture of the forearm may be associated with dislocation of the other bone.

Monteggia's fracture

- ❑ Shaft of the ulna fractured.
- ❑ Anterior dislocation of the radial head with rupture of anular ligament.

Fracture of the radius and ulna cont'd


Galezzi's fracture

- ❑ Proximal 3rd radius and distal ulnar dislocation at distal radio ulnar joint.

Fracture of the radius and ulna Cont'd

Colle's Fracture

- ❑ Complete fracture distal end of the radius.
- ❑ Most common of the forearm.
- ❑ Results from forced dorsiflexion of the hand.
- ❑ Ulnar styloid process avulsed



Fracture of the radius and ulna cont'd

Dinner fork deformity

- ❖ Posterior displacement of the fragment at distal end of the radius.

Smith's Fracture

- ❖ Distal end of the Radius fall on dorsum of hand.
- ❖ Distal fragment displaced anteriorly or reversed colle's fracture.


Fracture to bone of the hand

- ❖ Common in young adult.
- ❖ Unless treated properly, the fragment will not unite.
- ❖ Permanent weak and pain of wrist.
- ❖ Median nerve commonly involved.

Fracture to bone of the hand cont'd

Fracture of the Scaphoid

- Most frequently fractured carpal bone.
- Fall on the palm fracture occur at narrow part/ waist of the scaphoid.
- Radial artery and nerve involved.



Fracture to bone of the hand cont'd

Fracture of the hamate

- ☐ Occurred: non- splint of the fractured bony parts cause of the grip produced because the attached muscles.
- ☐ Ulnar nerve may be injured & results in decreased in the grip strength of the hand.
- ☐ The ulnar artery may also be damaged.

Fracture to bone of the hand cont'd

Fracture of Lunate

- ☐ Fall on stretched hand which causes the hyperextension of the wrist joint.
- ☐ Median nerve is commonly involved.

Fracture to bone of the hand cont'd

Fracture of the metacarpals bone

- ❖ May occur as a result of direct violence.
- ❖ Clenched fist striking a hard object.

Boxer's fracture

- ❖ Oblique fracture of 4th & 5th metacarpals at neck.
- ❖ The distal fragment is displaced proximally.
- ❖ Shortening of the finger posteriorly.

Fracture to bone of the hand cont'd

Bennett's Fracture

- ❖ Base of the thumb metacarpal fractured.
- ❖ Occurs when the thumb is forcefully abducted.
- ❖ Causes instability of the carpometacarpal joint of the thumb.

Brachial plexus injuries

- Affect movements/paralysis and cutaneous sensation/anesthesia of the upper limb.
- Axillary injury, Lateral cervical region wounds, stretching and disease are the leading causes.

Compression of cords of the brachial plexuses

- ❖ May result from prolonged hyperabduction of the arm.
- ❖ Compressed between the coracoid process of the scapula and the pectoralis minor tendon.
- ❖ Symptoms are radiating pain, numbness, paresthesia (tingling), erythema & weakness of the hand.


Injuries to the superior part of brachial plexuses

- C5 & C6 injuries results from excessive increase in angle between the neck & the shoulder.
- Thrown from horse and motorcycle.
- Stretching of the neck during delivery.



Erb-Duchenne paralysis or erb's paralysis

- Superior brachial plexuses injury(C6 & C7)
- The limb are medially rotated by the sternocostal part of the pectoralis major.
- Pronated forearm or action of the biceps brachi lost. **Waiter's tip position**



Muscle affected during erb palsy

Muscle Paralysis	Loss of Function	Effects
Deltoid	Loss of abduction of shoulder	Arm is adducted
Supraspinatus, Infraspinatus & Teres Minor	Loss of lateral rotation of shoulder	Arm is medially rotated
Biceps brachii & Brachialis	Elbow is extended	Loss of flexion of elbow
Biceps brachii & Supinator	Loss of supination	Forearm is pronated

Klumpke paralysis

- ❖ Inferior parts of the brachial plexuses(C8 & T1) are injured.
- ❖ Upper limbs pulled up superior. E.g. during delivery & grasping something over the head.
- ❖ Less common & muscles of the hands are involved.



Injuries of the brachial plexuses at root

Long thoracic nerve

❖ Serratus anterior

paralysed medial boarder
of the the scapula moves
laterally and posteriorly.

“Winged Scapula.”



Dropped Shoulder

- ❑ Position of the Scapula on the posterior is maintained by tone and balance of the muscle attached to it.
- ❑ Paralysis of Trapezius or Cranial nerve XI.

Injury of brachial plexuses at root

Dorsal scapular nerve injury

- ❑ Paralyzed rhomboids (minor & major) on the side of affected scapula located further from midline.

Injury to axillary nerve in the arm

- Surgical neck of the humerus fractured and deltoid atrophies.
- Unable to abduct the arm/resistance above 15°.

Injury to Radial Nerve in the Arm

- Superior to origin of its branch.
- All the muscle of the posterior compartments are paralyzed. e.g. triceps, brachioradialis, extensor. etc.
- Loss of sensation to skin supplied by the nerve.

Radial nerve injury cont'd

- ❑ At radial groove muscle of the posterior compartment of the forearm are paralyzed.
- ❑ Unable to extend elbow and wrist joint “wrist drop.”




Injury to musculocutaneous nerve

- It is beneath the biceps brachii muscle and protected so rarely injured.
- Paralysis of coracobrachialis, biceps and brachialis (weakened) little supplied by radial nerve.
- Elbow flexion, supination of forearm weakened, Loss of sensation.

Injury to median nerve at elbow

- Compression from the two head of pronator teres. (**Pronator syndrome**)
- Except, flexor carpi ulnaris and median half of the flexor digitorum profundus all flexor compartment of the forearm are paralyzed.
- Forearm kept in supine position.
- No flexion is occur at interphalangeal joint.

Injury of median nerve at elbow cont'd




Median nerve deformity

- Hand looks flattened and looks “Ape hand.”
- Opposition movement is impossible.



Olecranon bursitis


- is inflammation of the bursa overlying the olecranon process of the ulna associated with prolonged pressure at this point.



Posterior interosseous nerve entrapment

- Occurs where this branch of the radial nerve passes through two planes of fibers within the supinator muscle, often following elbow trauma or a fibrous band within the supinator.

Posterior interosseous nerve... cont'd



Ulnar nerve injury

- Posterior median epicondyle of the humerus most common.

At elbow and in forearm

- ✓ Fracturing of the medial epicondyle
- ✓ Funny or Crazy bone/ Claw hand

Ulnar nerve injury cont'd

- ✓ Difficulty in making fist
- ✓ Metacarpophalangeal joint become hyperextended.

Claw hand




Claw hand

Carpal tunnel syndrome

- ❑ Concave anterior surface of the carpal bone and closed by the flexor retinaculum.
- ❑ Compression of the median nerve '**burning pain.**'

Carpal tunnel syndrome cont'd




De Quervain's disease

- is a chronic inflammatory thickening of the common tendon sheath of the abductor pollicis longus and extensor pollicis brevis muscles.
- As they run across the lower end of the radius near the radial styloid.

De Quervain's disease cont'd


De Quervain's Tenosynovitis



Dupuytren's contracture

- is an hereditary deformity of the hand
- Due to thickening of the palmar aponeurosis with resultant fibrosis and eventual contracture of the fingers.


Dupuytren's contracture cont'd



Gamekeeper's thumb

- Is caused by a force directed on the thumb metacarpophalangeal (MCP) joint to produce a failure of the ulnar collateral ligament resulting in instability accompanied by pain and weakness of the pinch grasp.

Gamekeeper's thumb cont'd



Ulnar canal syndrome

- ❑ Compression of the ulnar nerve at wrist between pisiform and hook of hamate.
- ❑ Hypoesthesia in the medial one and half fingers.
- ❑ weakness of the intrinsic muscles of the hand.

Absent Palmaris Longus

- ✓ The palmaris longus muscle may be absent on one or both sides of the forearm in about 14% of persons.
- ✓ Others show variation such as *centrally* or *distally* placed.
- ✓ The muscle is relatively weak, its absence produces no disability.
- ✓ **Its not a deformity rather its anatomical variation.**

Absence of Pectoral Muscles

Poland syndrome: both the pectoralis major and minor are absent.

- ❖ breast hypoplasia
- ❖ absence of two to four rib segments.

Anterior interosseous nerve entrapment

- ❑ The deep branch of the median nerve, the anterior interosseous, may be trapped around the elbow following a fracture.
- ❑ The result is weakness in the **flexor pollicis longus** or **flexor profundus** muscles of the index and middle finger, making it impossible to flex the distal phalanx.


Anterior interosseous nerve.. cont'd



Mallet finger

- is avulsion of the extensor attachment of the extensor digitorum where it inserts into the dorsal terminal phalanx of the fingers.
- This is often caused by stubbing the finger against a hard object.

Mallet Finger cont'd



©MMG 2001

Summary for nerve injury

- ❖ Erb's palsy → upper trunk.
- ❖ Klumpke's palsy → lower trunk.
- ❖ Winging of scapula → long thoracic nerve.
- ❖ Ape's hand → median nerve → supracondylar fracture.
- ❖ Wrist drop → Radial nerve → fracture of spiral groove.
- ❖ Claw hand → ulnar nerve → fracture of medial epicondyle.


Occlusion of the axillary artery

- ❑ Ligation of lacerated subclavian or axillary artery necessary when stenosis occur.
- ❑ Collateral circulation is made by anastomosis.
- ❑ E.g. Suprascapular artery receives blood through anastomosis of:
 - ✓ supraclavicular, Transverse cervical, Intercostal artery.

Volkmann's contracture

- Muscular ischaemia following injury to the brachial artery typically in a supracondylar fracture of the humerus.
- Clinically, the finger flexors are usually the most severely affected, the patient being unable to extend the fingers when the wrist is flexed.

Volkman's contracture cont'd



Compression of Axillary Artery

- Can be palpated inferior part of the lateral wall of the axilla.
- Compression at 3rd part of axillary artery, humerus may be important to control profuse bleeding.

Aneurysm of Axillary Artery

- ✓ Enlargement of the first part of axillary artery.
- ✓ Compress the trunks of the brachial plexus.
- ✓ Causing pain and anesthesia.

Injuries of the axillary veins

- Wound in the axilla involves axillary vein due to large size and exposed position.
- Proximal part of axillary vein is most dangerous i.e. air enter and causes emboli.

Venipuncture and Blood transfusion

Superficial veins are clinically important for:


- ❖ Vein puncture

- ❖ Transfusion

- ❖ Cardiac catheterization

- ❑ Median cubital vein is most common site because easily visible.

Venipuncture cont'd



Conclusions

- ❑ Knowledge of the normal anatomy of the upper limb is crucial to identify the clinical conditions.
- ❑ Clinical anatomy of the upper limb includes Fracture, nerve & blood vessels injuries & Muscle paralysis.
- ❑ Fracture is the most frequently appear as clinical condition of the upper limb.

References

- 1) *Keize L., Aurther F., and Anne M., MOORE clinically oriented anatomy (2014) seventh edition.*
- 2) *Clinical anatomy by regions RICHARD S. SNELL (2012) ninth edition.*
- 3) *FRANK H. NETTER atlas of human anatomy.*
- 4) *Keith L & Anne M. Essential clinical anatomy (2007), 3rd edition.*
- 5) *Richard L, Wayne V & Adam W. Grays Anatomy for students (2007)*

THANK YOU!!!!