

Jimma University
Faculty of Health Sciences
School of Medical Laboratory Sciences

Module Name: Respiratory System

Module code: PC

Program: Regular Medicine

Academic Year: 2021/2022

Academic Level: Pre-clerkship I

Instructor: Dr. Muluaem Tadesse (PhD, Assistant professor)

Course Name: Medical Microbiology

Respiratory tract infections (RTI)

Objectives


At the end of this topic, students will be able to;

- Describe common microbes affecting respiratory tract
- Explain the clinical pictures of respiratory tract infections
- Identify diagnostic techniques to diagnose RTI
- Describe common prevention/control methods of RTI

Respiratory tract infections (RTI)


What is the respiratory system?

- The respiratory system consists of
 - the nose,
 - pharynx (FAIR inks),
 - larynx (LAIR inks),
 - trachea (TRAY kee ah),
 - bronchi (BRAHN kye), and
 - Lungs



The primary function of this system is to furnish oxygen for individual tissue cells, and to take away the waste products and carbon dioxide produced by those same cells

Parts of RTI



The Respiratory Tract and Its Defenses

- Most common place for infectious agents to gain access to the body
- Upper respiratory tract:
 - mouth, nose, nasal cavity, sinuses, pharynx, epiglottis, larynx
- Lower respiratory tract:
 - trachea, bronchi, bronchioles, lungs, alveoli
- Defenses
 - Nasal hair
 - Cilia
 - Mucus
 - Involuntary responses such as coughing, sneezing, and swallowing
 - Macrophages
 - Secretory IgA against specific pathogens

Normal Microflora of the Respiratory Tract

- Generally limited to the upper respiratory tract
- Gram-positive bacteria (streptococci and staphylococci) very common
- Disease-causing bacteria are present as normal biota;
 - can cause disease if their host becomes immunocompromised or
 - if they are transferred to other hosts
- Normal biota perform microbial antagonism

Copyright © The McGraw-Hill Companies, Inc. Permission required for reproduction or display.


Respiratory Tract Defenses and Normal Biota

	Defenses	Normal Biota
Upper Respiratory Tract	Nasal hair, ciliary escalator, mucus, involuntary responses such as coughing and sneezing, secretory IgA	<i>Moraxella</i> , nonhemolytic and alpha-hemolytic streptococci, <i>Corynebacterium</i> and other diphtheroids, <i>Candida albicans</i> Note: <i>Streptococcus pyogenes</i> , <i>Streptococcus pneumoniae</i> , <i>Haemophilus influenzae</i> , <i>Neisseria meningitidis</i> , and <i>Staphylococcus aureus</i> often present as “normal” biota.
Lower Respiratory Tract	Mucus, alveolar macrophages, secretory IgA	None

Respiratory tract infections (RTI)

- Upper respiratory tract infection (URTI)
 - Viral, bacterial
- Lower respiratory tract infection (LRTI)
 - Bacterial, viral, fungal, parasite

Diseases of the respiratory system



Upper Respiratory Tract Infections

- Most caused by viruses & usually nothing more than irritation
- Direct invasion of respiratory epithelium results in symptoms corresponding to the area(s) involved
- May spread down and cause more severe infections

Infections of **airway above glottis** **or vocal cords**

- Tonsillitis
- Pharyngitis
- Laryngitis
- Sinusitis
- Otitis media
- Rhinitis

- 1) Cough
- 2) Sore throat
- 3) Running nose
- 4) Nasal congestion
- 5) Headache
- 6) Low-grade fever
- 7) Facial pressure
- 8) Sneezing

Bacterial agents of URTI...

1. *Corynebacterium diphtheriae*

General Characteristics

- Gram-positive bacilli,
- Non Endospore-Forming Bacilli
- Pleomorphic (Curved or straight).
- Non-motile, non-acid fast and aerobic.
- Club-shaped forms = Swelled end
 - Chinese characters
- Found as free-living saprophytes
- Members of the usual flora of humans and animals
 - colonize skin, URT, urogenital tract of healthy people

Virulence Factors *C. diphtheriae*

- Exotoxin as the virulence factor
- Diphtheria toxin is secreted at the site of infection

A: Active fragment

Inhibits protein synthesis

Leads to cell/tissue death


B binds to specific cell membrane receptors

Mediates entry of fragment A

PATHOGENESIS

Entry into nose or mouth

The organism remains in the superficial layers of skin lesions or respiratory tract mucosa, inducing local inflammatory reaction



The major virulence of the organism lies in its ability to produce the potent 62-kd polypeptide exotoxin, which inhibits protein synthesis and causes local tissue necrosis



Within the first few days of respiratory tract infection , a dense necrotic coagulum of organisms, epithelial cells, fibrin, leukocytes and erythrocytes forms, advances, and becomes a gray-brown, leather-like adherent pseudomembrane . Removal is difficult and reveals a bleeding edematous submucosa

Clinical presentations

- ▶ Asymptomatic
- ▶ Respiratory diphtheria
 - Anterior nasal diphtheria
 - Pharyngeal and tonsillar diphtheria
 - Laryngeal diphtheria
- ▶ Cutaneous diphtheria

a. Diphtheriae

Local lesion in URT →

necrotic injury to epithelial cells

→ blood plasma leaks

fibrin network interlacing with
growing bacteria

pseudomembrane

Toxin causes inhibition of protein synthesis
in cells death of eukaryotic cells and
tissues

Diphtheria - pseudomembrane

- Thick grey 'pseudomembrane' composed of fibrin, epithelial cells, bacteria and polymorph neutrophils
- Pseudomembrane may cause blockage, suffocation




Figure 25.20b Microbiology: An Evolving Science
©Princess Margaret Hospital, Hong Kong

Tonsillar and pharyngeal diphtheria: sore throat is the universal early symptom

- Mild pharyngeal injection → unilateral or bilateral tonsillar membrane formation → extend to involve the uvula, soft palate, posterior oropharynx, hypopharynx, or glottic areas
- Underlying soft tissue edema and enlarged lymph nodes: **bull-neck appearance**
 - Classic *cutaneous diphtheria* is an indolent, nonprogressive infection characterized by a superficial, ecthymic, nonhealing ulcer with a gray-brown membrane



URTI....cont'd

b. Tonsilar diphtheria



Lab diagnosis

I. Diagnosis of case: aims at confirming clinical diagnosis

- Throat swabs from the membranes are examined as:
 - a. **Direct smears:** stained with gram and Methylene blue G+ve bacilli with characteristic morphology may be seen
 - b. **Cultures on** loeffler's serum and blood tellurite media. Colonies are picked & stained with gram & Methylene blue
 - c. **Elek test:** rapid diagnosis (16-24 hrs)
 - Results must be confirmed by toxigenicity tests (in vivo & vitro virulence test and tissue culture tests)
 - d. **PCR** - to detect toxin gene (*tox gene*)
 - e. **ELISA** - to detect toxin from clinical isolates

2. Diagnosis of susceptibility by the Schick test:

- Done to determine if a person is susceptible or immune to diphtheria by detecting presence or absence of antitoxin Abs
- ID injection of a small amount of toxin in one forearm & an equal amount of heat inactivated toxin in the other forearm (control)
- If both sites give no reaction => **Schick negative** (person immune)
- If a necrotic area develops within 3 days at the site of injection in the test arm only => **Schick positive** => person susceptible
- If reaction develops in both arms; allergy to proteins of the toxin

Prevention, control and treatment of diphtheria

1. Prophylaxis: Active immunization

- a. Fluid toxoid:** A filtrate of broth culture (toxigenic strain) treated with 0.3% formalin (remove toxicity, immunogenicity)
- b. Aluminium precipitated toxoid:** Fluid toxoid adsorbed onto aluminium hydroxide or aluminium phosphate
- Toxoids are commonly combined with tetanus toxoid & pertussis vaccine (DPT) and given IM to children at the age of 2,4 & 6 month

2. Diphtheria antitoxin serum given without delay

- Antitoxin neutralizes toxin before causes irreversible damage
- Dose: 20,000 - 100,000U (IM or IV); effective with antibiotics

3. Antibiotic

- Penicillin & erythromycin (when given with antiserum; reduce number of diphtheria in throat & decrease incidence of carrier

2. *Streptococcus pyogenes*

- Gram positive cocci in chain
- Group A streptococcus (GAS) with 40 antigenic types
- *S. pyogenes* is β hemolytic
- 90% of cases of pharyngitis
- Flesh eating bacteria
- Most important human pathogen causing disease including:
 - Suppurative conditions/skin infections
 - Throat infections
 - Systemic infections
 - Non-suppurative sequelae
- About 5-15% of normal individuals harbor in their respiratory tract, without signs of disease

Determinants of Pathogenicity--GAB

1. **avoid phagocytosis** - mediated primarily by capsule
2. Adhere to and invade host cells- M protein, lipoteichoic acid,
3. produce toxins and Enzymes:-
4. streptococcal pyrogenic exotoxins,
 - The toxins act as **superantigens**-- with the enhanced release of proinflammatory cytokines.
5. streptolysin S -- an oxygen-stable, nonimmunogenic,
6. streptolysin O--**It is immunogenic** (antistreptolysin O [ASO])
7. streptokinase, **lyse blood clots and fibrin deposits** and facilitate the rapid spread of **S. pyogenes in infected tissues**
8. DNases

Suppurative Streptococcal Disease

1. **Sore throat** and follicular tonsillitis (**pharyngitis**)

- Most common infection; enlarged tonsils with purulent exudates, high fever and enlarged cervical lymph nodes

2. **Scarlet fever**: Disease of children characterized by sore throat and erythematous rash; erythrogenic toxin

3. Pyoderma (impetigo) is a confined, purulent (“pyo”) infection of the skin (“derma”)

4. Erysipelas (erythros, “red”; pella, “skin”) is an acute infection of the skin.

5. Pyoderma (impetigo) is a confined, purulent (“pyo”) infection of the skin (“derma”) that primarily affects exposed areas (i.e., face, arms, legs).

Scarlet fever

Strawberry tongue:
papilla hypertrophy on
the tongue




Fig. 8 Scarlet fever: initially the tongue is covered by a white exudate through which the papillae project (white strawberry tongue - above). Later the exudate is shed to reveal the bright red inflammation of the underlying tissue (red strawberry tongue - left).

Exanthems
on the skin



Flesh Eating Bacteria

- Cellulitis- leading to Necrotising Fasciitis
- Leads to necrosis of subcutaneous tissues and muscular tissues and adjacent fascia
- Other complications
Toxic shock syndrome





Non Suppurative complications or sequelae

- Post-infection complications of Group A streptococcal disease; Serious complications in pre-antibiotic era
- **Acute rheumatic fever (ARF):**
 - following pharyngitis
 - Morbidity & mortality linked to subsequent disease of heart valve (Rheumatic Heart Disease)
 - Poorly understood pathogenesis with several proposed theories including **cross-reactivity of heart tissues & strep AGNs**
 - M protein cross-reacts heart myosin
 - Autoimmunity and potentially fatal
- **Acute Glomerulonephritis**- following skin infection
 - Immune complex mediated disease
 - inflammation of glomeruli due to Ag-Ab complex deposit on basement membrane
 - Clinically- Hematuria, Proteinuria, Hypertension

Lab diagnosis of streptococcal diseases:

- **Specimens:** swabs from throat or other lesions, pus, or blood (bacteraemia)
- **Direct gram smears:** Gram-positive cocci in chains
- **Culture on BA:** colonies producing complete haemolysis



Bacitracin test:

- Differentiated GAS from other β -hemolytic streptococci (B, C, G) by their sensitivity to bacitracin
- Done by placing filter paper disc containing bacitracin on blood agar inoculated with organism (zone of inhibition)

Antigen detection tests:

- Rapid detection of GAS Ags from throat swab can be done by ELISA or agglutination tests

Serologic tests: to diagnose post-streptococcal d/ses

- ASO test

Other URTI causing bacteria

- *Haemophilus influenza*
- *Moraxella catarrhalis*
- *Streptococcus pneumoniae*
- *Staphylococcus aureus*


Viral causative agents of URTI

1. Corona viruses

- *Corona* = crown like structure, pleomorphic, spiked
- Non-segmented, linear, ssRNA
- Enveloped +ve stranded RNA
- mRNA encased in nucleocapsid
- Lipid Bilayer – Soap works to disrupt this!
- *Causes diseases in Man, animals and birds*

three glycoproteins:

- ✓ **S - Spike protein:** receptor binding, cell fusion, major antigen
- ✓ **M - Membrane protein:** transmembrane - budding & envelope formation
- ✓ **HE – Hemagglutinin:** hemagglutination, acetylsterase activity



Corona viruses– pathogenesis

- Normal human coronaviruses cause 5-10% of common cold/URIs, with outbreaks to 30% of common cold
- Display a tropism for **epithelial cells** of the **respiratory or gastrointestinal tract**
- Remain limited to the upper respiratory tract
- a slow , patchy destruction of ciliated epithelial cells and the loss of beating cilia → disease development
- **High frequency of recombination during replication - evolution of new strains**

2. Rhinoviruses

- Common cold viruses
- Mild URT illnesses
- Nasal secretions, throat and oral secretions

Properties

- inactivated by low pH (pH 3)
- More thermo stable than enteroviruses and survive for days on environmental surfaces
- Infect only **humans, chimpanzee**
- Replication is limited to the **epithelium of the nasal mucosa**
- Human cell lines - **grow better at 33°C** (the temperature of the nasopharynx in human)
- **Common cold viruses with >100 antigenic types**

Rhinoviruses

Clinical findings


- Common cold
- IP → 2-4 days
- Acute illness usually lasts for 7 days
- Nonproductive cough may persist for 2-3 weeks
- Sneezing, nasal obstruction, nasal discharge & sore throat
- headache, malaise, mild cough, & a chilly sensation
- Little or no fever
- The nasal and nasopharyngeal mucosa become red and swollen and the sense of smell is decreased
- **No specific treatment/vaccine**

3. Adenoviruses

- Family: **Adenoviridae**
- Non-enveloped, dsDNA virus
- Linear ds DNA Genome
- 51- human adenoviruses, grouped in 6-species (A-F)
- Encode proteins that block MHC class I expression
- Cough, nasal congestion, fever, sore throat and rarely pneumonia
- Rounding, enlargement and aggregation of affected cells into grape-like clusters in cell cultures

Pathogenesis- Adenovirus

- Infect and replicate in **epithelial cells** of respiratory tract, eye, gastrointestinal tract, urinary bladder, and liver.
- Usually do not spread beyond the **regional lymph nodes**
- Group **C viruses** persist as latent infections for years in **adenoids and tonsils** and shed in the **feces** for many months after the initial infection



URTI....cont'd

Other URTI causing viruses

- Influenza viruses (type A and B)
- Parainfluenza virus
- Coxsackie A virus
- Epstein Bar Virus
- Herpes Simplex Virus
- Respiratory Syncytial Virus
- Rubella virus
- Varicella -zoster virus
- Cytomegalo virus
- Human metapneumovirus

Summary

- Non-specific, often combinations of viral & bacterial infections
- Diagnosis is mainly based on clinical manifestations
- Treatment: to maximize relief of most prominent symptoms
- Most are self-limiting but
 - Doctor, family & patient factors - over prescribe antibiotic
 - Treat only proven group A strep with antibiotic
- Common cold, sinusitis, pharyngitis, epiglottitis & laryngitis

Lower respiratory tract infections (LRTI)

Lower respiratory tract infections (LRTI)

- Lower respiratory tract
 - Trachea, bronchi, bronchioles, and alveoli in the lungs
 - Most are bacterial causes
 - Far more serious than upper respiratory tract infections
 - Many of the microbes infecting upper respiratory system can also infect the lower respiratory system
 - Bacterial, viral, fungal and parasitic infections can cause Inflammation of the LRT

Bacterial infections of lower respiratory tract

- Bacterial pneumonia
- Pertussis
- Inhalation anthrax
- Q fever
- Tuberculosis

1. Bacterial pneumonia

- Inflammatory condition of the lung in which fluid fills the alveoli
- Can be caused by a wide variety of different microorganisms
- Viral pneumonias are usually milder than bacterial
 - One of the most serious lower respiratory tract infections
 - Bacterial pneumonia can be divided into two types:
 - Nosocomial/hospital acquired
 - Community-acquired
- Each type can be caused by a variety of organisms

Bacterial pneumonia

* Classification of pneumonia:

1. Community-Acquired Acute Pneumonia:

- *Streptococcus pneumonia*
- *Haemophilus influenza pneumonia*
- *Staphylococcus aureus pneumonia*
- *Legionella pneumophila pneumonia*

2. Community-Acquired Atypical Pneumonia:

- *Mycoplasma pneumonia*
- Viral *pneumonia*: Respiratory syncytial virus, para-influenza virus (children); influenza A and B (adults); adenovirus.

3. Nosocomial Pneumonia.

4. Aspiration Pneumonia.

5. Chronic Pneumonia.

- Granulomatous: *Mycobacterium tuberculosis* and atypical mycobacteria.
- *Nocardia*.
- *Actinomyces*.
- *Histoplasma capsulatum*.

....Bacterial pneumonia

■ Nosocomial pneumonia

- Occurs ~ 48 hours after admission to hospital
- Usually associated with *Staphylococcus aureus*
- Also caused by gram-negative bacteria
- Particularly difficult to deal with antibiotic resistant s

■ Community-acquired pneumonia

- Usually presents as a lobar pneumonia
- Accompanied by fever, chest pain, and production of purulent sputum

Community-acquired pneumonia:

- Require enough pathogens to overwhelm resident defenses
- Establishment of an infection in the lungs depends on:
 - Number of pathogen entering and competence of immunity
 - *Streptococcus pneumoniae*
 - *Haemophilus influenzae*
 - *Staphylococcus aureus*
 - *Legionella pneumophila*
 - *Mycoplasma pneumoniae*
 - *Chlamydia pneumoniae*,
 - *chlamydia psittaci*

Pneumonia...cont'd

1. *Streptococcus pneumonia*

- Capsulated, G+ve, lancet-shaped cocci (usually in pairs)
- They do not display an M protein
- Common cause of pneumonia (usually of lobar type)
- Pneumococcus (morphology, consistent) - pneumococcal
- Can also cause sinusitis, otitis media, meningitis
- Leading cause of invasive d/se in children and the elderly

Pneumococcal Pneumonia: Pathogenesis

- Nasopharyngeal colonization: ~40% of population
- Reaching to lower respiratory tract by aerosol
- Progress to alveolus and associate with specific alveolar cells (produce choline-containing surfactant)
- Activation of inflammatory cascades
- Altered vascular permeability, arrival of inflammatory exudates, arrival of leukocytes (switch exudates from serous to purulent)
- Pneumolysin and hydrogen peroxide kill cells

Clinical diseases

- Colonizes oropharynx then can spread to lung sinus, middle ear, can be transported to blood and then different organs (meningitis, endocarditis, septic arthritis)
- Causes 60-70% of all ***bacterial pneumonias***
 - One of the most common causes of **community acquired disease**
 - after damage to upper respiratory tract --following viral infection
 - multiply & induce an overwhelming inflammatory response
 - Acute chills, fever, productive cough with blood mixed sputum
- Meningitis – Most common causes of bacterial meningitis
- Common cause of sinusitis, **otitis media**
- Can cause a variety of systemic infections like bacteremia and endocarditis

Community-acquired *Bacterial* pneumonia

TWO types:

A . Atypical pneumonia

- Coughing without sputum
- Caused by a variety of bacteria
- Bacterial pneumonia can progress to the production of lung abscesses
- Caused by viruses, Mycoplasma, Chlamydia

Atypical Pneumonia

- Fever and malaise precede respiratory symptoms few days
- Severe headache, malaise, anorexia
- No localized signs on chest exam,
- No consolidation on chest x-ray
- Spleen may be enlarged
- WBC normal, cultures negative
- No improvement with Penicillin

Atypical Pneumonia

- Atypical Pneumonia (community-acquired)
 - Mycoplasma
 - Sporadic or epidemics
 - Viruses
 - Influenza, Parainfluenza, Adenovirus, respiratory syncytial virus, measles, chicken pox
 - Chlamydia

Atypical pneumonia

- Morphology:
 - Patchy or involve whole lobe
 - Inflammation is confined to the alveolar walls
 - Widening of alveolar walls by edema, mononuclear cell infiltration (lymphocytes, plasma cells, macrophages)

B. Typical *pneumonia*

Lobar Pneumonia:

- *S. pneumoniae* that affects part of a lobe in the lung or it may affect more than one lobe

Bronchial Pneumonia:

- Pneumonia spreads to several patches in one or both lungs
- Most prevalent in infants, young children and aged adults
- Cough (with or without mucus), chest pain, rapid breathing, and shortness of breath
- Transmitted by respiratory droplets

Treatment...Bacterial pneumonia

- Course of treatment depends on:
 - Severity of the infection
 - Type of organism causing the infection
- Most common pathogen: *Streptococcus pneumoniae*
 - Treated with penicillin, amoxicillin-clavulanate, and erythromycin

Hospital acquired (Nosocomial) pneumonia

- Nosocomial, or hospital-acquired pneumonias are defined as pulmonary infections acquired in the course of a hospital stay.
- Nosocomial infections are common in hospitalized persons with severe underlying disease, those who are immunosuppressed, or those on prolonged antibiotic regimens.
- Those on mechanical ventilation represent a particularly high-risk group, and infections acquired in this setting are given the distinctive designation *ventilator-associated pneumonia*.

* Causative agents:

1. Gram-negative rods (Enterobacteriaceae and *Pseudomonas* spp.)
2. *Staph. aureus*.

Staphylococcus aureus pneumonia

- *S. aureus* is an important cause of secondary bacterial pneumonia in children and healthy adults after viral respiratory illnesses (e.g., measles in children and influenza in both children and adults).
- Staphylococcal pneumonia is associated with a high incidence of complications, such as lung abscess and empyema.
- Staphylococcal pneumonia occurring in association with right-sided staphylococcal endocarditis is a serious complication of *intravenous drug abuse*.

Klebsiella pneumoniae

- *K. pneumoniae* is the most frequent cause of gram negative bacterial pneumonia.
- *Klebsiella*-related pneumonia frequently afflicts debilitated and malnourished persons, particularly *chronic alcoholics*.
- Thick and gelatinous sputum is characteristic, because the organism produces an abundant viscid capsular polysaccharide, which the patient may have difficulty coughing up.

Pseudomonas aeruginosa

- *Pseudomonas* pneumonia is common in persons who are neutropenic, usually secondary to chemotherapy; in victims of extensive burns; and in patients requiring mechanical ventilation.
- *P. aeruginosa* has a propensity to invade blood vessels at the site of infection, with consequent extra-pulmonary spread; *Pseudomonas* bacteremia is a fulminant disease, with death often occurring within a matter of days.

2. Tuberculosis

- Caused by *Mycobacterium tuberculosis*
 - Rod-shaped, Acid-fast bacillus, non-spore forming
 - Produces mycolic acid
 - Makes it difficult to Gram stain
 - Protects pathogen from antibiotic therapy and host defenses




Figure 21.12 Microbiology: A Clinical Approach (© Garland Science)

Pathogenesis...Tuberculosis

- *MTB* cell wall interferes with macrophage function and immune cell activation
 - Inhibits the formation of the phagolysosome
- This allows *MTB* to escape into the cytoplasm where it:
 - Increases in number & eventually spreads to lymph nodes, enters blood and distributed throughout body
- Fever, Fatigue, Weight loss, Chest pain, Shortness of breath, Congestion with coughing

Pathogenesis...Tuberculosis

- Two basic types of tuberculosis

- Primary

- Follows initial exposure to the pathogen

- Secondary

- Can occur years later




Figure 21.13 Microbiology: A Clinical Approach (© Garland Science)

Diagnosis of TB

- Depends on detection and identification of tubercle bacilli (TB) and their isolation from pathologic specimens
- TB can affect every tissue in the body, specimen differs
- Sputum (Pulmonary), Urine (renal), CSF (meningeal), Stools (intestinal)

1. Direct smears:

- Specimens: stained with Ziehl-Neelsen stain.
- Detection of AFB in sputum gives a fairly strong indication of PTB, but are detected only if they are present in large numbers

Diagnosiscont'd

2. Concentration (Petroff's method):

- Negative direct smear and contaminated specimens are subjected to concentration before processing
- Specimen is mixed with equal volume of 4% NaOH
- Mixture is incubated at 37°C for 30 min, shake every 5 min
- Centrifugation and deposit is neutralized by 8% HCl

What are the importances of concentration?

Diagnosiscont'd

3.Tuberculin Test (Mantoux test)

- Delayed hypersensitivity skin test; used to detect cell mediated immunity to TB
- Intradermal (ID) injection purified protein derivative (PPD) of the tubercle bacilli containing 5 tuberculin units (TU).
- In positive tests, local area of induration 10 mm in diameter or more develops 48-72 hrs after injection.
- Positive test in adults indicates that individual has been previously exposed to tubercle bacilli and continues to carry viable bacilli.

Treatment

1. **Prolonged treatment:** slow response of TB treatment and should be continued for 6-12 mon (DOTs=6-8 mon): **Why?**
 - a. Most bacilli are found intracellularly
 - b. Caseous material interfere with the drug
 - c. In chronic TB, bacilli are not dividing, i.e. "metabolically inactive", hence resistant to drugs.
2. **Combination of drugs:** to reduce toxicity due to prolonged course of treatment and rapid emergence of resistant strains.
 - ✓ Drug of choice: isoniazide (INH), rifampicin, ethambutol, streptomycin & para-amino-salicylic acid

Prevention

A. Public health measures

- Early diagnosis and treatment until they become non-infectious
- Control of transmission

B. Vaccination:

- A live-attenuated vaccine – BCG commonly used
- Prepared from bovine strain with a fixed low virulence
- Vaccine is given in a single dose of 0.1ml ID in the deltoid region to stimulate hypersensitivity and CMI against infection

3. Pertussis (whooping cough)

- *B. pertussis*: etiologic agent of **whooping cough**
- *B. parapertussis* & *B. bronchiseptica*: mild form of whooping cough


Bordetella pertussis

- Short, gram-negative bacilli
- **Pili & haemagglutinin**: adheres to ciliated epithelia of RT
- **Pertussis toxin** - promotes phagocytosis
- **Tracheal cytotoxin** inhibit DNA synthesis in ciliated cells
- **LPS**: causes damage to epithelial cells of the RT

B. pertussis

- Adheres and multiplies rapidly on epithelial surface of trachea & bronchi and interfere with ciliary action.
- Liberate toxins & substances that irritate surface cells, causing cough & marked lymphocytosis.
- Cough is intermittent with bursting attacks accompanied by whoop followed by vomiting
- Complications: bronchopneumonia, subconjunctival or cerebral hemorrhage due to paroxysms of severe cough.

Colonization of tracheal epithelial cells by *B.pertussis*



Whooping cough

Diagnosis:

- Isolation of bacteria: positive during the 1st week only
- Culture from mucus or droplets: **Mercury drops colony**
- Serology: using specific anti-sera or specific antigens

Prophylaxis:

- Heat killed vaccine to children during 1st year of life in combination with diphtheria and tetanus (DPT)

Treatment:

- Antibiotics: erythromycin effective if given early
- Recovery is followed by long lasting immunity

4. Inhalation anthrax

- Produces a fulminate pneumonia
 - Comes suddenly with great severity
 - Leads to respiratory failure and death
- Anthrax primarily a disease of herbivores
 - Acquired from spores found in pastures
 - If spores inhaled, anthrax can occur in respiratory tract
- Infection is infrequently seen in healthy individuals
 - Usually presents as localized lesions where it occurs
- Recent interest: as a biological weapon

Pathogenesis...Inhalation anthrax

- The causative agent is *Bacillus anthracis*.
 - Gram-positive rod
 - Spore-forming
- Spores germinate in human tissues
- Antiphagocytic properties of the capsule aid its survival and growth in large numbers

Pathogenesis...Inhalation anthrax

- Pathogenesis results from the powerful **exotoxin** produced
- Symptoms include:
 - Nonspecific malaise, mild fever, nonproductive cough
 - Progressive respiratory distress and cyanosis
- Rapid and massive spread to blood and CNS - death
- Antibiotic therapy can be successful
 - *B. anthracis* is susceptible to penicillin
 - Doxycycline and ciprofloxacin are alternative prophylactics

5. *Legionella* pneumonia (legionnaires' disease)

- Caused by *Legionella pneumophila*
 - Gram-negative rod
 - Cannot be stained or grown using normal techniques
- Transmitted to humans as a humidified aerosol; not person to person
- *Legionella* is ubiquitous in fresh water
- Erythromycin is better than penicillin





Figure 21.15 Microbiology: A Clinical Approach (© Garland Science)

Spread of Legionnaires' Disease

Figure 1. Conditions for Contracting Legionnaires' Disease



6. Q fever

- A zoonotic infection seen world wide
- Caused by *Coxiella burnetii*
 - Obligate intracellular pathogen
 - Stable and resistant
 - Killed by pasteurization
- Grows well in placenta of animals and large numbers can be transmitted by inhalation during animal births
- Transmission can also be by ingestion of unpasteurized milk
- Mild hacking cough and patchy interstitial pneumonia
- Most cases resolve spontaneously


6. Q fever

- A zoonotic infection seen world wide
- Caused by *Coxiella burnetii*
 - Obligate intracellular pathogen
 - Stable and resistant
 - Killed by pasteurization
- **Transmission**
 - Grows well in **placenta of animals** and large numbers can be transmitted by inhalation during animal births
 - Aerosol
 - Urine, feces, milk
 - Direct contact
 - Fomites
 - Ingestion
 - Arthropods (ticks)




6. Q fever

- A zoonotic infection seen world wide
- Caused by *Coxiella burnetii*
 - Obligate intracellular pathogen
 - Stable and resistant
 - Killed by pasteurization
- Grows well in placenta of animals and large numbers can be transmitted by inhalation during animal births
- Transmission can also be by ingestion of unpasteurized milk
- Mild hacking cough and patchy interstitial pneumonia
- Most cases resolve spontaneously
- *Tetracycline* can be given to shorten fever



Inhalation of contaminated aerosols



-  Small cell variants
-  Spores

Acute Q fever

60% cases → Asymptomatic

40% cases → Symptomatic

(95-98) % cases → Mild
(Undiagnosed)

(2-5) % cases → Severe
(Hospitalized)

- Severe fever
- Pneumonia
- Hepatitis

On long run 1-5% acute
Q fever cases develops
into chronic Q fever

Predisposing factors

- Immunosuppression
- Pregnancy
- Heart valve lesions
- Vascular abnormalities

Chronic Q fever

Valve lesion (2% cases)

- Endocarditis (30-60% within two years)

- Chronic fatigue syndrome
- Osteomyelitis
- Chronic hepatitis
- Interstitial lung disease
- Repeated abortions

Viral infections of the lower respiratory tract

Viral infections of lower respiratory tract

- Majority of infections in the lower respiratory tract are caused by:
 - Influenza virus
 - Respiratory syncytial virus
- Common characteristics of infection are:
 - Short incubation period of 1 to 4 days
 - Transmission from person to person
- Transmission can be direct or indirect
 - Direct – through droplets
 - Indirect – through hand transfer of contaminated secretions

Why has viral pneumonias become important?

- Increase in immunocompromised patients and population at-risk groups
- Discovery of new respiratory viruses
 - Human metapneumovirus
 - Corona viruses - NL63 and HKU1
 - Hantavirus
 - Human bocavirus
- Emergence of new viruses
 - Severe acute respiratory syndrome (SARS)
 - Avian influenza A (H5N1) virus,
 - 2009 pandemic influenza A (H1N1) virus

1. Severe Acute Respiratory Syndrome (SARS)

- New emerged respiratory disease (2002)
- Symptoms are initially usually mild but latter patients may develop dry non-productive cough and breathing may become difficult (dyspnea)
- Shortness of breath → hypoxia
- Respiratory distress leads to death in 3-30 % of cases
- Laboratory tests show a reduction in lymphocyte and rise in aminotransferase activity - damage to liver

SARS... cont'd

SARS Laboratory findings: CDC recommendations

- Chest radiograph, pulse oximetry, tissue culture
- Sputum gram stain/culture (for Legionella and pneumococcal)
- Testing for viral respiratory pathogens (influenza A, B, RSV)
- Nucleic acid assays to detect SARS-CoV
- ELISA to detect SARS-CoV antibodies
- RT-PCR to detect SARS-CoV infection

2. Respiratory syncytial virus

- Common cause of fatal ARTI in infants/young children
- Highly contagious; period precedes symptoms
- Localized infections of LRT, no viremia
- In children < 1 yo; bronchiolitis - wheezing, dyspnea, decreased ventilation, hyperexpansion of lung, air trapping- resembles asthma
- Cause pneumonia & common cold in children/adults

3. Influenza viruses

- Three types (A, B, C); matrix & nucleoprotein antigens
- Febrile respiratory d/se with systemic symptoms (**flu**); fever, chills, cough, sore throat, runny nose, muscle or body aches, headache, fatigue (tiredness) & vomiting
- Type A infects humans, swine, horses, whales, birds
 - ✓ Primary reservoirs (birds); infection is mostly asymptomatic, virus replicate in lungs and intestinal mucosa; shed in feces => human respiratory infection
- Influenza B & C are human viruses; do not infect birds

Influenza... cont'd

- Orthomyxovirus, enveloped
- Segmented ssRNA virus; allows a high rate of mutation
 - Hemagglutinin (HA)
 - Neuraminidase (NA)
 - M proteins (matrix protein-M1, membrane protein-M2)
 - Nucleoprotein (NP)
 - Polymerase proteins (PB2, PB1, PA)
 - Non structural proteins (NS1, NS2)

Influenza...cont'd





Figure 21.17 Microbiology: A Clinical Approach (© Garland Science)

(b)



Panel B: © Dennis Kunkel

Influenza... cont'd

Hemagglutinin

- Required for virus binding to cell surface glycolipids and glycoproteins, 15 HA subtypes
- Responsible for penetration; Abs to HA neutralize virus

Neuraminidase

- Removes sialic acid from glycoconjugate; aiding spread and prevents virus clustering at cell surface upon release
- High concentration of anti-NA antibody are necessary for virus neutralization; 9 NA subtypes

Pathogenesis...Influenza

- Influenza virus prefers the respiratory epithelium
 - Viremia is rare
- Virus multiplies in the ciliated cells of lower respiratory tract
 - Results in functional and structural abnormalities
- Cellular synthesis of nucleic acids and proteins is shut down
- Ciliated and mucus-producing epithelial cells are shed.
 - Substantial interference with clearance mechanisms
 - Localized inflammation; both in URT and LRT

Pathogenesis: *Influenza... cont'd*

- Local symptoms result from epithelial cell damage, including ciliated and mucus-secreting cells
- Systemic symptoms: interferon and lymphokine response
- Infected people predisposed to bacterial super infection
- HA & NA of Influenza A: undergo **major (reassortment: shift)** and **minor (mutation: drift)** antigenic changes to ensure presence of immunologically naïve, susceptible people

Antigenic shift: *Influenza... cont'd*

- Reassortment (recombination); individual RNA segments segregate independently in mixed infections => Antigenic shift; abrupt, major change in virus => produce new combinations of HA and NA proteins
- Viruses may reassort in non-human species, shielded from human immunity and may involve interspecies transmission
- Reassorted viruses express new HA, population has no immunity for the newly expressed antigen
- Shift accounts for major pandemics

Antigenic drift: *Influenza... cont'd*

- Gradual, continuous change occurred when virus makes small 'mistakes' during replication (high mutation rate)
- Antigen variation cannot explained by high mutation rate alone
- Point mutations occur in HA (and NA) in antigenic sites => slight change in HA or NA proteins => immune evasion
- New viruses escape; got selective advantage
- Drift accounts for frequent epidemics

Influenza... cont'd

	Type A	Type B	Type C
Severity of illness	++++	++	+
Animal reservoir	yes	no	no
Human pandemics	yes	no	no
Human epidemics	yes	yes	no (sporadic)
Antigenic changes	shift, drift	drift	drift
Amantadine/rimantidine	sensitive	no effect	no effect
Zanamivir	sensitive	sensitive	--
Surface glycoproteins	2	2	1

Lab diagnosis: *Influenza... cont'd*

Diagnosis:

- Cell or tissue culture
- Hemadsorption
- Viral antigen detection

Treatment: *Influenza...cont'd*

- Two basic approaches
 - Symptomatic care & anticipation of potential complications
- Best treatments include:
 - Rest and fluid intake, cough suppressants
 - Conservative use of analgesics for myalgia and headache
- Amantidine & rimantadine (diagnosed within 12-24 hrs) (M2)
- Zanamivir and tamiflu (oseltamivir) target neuraminidase

Prevention: *influenza...cont'd*

Vaccination

- Trivalent: two current A strains and one current B strain
- Formalin fixed wild type virus approved for parenterally administered vaccination.
- Live attenuated vaccine: temperature sensitive recombinant bearing relevant HA and NA genes
- Must anticipate shift and drift in order to identify appropriate vaccine strain

Fungal infections of the Lower respiratory tract

Fungal infections of lower respiratory tract

- Fungal spores are easily inhaled and may germinate in LRT
- Two major factors govern incidence and spread of fungal infection
 - Ubiquity of the infectious organisms
 - Found in soil
 - Resident flora
 - Adaptive immune response
 - Usually keeps these infections under control
 - Immunocompromised patients at much greater risk
- Incidence is being increased in recent years. Why?

1. *Pneumocystis* pneumonia (pcp)

- A lethal pneumonia

- Common in AIDS patients



- Caused by the fungus *Pneumocystis (carinii) jiroveci*

- Never been grown in culture

- Most information comes from clinical information of patients


Pneumocystis Pneumonia

- *Pneumocystis jirovecii* (formerly *P.carinii*) is sometimes found in healthy human lung
- Pneumonia; a common serious complication of AIDS



2. Blastomycosis

- Caused by *Blastomyces dermatitidis*
- Spores of the fungi enter through the respiratory system
- Primarily affect the lungs
 - Can spread through bloodstream and affect other parts
- Men between ages of 20 and 40 years are the most commonly infected



Pathogenesis...Blastomycosis


- Infection of the lungs is gradual
 - Fever, chills, and drenching sweats develop
 - Chest pain, difficulty breathing, and cough may also develop
 - Can sometimes heal without treatment
 - Skin: warty patches develop surrounded by tiny painless abscesses
 - Bones: painful swellings
 - GUT: prostatitis or painful swelling of epididymis

3. Histoplasmosis

- Caused by *Histoplasma capsulatum*
 - Occurs in soil contaminated with bat or bird droppings
 - Commonly found in temperate, subtropical and tropical zones
 - 50% - 90% of residents in these areas test positive for exposure
 - People who live and work in the vicinity of bat or bird droppings are at increased risk of infection
 - Amphotericin B is the treatment of choice if necessary

...Histoplasmosis

- *Histoplasma capsulatum*: subclinical respiratory infection that only occasionally progresses to severe, generalized disease
- Acquired by inhalation of airborne conidia
- Isolation of the fungus or identification in tissues sample is necessary for diagnosis
- Treatment : Amphotericin B



Pathogenesis...Histoplasmosis


- **Transmission** is through inhalation of conidia
 - Small enough to reach bronchioles and alveoli
- **After inhalation:**
 - Microconidia convert to yeast form, phagocytosis, tubercle formation
 - Severe cases may develop chills, malaise, chest pain, and extensive pulmonary infiltration

5. Coccidioidomycosis

- Caused by *Coccidioides immitis*
- Infection can be symptomatic or asymptomatic
 - Symptomatic form known as Valley Fever
- Arthroconidia of the fungus are inhaled.
 - Small enough to bypass defenses of the upper tract.
 - Lodge directly in bronchioles.
- Fungal outer wall has antiphagocytic properties
 - Prevents elimination

...*Coccidioidomycosis*

- Inhalation of the airborne arthroconidia of *Coccidioides immitis*
- Most cases are subclinical, but when there are predisposing factors such as fatigue and poor nutrition, a progressive disease resembling tuberculosis can result → fever, coughing, weight loss; occasionally fatal



Pathogenesis...coccidioidomycosis

- **Arthroconidia** convert to spherules which grow slowly
 - Completely inhibit phagocytosis
- Disseminated coccidioidomycosis: AIDS patients and on immunosuppressive therapy
- Can also cause a form of coccidioidal **meningitis**
 - Can be fatal if not treated aggressively
- Usually self-limiting and no treatment is required
- Progressive pulmonary infection or infection of central nervous system is treated with amphotericin B

6. Aspergillosis

- Invasive aspergillosis shows a rapid progression to death
- Typically seen in the immunocompromised
 - Particularly patients with leukemia or AIDS
 - Patients undergoing bone marrow transplantation
- Also seen in individuals with preexisting pulmonary disease
 - Chronic bronchitis, asthma, and tuberculosis
 - Fungus produces extracellular proteases, phospholipases, and toxic metabolites

...*Aspergillosis*

- Caused by the fungus *Aspergillus*
 - Widely distributed and found throughout world
 - Dispersal is through inhalation of resistant conidia.
 - Seen more and more in nosocomial infections associated with air-conditioning systems.


...*Aspergillosis*

Many other opportunistic fungi may cause respiratory d/se

Aspergillus fumigatus can cause:

- Allergic bronchopulmonary aspergillosis
- Aspergilloma: in patients with pre-existing lung disease → mass of hyphae produce fungus ball
- Disseminated aspergillosis


...Aspergillosis



Parasitic infections of lower respiratory tract

Paragonimus westermani

- Lung Fluke
- **Adults:** in the lung
- **Eggs:** in the sputum
- **Larval forms:** fresh water snails
- **Metacercariae:** fresh water crabs & crayfish
- Symptoms severe pulmonary paragonimiasis: chest pain , cough, night sweats, pleural effusion, & coughing up blood



Life cycle of *P.westermani*

Paragonimus westermani

Laboratory Diagnosis

1. Finding of eggs in the sputum

- ✓ sputum is usually bloody, mucoid & rusty brown

2. Finding of eggs in aspirates of pleural fluid & in faeces

Treatment: Praziquantel