

Jimma University
Faculty of Health Sciences
School of Medical Laboratory Sciences

Module Name: Respiratory System

Module code: PC

Program: Regular Medicine

Academic Year: 2021/2022

Academic Level: Pre-clerkship I

Instructor: Dr. Muluaem Tadesse (PhD, Assistant professor)

Course Name: Medical Microbiology

Respiratory tract infections (RTI)

Objectives


At the end of this topic, students will be able to;

- Describe common microbes affecting respiratory tract
- Explain the clinical pictures of respiratory tract infections
- Identify diagnostic techniques to diagnose RTI
- Describe common prevention/control methods of RTI

Respiratory tract infections (RTI)


What is the respiratory system?

- The respiratory system consists of
 - the nose,
 - pharynx (FAIR inks),
 - larynx (LAIR inks),
 - trachea (TRAY kee ah),
 - bronchi (BRAHN kye), and
 - Lungs



The primary function of this system is to furnish oxygen for individual tissue cells, and to take away the waste products and carbon dioxide produced by those same cells

Parts of RTI



The Respiratory Tract and Its Defenses

- Most common place for infectious agents to gain access to the body
- Upper respiratory tract:
 - mouth, nose, nasal cavity, sinuses, pharynx, epiglottis, larynx
- Lower respiratory tract:
 - trachea, bronchi, bronchioles, lungs, alveoli
- Defenses
 - Nasal hair
 - Cilia
 - Mucus
 - Involuntary responses such as coughing, sneezing, and swallowing
 - Macrophages
 - Secretory IgA against specific pathogens

Normal Microflora of the Respiratory Tract

- Generally limited to the upper respiratory tract
- Gram-positive bacteria (streptococci and staphylococci) very common
- Disease-causing bacteria are present as normal biota;
 - can cause disease if their host becomes immunocompromised or
 - if they are transferred to other hosts
- Normal biota perform microbial antagonism

Copyright © The McGraw-Hill Companies, Inc. Permission required for reproduction or display.


Respiratory Tract Defenses and Normal Biota

	Defenses	Normal Biota
Upper Respiratory Tract	Nasal hair, ciliary escalator, mucus, involuntary responses such as coughing and sneezing, secretory IgA	<i>Moraxella</i> , nonhemolytic and alpha-hemolytic streptococci, <i>Corynebacterium</i> and other diphtheroids, <i>Candida albicans</i> Note: <i>Streptococcus pyogenes</i> , <i>Streptococcus pneumoniae</i> , <i>Haemophilus influenzae</i> , <i>Neisseria meningitidis</i> , and <i>Staphylococcus aureus</i> often present as “normal” biota.
Lower Respiratory Tract	Mucus, alveolar macrophages, secretory IgA	None

Respiratory tract infections (RTI)

- Upper respiratory tract infection (URTI)
 - Viral, bacterial
- Lower respiratory tract infection (LRTI)
 - Bacterial, viral, fungal, parasite

Diseases of the respiratory system



Upper Respiratory Tract Infections

- Most caused by viruses & usually nothing more than irritation
- Direct invasion of respiratory epithelium results in symptoms corresponding to the area(s) involved
- May spread down and cause more severe infections

Infections of **airway above glottis** **or vocal cords**

- Tonsillitis
- Pharyngitis
- Laryngitis
- Sinusitis
- Otitis media
- Rhinitis

- 1) Cough
- 2) Sore throat
- 3) Running nose
- 4) Nasal congestion
- 5) Headache
- 6) Low-grade fever
- 7) Facial pressure
- 8) Sneezing

Bacterial agents of URTI...

1. *Corynebacterium diphtheriae*

General Characteristics

- Gram-positive bacilli,
- Non Endospore-Forming Bacilli
- Pleomorphic (Curved or straight).
- Non-motile, non-acid fast and aerobic.
- Club-shaped forms = Swelled end
 - Chinese characters
- Found as free-living saprophytes
- Members of the usual flora of humans and animals
 - colonize skin, URT, urogenital tract of healthy people

Virulence Factors *C. diphtheriae*

- Exotoxin as the virulence factor
- Diphtheria toxin is secreted at the site of infection

A: Active fragment

Inhibits protein synthesis

Leads to cell/tissue death


B binds to specific cell membrane receptors

Mediates entry of fragment A


PATHOGENESIS

Entry into nose or mouth

The organism remains in the superficial layers of skin lesions or respiratory tract mucosa, inducing local inflammatory reaction



The major virulence of the organism lies in its ability to produce the potent 62-kd polypeptide exotoxin, which inhibits protein synthesis and causes local tissue necrosis



Within the first few days of respiratory tract infection , a dense necrotic coagulum of organisms, epithelial cells, fibrin, leukocytes and erythrocytes forms, advances, and becomes a gray-brown, leather-like adherent pseudomembrane . Removal is difficult and reveals a bleeding edematous submucosa

Clinical presentations

- ▶ Asymptomatic
- ▶ Respiratory diphtheria
 - Anterior nasal diphtheria
 - Pharyngeal and tonsillar diphtheria
 - Laryngeal diphtheria
- ▶ Cutaneous diphtheria

a. Diphtheriae

Local lesion in URT →

necrotic injury to epithelial cells

→ blood plasma leaks

fibrin network interlacing with
growing bacteria

pseudomembrane

Toxin causes inhibition of protein synthesis
in cells death of eukaryotic cells and
tissues

Diphtheria - pseudomembrane

- Thick grey 'pseudomembrane' composed of fibrin, epithelial cells, bacteria and polymorph neutrophils
- Pseudomembrane may cause blockage, suffocation




Figure 25.20b Microbiology: An Evolving Science
©Princess Margaret Hospital, Hong Kong

Tonsillar and pharyngeal diphtheria: sore throat is the universal early symptom

- Mild pharyngeal injection → unilateral or bilateral tonsillar membrane formation → extend to involve the uvula, soft palate, posterior oropharynx, hypopharynx, or glottic areas
- Underlying soft tissue edema and enlarged lymph nodes: **bull-neck appearance**
 - Classic *cutaneous diphtheria* is an indolent, nonprogressive infection characterized by a superficial, ecthymic, nonhealing ulcer with a gray-brown membrane



URTI....cont'd

b. Tonsilar diphtheria



Lab diagnosis

I. Diagnosis of case: aims at confirming clinical diagnosis

- Throat swabs from the membranes are examined as:
 - a. **Direct smears:** stained with gram and Methylene blue G+ve bacilli with characteristic morphology may be seen
 - b. **Cultures on** loeffler's serum and blood tellurite media. Colonies are picked & stained with gram & Methylene blue
 - c. **Elek test:** rapid diagnosis (16-24 hrs)
 - Results must be confirmed by toxigenicity tests (in vivo & vitro virulence test and tissue culture tests)
 - d. **PCR** - to detect toxin gene (*tox gene*)
 - e. **ELISA** - to detect toxin from clinical isolates

2. Diagnosis of susceptibility by the Schick test:

- Done to determine if a person is susceptible or immune to diphtheria by detecting presence or absence of antitoxin Abs
- ID injection of a small amount of toxin in one forearm & an equal amount of heat inactivated toxin in the other forearm (control)
- If both sites give no reaction => **Schick negative** (person immune)
- If a necrotic area develops within 3 days at the site of injection in the test arm only => **Schick positive** => person susceptible
- If reaction develops in both arms; allergy to proteins of the toxin

Prevention, control and treatment of diphtheria

1. Prophylaxis: Active immunization

- a. Fluid toxoid:** A filtrate of broth culture (toxigenic strain) treated with 0.3% formalin (remove toxicity, immunogenicity)
- b. Aluminium precipitated toxoid:** Fluid toxoid adsorbed onto aluminium hydroxide or aluminium phosphate
- Toxoids are commonly combined with tetanus toxoid & pertussis vaccine (DPT) and given IM to children at the age of 2,4 & 6 month

2. Diphtheria antitoxin serum given without delay

- Antitoxin neutralizes toxin before causes irreversible damage
- Dose: 20,000 - 100,000U (IM or IV); effective with antibiotics

3. Antibiotic

- Penicillin & erythromycin (when given with antiserum; reduce number of diphtheria in throat & decrease incidence of carrier

2. *Streptococcus pyogenes*

- Gram positive cocci in chain
- Group A streptococcus (GAS) with 40 antigenic types
- *S. pyogenes* is β hemolytic
- 90% of cases of pharyngitis
- Flesh eating bacteria
- Most important human pathogen causing disease including:
 - Suppurative conditions/skin infections
 - Throat infections
 - Systemic infections
 - Non-suppurative sequelae
- About 5-15% of normal individuals harbor in their respiratory tract, without signs of disease

Determinants of Pathogenicity--GAB

1. **avoid phagocytosis** - mediated primarily by capsule
2. Adhere to and invade host cells- M protein, lipoteichoic acid,
3. produce toxins and Enzymes:-
4. streptococcal pyrogenic exotoxins,
 - The toxins act as **superantigens**-- with the enhanced release of proinflammatory cytokines.
5. streptolysin S -- an oxygen-stable, nonimmunogenic,
6. streptolysin O--**It is immunogenic** (antistreptolysin O [ASO])
7. streptokinase, **lyse blood clots and fibrin deposits** and facilitate the rapid spread of **S. pyogenes in infected tissues**
8. DNases

Suppurative Streptococcal Disease

1. **Sore throat** and follicular tonsillitis (**pharyngitis**)

- Most common infection; enlarged tonsils with purulent exudates, high fever and enlarged cervical lymph nodes

2. **Scarlet fever**: Disease of children characterized by sore throat and erythematous rash; erythrogenic toxin

3. Pyoderma (impetigo) is a confined, purulent (“pyo”) infection of the skin (“derma”)

4. Erysipelas (erythros, “red”; pella, “skin”) is an acute infection of the skin.

5. Pyoderma (impetigo) is a confined, purulent (“pyo”) infection of the skin (“derma”) that primarily affects exposed areas (i.e., face, arms, legs).

Scarlet fever

Strawberry tongue:
papilla hypertrophy on
the tongue




Fig. 8 Scarlet fever: Initially the tongue is covered by a white exudate through which the papillae project (white strawberry tongue - above). Later the exudate is shed to reveal the bright red inflammation of the underlying tissue (red strawberry tongue - left).

Exanthems
on the skin



Flesh Eating Bacteria

- Cellulitis- leading to Necrotising Fasciitis
- Leads to necrosis of subcutaneous tissues and muscular tissues and adjacent fascia
- Other complications
Toxic shock syndrome





Non Suppurative complications or sequelae

- Post-infection complications of Group A streptococcal disease; Serious complications in pre-antibiotic era
- **Acute rheumatic fever (ARF):**
 - following pharyngitis
 - Morbidity & mortality linked to subsequent disease of heart valve (Rheumatic Heart Disease)
 - Poorly understood pathogenesis with several proposed theories including **cross-reactivity of heart tissues & strep AGNs**
 - M protein cross-reacts heart myosin
 - Autoimmunity and potentially fatal
- **Acute Glomerulonephritis**- following skin infection
 - Immune complex mediated disease
 - inflammation of glomeruli due to Ag-Ab complex deposit on basement membrane
 - Clinically- Hematuria, Proteinuria, Hypertension

Lab diagnosis of streptococcal diseases:

- **Specimens:** swabs from throat or other lesions, pus, or blood (bacteraemia)
- **Direct gram smears:** Gram-positive cocci in chains
- **Culture on BA:** colonies producing complete haemolysis



Bacitracin test:

- Differentiated GAS from other β -hemolytic streptococci (B, C, G) by their sensitivity to bacitracin
- Done by placing filter paper disc containing bacitracin on blood agar inoculated with organism (zone of inhibition)

Antigen detection tests:

- Rapid detection of GAS Ags from throat swab can be done by ELISA or agglutination tests

Serologic tests: to diagnose post-streptococcal d/ses

- ASO test

Other URTI causing bacteria

- *Haemophilus influenza*
- *Moraxella catarrhalis*
- *Streptococcus pneumoniae*
- *Staphylococcus aureus*


Viral causative agents of URTI

1. Corona viruses

- *Corona* = crown like structure, pleomorphic, spiked
- Non-segmented, linear, ssRNA
- Enveloped +ve stranded RNA
- mRNA encased in nucleocapsid
- Lipid Bilayer – Soap works to disrupt this!
- *Causes diseases in Man, animals and birds*

three glycoproteins:

- ✓ **S - Spike protein:** receptor binding, cell fusion, major antigen
- ✓ **M - Membrane protein:** transmembrane - budding & envelope formation
- ✓ **HE – Hemagglutinin:** hemagglutination, acetylsterase activity



Corona viruses– pathogenesis

- Normal human coronaviruses cause 5-10% of common cold/URIs, with outbreaks to 30% of common cold
- Display a tropism for **epithelial cells** of the **respiratory or gastrointestinal tract**
- Remain limited to the upper respiratory tract
- a slow , patchy destruction of ciliated epithelial cells and the loss of beating cilia → disease development
- **High frequency of recombination during replication - evolution of new strains**

2. Rhinoviruses

- Common cold viruses
- Mild URT illnesses
- Nasal secretions, throat and oral secretions

Properties

- inactivated by low pH (pH 3)
- More thermo stable than enteroviruses and survive for days on environmental surfaces
- Infect only **humans, chimpanzee**
- Replication is limited to the **epithelium of the nasal mucosa**
- Human cell lines - **grow better at 33°C** (the temperature of the nasopharynx in human)
- **Common cold viruses with >100 antigenic types**

Rhinoviruses

Clinical findings


- Common cold
- IP → 2-4 days
- Acute illness usually lasts for 7 days
- Nonproductive cough may persist for 2-3 weeks
- Sneezing, nasal obstruction, nasal discharge & sore throat
- headache, malaise, mild cough, & a chilly sensation
- Little or no fever
- The nasal and nasopharyngeal mucosa become red and swollen and the sense of smell is decreased
- **No specific treatment/vaccine**

3. Adenoviruses

- Family: **Adenoviridae**
- Non-enveloped, dsDNA virus
- Linear ds DNA Genome
- 51- human adenoviruses, grouped in 6-species (A-F)
- Encode proteins that block MHC class I expression
- Cough, nasal congestion, fever, sore throat and rarely pneumonia
- Rounding, enlargement and aggregation of affected cells into grape-like clusters in cell cultures

Pathogenesis- Adenovirus

- Infect and replicate in **epithelial cells** of respiratory tract, eye, gastrointestinal tract, urinary bladder, and liver.
- Usually do not spread beyond the **regional lymph nodes**
- Group **C viruses** persist as latent infections for years in **adenoids and tonsils** and shed in the **feces** for many months after the initial infection



URTI....cont'd

Other URTI causing viruses

- Influenza viruses (type A and B)
- Parainfluenza virus
- Coxsackie A virus
- Epstein Bar Virus
- Herpes Simplex Virus
- Respiratory Syncytial Virus
- Rubella virus
- Varicella -zoster virus
- Cytomegalo virus
- Human metapneumovirus

Summary

- Non-specific, often combinations of viral & bacterial infections
- Diagnosis is mainly based on clinical manifestations
- Treatment: to maximize relief of most prominent symptoms
- Most are self-limiting but
 - Doctor, family & patient factors - over prescribe antibiotic
 - Treat only proven group A strep with antibiotic
- Common cold, sinusitis, pharyngitis, epiglottitis & laryngitis

Lower respiratory tract infections (LRTI)

Lower respiratory tract infections (LRTI)

- Lower respiratory tract
 - Trachea, bronchi, bronchioles, and alveoli in the lungs
 - Most are bacterial causes
 - Far more serious than upper respiratory tract infections
 - Many of the microbes infecting upper respiratory system can also infect the lower respiratory system
 - Bacterial, viral, fungal and parasitic infections can cause Inflammation of the LRT

Bacterial infections of lower respiratory tract

- Bacterial pneumonia
- Pertussis
- Inhalation anthrax
- Q fever
- Tuberculosis

1. Bacterial pneumonia

- Inflammatory condition of the lung in which fluid fills the alveoli
- Can be caused by a wide variety of different microorganisms
- Viral pneumonias are usually milder than bacterial
- One of the most serious lower respiratory tract infections

* Classification of pneumonia:

1. Community-Acquired Acute Pneumonia:

- *Streptococcus pneumonia*
- *Haemophilus influenza pneumonia*
- *Staphylococcus aureus pneumonia*
- *Legionella pneumophila pneumonia*

2. Community-Acquired Atypical Pneumonia:

- *Mycoplasma pneumonia*
- Viral pneumonia: Respiratory syncytial virus, para-influenza virus (children); influenza A and B (adults); adenovirus.

3. Nosocomial Pneumonia.

4. Aspiration Pneumonia.

5. Chronic Pneumonia.

- Granulomatous: *Mycobacterium tuberculosis* and atypical mycobacteria.
- *Nocardia*.
- *Actinomyces*.
- *Histoplasma capsulatum*.

....Bacterial pneumonia

■ Nosocomial pneumonia

- Occurs ~ 48 hours after admission to hospital
- Usually associated with *Staphylococcus aureus*
- Also caused by gram-negative bacteria
- Particularly difficult to deal with antibiotic resistant

■ Community-acquired pneumonia

- Usually presents as a lobar pneumonia
- Accompanied by fever, chest pain, and production of purulent sputum

Community-acquired pneumonia:

- Require enough pathogens to overwhelm resident defenses
- Establishment of an infection in the lungs depends on:
 - ❖ Number of pathogen entering and competence of immunity
- Bacterial causative agents of LRTIs
 - *Streptococcus pneumoniae*
 - *Haemophilus influenzae*
 - *Staphylococcus aureus*
 - *Legionella pneumophila*
 - *Mycoplasma pneumoniae*
 - *Chlamydia pneumoniae*,
 - *chlamydia psittaci*

Pneumonia...cont'd

1. *Streptococcus pneumonia*

- Encapsulated, Gram positive coccus, Lancet shaped, **arranged in pairs**
- 5-50% of all people carry it as normal flora in pharynx
- **Very delicate, does not survive long outside of its habitat**
- Virulent strains are covered with polysaccharide capsule
- All colonies undergo autolysis
- α hemolytic
- Progress to alveolus and associate with specific alveolar cells (produce choline-containing surfactant)
- Can also cause sinusitis, otitis media, meningitis

Streptococcus pneumoniae

Virulent factors

a. **Secretory IgA protease**

b. **Teichoic acid and peptidoglycan fragment**

activate alternate complement system which results production of C5a, C3a that result inflammation

c. **Amidase** increases release of cell wall fragments

Autolytic enzyme

d. **Pneumolysin**

Cytotoxin (bind to cholesterol in CM and form pore) destroy ciliated epithelial cells and phagocytic cells

Activates classical complement system $\Rightarrow$ C5a, C3a $\Rightarrow$ activate macrophage $\Rightarrow$ $\uparrow$ Inf. cytokines $\Rightarrow$ fever, tissue damage etc

■ Pathological effect is mainly due to:- Activation of inflammatory cascades--Host responses, Toxic factors

Clinical diseases *Streptococcus pneumonia*

- Colonizes oropharynx → then can spread to lung sinus → middle ear
→ can be transported to blood and then different organs (meningitis, endocarditis, septic arthritis)
- Causes 60-70% of all ***bacterial pneumonias***
 - One of the most common causes of **community acquired disease**
 - multiply & induce an overwhelming inflammatory response
 - Acute chills, fever, productive cough with blood mixed sputum
- Meningitis – Most common causes of bacterial meningitis
- Common cause of sinusitis, **otitis media**
- Can cause a variety of systemic infections like bacteremia and endocarditis

Community-acquired *Bacterial* pneumonia

TWO types:

A . Atypical pneumonia

- Coughing without sputum
- Caused by a variety of bacteria
- Bacterial pneumonia can progress to the production of lung abscesses
- Caused by **viruses, Mycoplasma, Chlamydia**

Atypical Pneumonia

- Fever and malaise precede respiratory symptoms few days
- Severe headache, malaise, anorexia
- No localized signs on chest exam,
- No consolidation on chest x-ray
- Spleen may be enlarged
- WBC normal, cultures negative
- No improvement with Penicillin

Atypical Pneumonia

- Atypical Pneumonia (community-acquired)
 - Mycoplasma
 - Sporadic or epidemics
 - Viruses
 - Influenza, Parainfluenza, Adenovirus, respiratory syncytial virus, measles, chicken pox
 - Chlamydia
- Morphology:
 - Patchy or involve whole lobe
 - Inflammation is confined to the alveolar walls
 - Widening of alveolar walls by edema, mononuclear cell infiltration (lymphocytes, plasma cells, macrophages)

B. Typical *pneumonia*

Lobar Pneumonia:

- *S. pneumoniae* that affects part of a lobe in the lung or it may affect more than one lobe

Bronchial Pneumonia:

- Pneumonia spreads to several patches in one or both lungs
- Most prevalent in infants, young children and aged adults
- Cough (with or without mucus), chest pain, rapid breathing, and shortness of breath
- Transmitted by respiratory droplets

Treatment...Bacterial pneumonia

- Course of treatment depends on:
 - Severity of the infection
 - Type of organism causing the infection
- Most common pathogen: *Streptococcus pneumoniae*
 - Treated with penicillin, amoxicillin-clavulanate, and erythromycin

Hospital acquired (Nosocomial) pneumonia

- Nosocomial, or hospital-acquired pneumonias are defined as pulmonary infections acquired in the course of a hospital stay.
- Nosocomial infections are common in hospitalized persons with severe underlying disease, those who are immunosuppressed, or those on prolonged antibiotic regimens.
- Those on mechanical ventilation represent a particularly high-risk group, and infections acquired in this setting are given the distinctive designation *ventilator-associated pneumonia*.

* Causative agents:

1. Gram-negative rods (Enterobacteriaceae and *Pseudomonas* spp.)
2. *Staph. aureus*.

Staphylococcus aureus pneumonia

- *S. aureus* is an important cause of secondary bacterial pneumonia in children and healthy adults after viral respiratory illnesses (e.g., measles in children and influenza in both children and adults).
- Staphylococcal pneumonia is associated with a high incidence of complications, such as lung abscess and empyema.
- Staphylococcal pneumonia occurring in association with right-sided staphylococcal endocarditis is a serious complication of *intravenous drug abuse*.

Klebsiella pneumoniae

- *K. pneumoniae* is the most frequent cause of gram negative bacterial pneumonia.
- *Klebsiella*-related pneumonia frequently afflicts debilitated and malnourished persons, particularly *chronic alcoholics*.
- Thick and gelatinous sputum is characteristic, because the organism produces an abundant viscid capsular polysaccharide, which the patient may have difficulty coughing up.

Pseudomonas aeruginosa

- *Pseudomonas* pneumonia is common in persons who are neutropenic, usually secondary to chemotherapy; in victims of extensive burns; and in patients requiring mechanical ventilation.
- *P. aeruginosa* has a propensity to invade blood vessels at the site of infection, with consequent extra-pulmonary spread; *Pseudomonas* bacteremia is a fulminant disease, with death often occurring within a matter of days.

Tuberculosis

- Caused by *Mycobacterium tuberculosis complex*
- Slender, slightly curved or straight rod-shaped organisms
- Non-motile
- Do not form spores
- Strictly aerobic
- do not form capsules, flagella or spores
- Extremely high lipid content
- possess mycolic acids & a unique type of peptidoglycan
- grow slowly

Epidemiology of TB

~ 1 in 3 people in world infected

~ 9.4 million new cases of active TB/year

About 2 million deaths/year

30 high-burden countries for TB, TB/HIV and multidrug-resistant TB

TB kills an estimated of 32,000 Ethiopians every year (more than 80 people per day)







Figure 21.12 Microbiology: A Clinical Approach (© Garland Science)

Natural History of TB Infection



Latent TB Infection (LTBI)



Person:

- Not ill
- Not contagious
- Normal chest x-ray
- Usually the tuberculin skin test is positive



Germs:

- **Sleeping but still alive--?**
- Surrounded (walled off) by body's immune system


Active TB Disease

Germ:


- Awake and multiplying
- Cause damage to the lungs

Person:

- Most often feels sick
- Contagious (before pills started)
- Usually have a positive tuberculin skin test
- Chest X-ray is often abnormal (with pulmonary TB)



Granuloma breaks down and tubercle escape and multiply



Pathogenesis...Tuberculosis

- *MTB* cell wall interferes with macrophage function and immune cell activation
 - Inhibits the formation of the phagolysosome
- This allows *MTB* to escape into the cytoplasm where it:
 - Increases in number & eventually spreads to lymph nodes, enters blood and distributed throughout body
- Fever, Fatigue, Weight loss, Chest pain, Shortness of breath, Congestion with coughing

Pathogenesis...Tuberculosis

- Two basic types of tuberculosis

- Primary



- Follows initial exposure to the pathogen

- Secondary

- Can occur years later

TB can spread to d/t parts of the body

1. Lungs (85% all cases)
2. Pleura
3. Central nervous system
 - (e.g., brain, meninges)
4. Lymph nodes
5. Genitourinary system
6. Bones and joints
7. Disseminated
 - (e.g., miliary)



Evaluation for TB

- Medical history
- Physical examination
- Mantoux tuberculin skin test/ IGRAs
- Chest x-ray
- Histopathology/cytology
- Bacteriologic exam
 - smear
 - Culture
 - NAAT/PCR

Symptoms of TB

- Productive prolonged cough(>2wks)*
- Chest pain
- Hemoptysis
- Fever and chills
- Night sweats
- Fatigue
- Loss of appetite
- Weight loss

Laboratory Diagnosis of Tuberculosis

Specimens: Sputum, Urine, pleural fluid, peritoneal fluid, C.S.F., gastric lavage, lymph node biopsy, tissue biopsy

Diagnosis of TB Disease

- Signs and Symptoms consistent with TB
- Chest X-ray
- Clinical Judgment
- Bacteriology
 - AFB smear microscopy
 - Nucleic Acid Amplification Testing (NAAT)
 - Culture and Identification
 - Drug susceptibility testing (DST)

4. Molecular Techniques

- Allow rapid detection of MTBc DNA
- Routine PCR: Gene amplification
- Real Time PCR: Gene amplification plus quantification
- Sensitivity: 80 - 85%, specificity ~ 99%
- **GeneXpert MTB/RIF assay**
 - Is the novel diagnostic tool endorsed by WHO in 2012---
 - **Detects TB and its resistance to rifampicin**
- It is simple and rapid---results within 2hrs



Mantoux test (TST)


Tests for exposure-- LTBI

Tests for hypersensitivity(type IV)

Intracutaneous injection of purified protein derivative (PPD)

Induration measured after 48-72h

Copyright © The McGraw-Hill Companies, Inc. Permission required for reproduction or display.



Treatment of active TB Disease

- Include four 1st-line drugs in initial regimen
 - Isoniazid, Rifampin, Pyrazinamide, Ethambutol
- 6 months of combination therapy (**2RIPE (intensive /4RI -continuation)**)
- **Prolonged treatment** --slow response of TB treatment
 - ❖ Most bacilli are found intracellularly
 - ❖ Caseous material interfere with the drug
- **Combination of drugs:** to reduce toxicity due to prolonged course of treatment and rapid emergence of resistant strains
- **Directly Observed Therapy (DOT)**-- Promote adherence and ensure treatment completion

Multidrug resistant TB- a MTBc strain resistant to RIF and INH

- Multi-drug resistant TB treatment: much more expensive, toxic and can go on for 2 years.

Prevention

A. Public health measures

- Early diagnosis and treatment until they become non-infectious
- Control of transmission

B. Vaccination:

- A live-attenuated vaccine – BCG commonly used
- Prepared from bovine strain with a fixed low virulence
- Vaccine is given in a single dose of 0.1ml ID in the deltoid region to stimulate hypersensitivity and CMI against infection

Pertussis (whooping cough)

- A highly contagious acute bacterial infection caused by the bacilli *Bordetella pertussis*
- *It continues to be an important respiratory disease afflicting unvaccinated infants*
- **Transmission:** through the respiratory route in the form of droplet infection
- Adolescents and adults are the reservoir.
- No animal or insect reservoir


Bordetella pertussis

- Short, gram-negative bacilli
- **Pili & haemagglutinin:** adheres to ciliated epithelia of RT
- **Pertussis toxin** - promotes phagocytosis
- **Tracheal cytotoxin** inhibit DNA synthesis in ciliated cells
- **LPS:** causes damage to epithelial cells of the RT

B. pertussis

- Adheres and multiplies rapidly on epithelial surface of trachea & bronchi and interfere with ciliary action.
- Liberate toxins & substances that irritate surface cells, causing cough & marked lymphocytosis.
- Cough is intermittent with bursting attacks accompanied by whoop followed by vomiting
- Complications: bronchopneumonia, subconjunctival or cerebral hemorrhage due to paroxysms of severe cough.
- **Whooping cough: whooping sound made when gasping for air after a fit of coughing**

Clinical Progression of Pertussis

	Incubation	Catarrhal	Paroxysmal	Convalescent
Duration	7–10 days	1–2 weeks	2–4 weeks	3–4 weeks (or longer)
Symptoms	None	Rhinorrhea, malaise, fever, sneezing, anorexia, inflammation of respiratory mucosal memb.	Repetitive cough with whoops, vomiting, leukocytosis	Diminished paroxysmal cough, development of secondary complications (pneumonia, seizures, encephalopathy) or death
Bacterial culture	 Most infectious, but generally not yet diagnosed			

Whooping cough

Diagnosis:

- Isolation of bacteria: positive during the 1st week only
- Culture from mucus or droplets: **Mercury drops colony**
- Serology: using specific anti-sera or specific antigens

Prophylaxis:

- Heat killed vaccine to children during 1st year of life in combination with diphtheria and tetanus (DPT)

Treatment:

- Antibiotics: erythromycin effective if given early
- Recovery is followed by long lasting immunity

Inhalation anthrax

- Zoonotic disease of herbivorous livestock



- Caused by *Bacillus anthracis*
- Gram-positive, endospore-forming, motile rods
- Mostly saprobic
- Aerobic and catalase positive
- Versatile in degrading complex macromolecules
- Source of antibiotics

Inhalation anthrax

- Produces a fulminate pneumonia
 - Comes suddenly with great severity
 - Leads to respiratory failure and death
- Anthrax primarily a disease of herbivores
 - Acquired from spores found in pastures
 - If spores inhaled, anthrax can occur in respiratory tract
- Infection is infrequently seen in healthy individuals
 - Usually presents as localized lesions where it occurs
- Recent interest: as a biological weapon

Virulence factors include

Capsule, plasmid encoded

Anthrax toxin (AB toxin)

1. Protective antigen (PA)
2. Edema factor (EF)
3. Lethal factor (LF)

PA+LF → lethal activity

EF+PA → edema

EF+LF → inactive

PA+LF+EF → edema & necrosis; lethal

Pathogenesis...Inhalation anthrax

- **The infection begins with the inhalation of the anthrax spore**
- Spores germinate in human tissues
- Antiphagocytic properties of the capsule aid its survival and growth in large numbers
 - Pathogenesis results from the powerful **exotoxin** produced
- Symptoms include:
 - Nonspecific malaise, mild fever, nonproductive cough
 - Progressive respiratory distress and cyanosis
- Antibiotic therapy can be successful
 - *B. anthracis* is susceptible to penicillin
 - Doxycycline and ciprofloxacin are alternative prophylactics

Q fever

- A zoonotic infection seen world wide
- Primarily it affects lung
- Caused by *Coxiella burnetii*
- Small pleomorphic coccobacilli
- Obligate intracellular, gram negative bacterium
- Multiply in phagolysosome (the acidic pH environment activates the bacterial metabolic machinery)
- Grows well in placenta of animals and large numbers can be transmitted by inhalation during animal births



Q fever


- Bacteria is excreted in: Feces, urine and milk of infected animals
- Released into Environment ;
- During birthing the organisms are shed in high numbers in amniotic fluids and the placenta
- 10^9 bacteria per gram of placenta

Transmission can also be by ingestion
of unpasteurized milk
Mild hacking cough and patchy
interstitial pneumonia

Do not touch!



Inhalation of contaminated aerosols



On long run 1-5% acute Q fever cases develops into chronic Q fever

Acute Q fever

60% cases → Asymptomatic

40% cases → Symptomatic

(95-98) % cases → Mild (Undiagnosed)

(2-5) % cases → Severe (Hospitalized)

- Severe fever
- Pneumonia
- Hepatitis

Predisposing factors

- Immunosuppression
- Pregnancy
- Heart valve lesions
- Vascular abnormalities

Chronic Q fever

Valve lesion (2% cases)

- Endocarditis (30-60% within two years)

- Chronic fatigue syndrome
- Osteomyelitis
- Chronic hepatitis
- Interstitial lung disease
- Repeated abortions

Small cell variants

Spores

Viral infections of the lower respiratory tract

Viral infections of lower respiratory tract

- Majority of infections in the lower respiratory tract are caused by:
 - Influenza virus
 - Respiratory syncytial virus
- Common characteristics of infection are:
 - Short incubation period of 1 to 4 days
 - Transmission from person to person
- Transmission can be direct or indirect
 - Direct – through droplets
 - Indirect – through hand transfer of contaminated secretions

Why has viral pneumonias become important?

- Increase in immunocompromised patients and population at-risk groups
- Discovery of new respiratory viruses
 - Human metapneumovirus
 - Corona viruses -
 - Hantavirus
 - Human bocavirus
- Emergence of new viruses
 - Severe acute respiratory syndrome (SARS)
 - Avian influenza A (H5N1) virus,
 - 2009 pandemic influenza A (H1N1) virus

1. Severe Acute Respiratory Syndrome (SARS)

- New emerged respiratory disease (2002)
- Symptoms are initially usually mild but latter patients may develop dry non-productive cough and breathing may become difficult (dyspnea)
- Shortness of breath → hypoxia
- Respiratory distress leads to death in 3-30 % of cases
- Laboratory tests show a reduction in lymphocyte and rise in aminotransferase activity - damage to liver

SARS... cont'd


SARS Laboratory findings: CDC recommendations

- Chest radiograph, pulse oximetry, tissue culture
- Sputum gram stain/culture (for Legionella and pneumococcal)
- Testing for viral respiratory pathogens (influenza A, B, RSV)
- Nucleic acid assays to detect SARS-CoV
- ELISA to detect SARS-CoV antibodies
- RT-PCR to detect SARS-CoV infection

SARS-CoV-2 (COVID-19)

- Coronavirus disease 2019 (COVID-19) is an infectious respiratory tract infection caused by SARS-CoV-2
- On 30 January 2020 WHO declared COVID-19 a public health emergency of international concern
- On 11 March declared it as a global pandemic

- Single-stranded RNA viruses
 - Enveloped positive sense
- SARS CoV-2 contains four structural proteins




Transmission, pathogenesis, replication of SARS-CoV-2 (COVID-19)''''''


- Modes of transmission

- Droplets transmission
- Contact transmission
- Aerosol transmission....????

Transmission Cycle of SARS CoV 2



Replication of Coronavirus (SARS-CoV-2)




Signs and Symptoms of COVID-19



Fever



Cough



Shortness of breath

The most commonly reported symptoms include:

1. Fever
2. Dry cough
3. Myalgia or fatigue
4. Dyspnea (shortness of breath)
5. Respiratory issues

The less common symptoms include:


- The incubation period, the duration from the exposure to the virus to the onset of symptoms ranges from 1-14 days (average 5 days).
- The virus is most contagious for the first three days after the appearance of symptoms.

1. Headache
2. Aches and pains
3. Sore throat
4. Diarrhea
5. Hemoptysis
6. Runny nose
7. Phlegm-producing cough

Risk factors of COVID-19

- The primary risk factor associated with coronavirus is age.
 - A case fatality rate of 0.32% is seen in those aged below 60 years
 - 6.4% in those aged ≥ 60 years
 - as high as 13.4% in those aged more than 80 years.
- Underlying disorders
 - hypertension, chronic obstructive pulmonary disease, diabetes, cardiovascular disease),
 - are found to be at more risk of developing severe symptoms.

Basic diagnostic methods



2. Respiratory syncytial virus

- Most important cause of lower respiratory tract illness in **infants and young children**
- Outranking all other microbial pathogens as the cause of **bronchiolitis and pneumonia** in infants under 1 year of age

Pathogenesis

- ✓ Replication occurs initially in epithelial cells of the **nasopharynx**
- ✓ Spread into the lower respiratory tract, probably carried there by secretions
- ✓ Viremia **has not been detected**
- ✓ An intact immune system seems to be important in resolving an infection
- ✓ Spread outside the respiratory epithelium (kidney, liver, etc)
 - in several fatal infections in individuals who lacked cell- mediated immunity

3. Influenza viruses

- Orthomyxovirus, enveloped
 - The single stranded RNA genomes
 - Spherical, Pleomorphic
 - Febrile respiratory d/se with systemic symptoms (**flu**); fever, chills, cough, sore throat, runny nose, muscle or body aches, headache, fatigue (tiredness) & vomiting
 - Type A infects humans, swine, horses, whales, birds
 - ✓ Primary reservoirs (birds); infection is mostly asymptomatic, virus replicate in lungs and intestinal mucosa; shed in feces => human respiratory infection
 - Influenza B & C are human viruses; do not infect birds

Influenza... cont'd

- **Segmented ssRNA virus**; allows a high rate of mutation
 - Hemagglutinin (HA)
 - Neuraminidase (NA)
 - M proteins (matrix protein-M1, membrane protein-M2)
 - Nucleoprotein (NP)
 - Polymerase proteins (PB2, PB1, PA)
 - Non structural proteins (NS1, NS2)
- Two glycoproteins (**hemagglutinin (HA) & neuraminidase (NA)**)
 - derived from the host cell
 - exposed as spikes
 - determine antigenic variation of influenza viruses and host immunity
 - undergo antigenic variation independent of each other

Influenza... cont'd






Figure 21.17 Microbiology: A Clinical Approach (© Garland Science)

Influenza... cont'd

Hemagglutinin

- Required for virus binding to cell surface glycolipids and glycoproteins , 15 HA subtypes
- Responsible for penetration; Abs to HA neutralize virus

Neuraminidase

- Removes sialic acid from glycoconjugate; aiding spread and prevents virus clustering at cell surface upon release
- High concentration of anti-NA antibody are necessary for virus neutralization; 9 NA subtypes

Pathogenesis...Influenza

- Influenza virus prefers the respiratory epithelium
 - Viremia is rare
- Virus multiplies in the ciliated cells of lower respiratory tract
 - Results in functional and structural abnormalities
- Cellular synthesis of nucleic acids and proteins is shut down
- Ciliated and mucus-producing epithelial cells are shed.
 - Substantial interference with clearance mechanisms
 - Localized inflammation; both in URT and LRT

Pathogenesis: *Influenza... cont'd*

- Local symptoms result from epithelial cell damage, including ciliated and mucus-secreting cells
- Systemic symptoms: interferon and lymphokine response
- Infected people predisposed to bacterial super infection
- HA & NA of Influenza A: undergo **major (reassortment: shift)** and **minor (mutation: drift)** antigenic changes to ensure presence of immunologically naïve, susceptible people

Pathogenesis: *Influenza... cont'd*

- Antigenic Drift → Minor antigenic changes
 - accumulation of point mutations in the gene
 - amino acid changes in the protein
 - alter antigenic sites on the molecule
 - a virion can escape recognition by the **host's immune system**
- Antigenic Shift → Major antigenic changes in HA or NA
 - a new subtype
 - segmented nature of the genome
 - cell is co-infected by 2 different viruses of a given type
 - mixtures of parental gene segments may be assembled in to progeny virions
 - **genetic reassortment** → sudden changes in viral surface antigens
- Explains the epidemiological features of influenza
- Poses significant problems for vaccine development

Lab diagnosis: *Influenza... cont'd*

Diagnosis:

- Cell or tissue culture
- Hemadsorption
- Viral antigen detection

Treatment: *Influenza...cont'd*

- Two basic approaches
 - Symptomatic care & anticipation of potential complications
- Best treatments include:
 - Rest and fluid intake, cough suppressants
 - Conservative use of analgesics for myalgia and headache
- **Amantidine & rimantadine** (diagnosed within 12-24 hrs) (M2)
- Zanamivir and tamiflu (oseltamivir) target neuraminidase

Prevention: *influenza...cont'd*

Vaccination

- Trivalent: two current A strains and one current B strain
- Formalin fixed wild type virus approved for parenterally administered vaccination.
- Live attenuated vaccine: temperature sensitive recombinant bearing relevant HA and NA genes
- Must anticipate shift and drift in order to identify appropriate vaccine strain

Fungal infections of the Lower respiratory tract

Fungal infections of lower respiratory tract


- Fungal spores are easily inhaled and may germinate in LRT
- Two major factors govern incidence and spread of fungal infection
 - Ubiquity of the infectious organisms
 - Found in soil
 - Resident flora
 - Adaptive immune response
 - Usually keeps these infections under control
 - Immunocompromised patients at much greater risk
- Incidence is being increased in recent years. Why?

1. Histoplasmosis

- Intracellular infection of the RES caused by *Histoplasma capsulatum*. Endemic in parts of USA
- Also called **Darling's disease**; 1st described by Samuel Darling.
 - "histio" within histiocytes
 - "plasma" resembled plasmodium.
- Present in soil, rotting areas and in feces of chicken, bats & other birds. (high N₂ content)

Histoplasma capsulatum

Copyright © The McGraw-Hill Companies, Inc. Permission required for reproduction or display.





Pathogenesis & Pathology

Inhalation of conidia or mycelial fragments



Converted into yeast in alveolar macrophages



Localized granulomatous inflammation



Granuloma with or
without caseation



assist in
dissemination to RES

- Involves all phagocytic cells of RES, cytoplasm being studded with fungal cells.



Clinical features


- Resembles TB - mainly asymptomatic
- Clinical types -
 1. Pulmonary
 2. Cutaneous & mucocutaneous
 3. Disseminated histoplasmosis - commonly seen in children below 2 yrs & adolescents
 - individuals with HIV are at a greater risk.

Laboratory Diagnosis

Specimen - sputum, BM, LNs, scrapings from lesions, biopsy & peripheral blood.


Direct Examination

- Blood smear - Giemsa or Wright stains.
 - small, oval yeast like cells, 2-4 μ within mononuclear or polymorphonuclear cells, narrow neck budding.
- Fluorescent Ab technique.



Fungal Culture

- SDA , BHI at 25° & 37°C.
- LPCB - White cottony mycelia with large (8-20 μ) thick walled, spherical spores with tubercles or finger like projections -
Tuberculate
Macroconidia.





Immunodiagnosis

- **Histoplasmin skin test** - I.D. test with 0.1 ml histoplasmin Ag - DTH response.
- **Serological tests** - LA

* titer of 1:32 or higher or 4-fold increase in titer of Abs is significant.



Treatment & Prophylaxis

- Amphotericin B - disseminated & other severe forms.
- Oral Itraconazole
- Regular cleaning of farm buildings, chicken houses for prevention.



BLASTOMYCOSIS

- Also called as *Gilchrist's disease* or *Chicago disease* due to its endemicity in N.America (N.American blastomycosis)
- Caused by Inhalation of the spores of *Blastomyces dermatitidis*
- Causes suppurative & granulomatous infection



Clinical features

- 1° infection resembles TB or histoplasmosis.
- Clinical types:
 1. Pulmonary
 2. **Cutaneous** - commonest form, hence the name "dermatitidis".
 - seen over exposed parts like face, neck & hands.
 3. Disseminated type - form multiple abscesses in different parts like bone, genitourinary system, breast etc

BLASTOMYCOSIS






Laboratory Diagnosis


Specimen - sputum, BAL, biopsy or pus from abscesses, urine.

Direct Examination

- **Wet mount** - KOH, CFW : double contoured, thick walled, multinucleate giant yeast cells with broad base budding daughter cells.



The yeast is round with a doubly refractive wall and a single broad-based bud.





Blastomyces dermatidis. **A:** Yeast with a broad-based bud at 37 °C. **B:** Mold with microconidia at 20 °C. (Reproduced, with permission, from Brooks GF et al: *Medical Microbiology*, 19th ed. Originally published by Appleton & Lange. Copyright © 1991 by The McGraw-Hill Companies, Inc.)



Fungal Culture

- Very slow growth - 2 to 4 weeks.
- Tissue & cultures at 37°C shows budding yeast cells.
- At 25°C - fine, branched septate hypha with **conidia** measuring 2-10 μ located **on short terminal or lateral branches**.






Diagnosis

- Immunodiffusion precipitation bands.
- EIA / RIA
- Skin test using blastomycin


Treatment & Prophylaxis

- Initial phase - Oral Ketoconazole & Itraconazole
- Life threatening infections - AMB



COCCIDIOIDOMYCOSIS

- Infection of the respiratory system caused by *Coccidioides immitis*.
- **Most virulent** of all the fungal pathogens but **not contagious**.
- More prevalent in western hemisphere.
- Fungus present in soil & in rodents.
- Infection occurs by
 - inhalation of arthroconidia or
 - reactivation of latent infection in immunocompromised patients.



Coccidioidomycosis

C. immitis is found in soil --enhanced by bat and rodent dropping.


C. immitis is probably the most virulent

Inhalation of a few arthroconidia produce 1^o infection which result asymptomatic or a self limited flue like illness

- Spores transform to **spherules** in lung, elicit cellular immunity as per TB

Like syphilis and tuberculosis coccidioidomycosis causes a wide varieties of lesions and has been called **the great imitator**.

Disseminated infection (Extra pulmonary infection) includes skin, soft tissues, bones, joints, and meninges



Clinical features



- Many develop influenza like fever - Valley fever or Desert Rheumatism
- < 1% develop chronic progression disseminated disease -
 - skin (commonest) : granuloma, cold abscess.
 - osteomyelitis & synovitis
 - CNS (meningitis)

1. *Pneumocystis pneumonia* (pcp)

- A lethal pneumonia
 - Common in AIDS patients
- Caused by the fungus *Pneumocystis (carinii) jiroveci*
 - Never been grown in culture
 - Most information comes from clinical information of patients

Pneumocystis Pneumonia

- *Pneumocystis jirovecii* (formerly *P. carinii*) is sometimes found in healthy human lung
- Pneumonia; a common serious complication of AIDS




6. Aspergillosis

- Invasive aspergillosis shows a rapid progression to death
- Typically seen in the immunocompromised
 - Particularly patients with leukemia or AIDS
 - Patients undergoing bone marrow transplantation
- Also seen in individuals with preexisting pulmonary disease
 - Chronic bronchitis, asthma, and tuberculosis
 - Fungus produces extracellular proteases, phospholipases, and toxic metabolites

...Aspergillosis

- Caused by the fungus *Aspergillus*
 - Widely distributed and found throughout world
 - Dispersal is through inhalation of resistant conidia.
 - Seen more and more in nosocomial infections associated with air-conditioning systems.

Fungal spores enters through respiratory tract



Aspergillosis

The *Aspergillus* species can cause a variety of clinical syndromes

1. Pulmonary aspergillosis

a) Allergic asthma

b) Bronchopulmonary aspergillosis

c) Aspergilloma

2. Invasive aspergillosis

3. Superficial infections

...*Aspergillosis*

Many other opportunistic fungi may cause respiratory d/se

Aspergillus fumigatus can cause:

- Allergic bronchopulmonary aspergillosis
- Aspergilloma: in patients with pre-existing lung disease → mass of hyphae produce fungus ball
- Disseminated aspergillosis

...Aspergillosis

