


Lower respiratory tract infections (LRTI)

- Lower respiratory tract
 - Trachea, bronchi, bronchioles, and alveoli in the lungs
 - Most are bacterial causes
 - Far more serious than upper respiratory tract infections
 - Many of the microbes infecting upper respiratory system can also infect the lower respiratory system
 - Bacterial, viral, fungal and parasitic infections can cause Inflammation of the LRT

Bacterial Infections of the Lower Respiratory Tract



- **Pertussis**

- *Bordetella pertussis*

- **Pulmonary Anthrax**

- *Bacillus anthracis*

- **Pneumonia**

- *Streptococcus pneumoniae*
- *Mycoplasma pneumoniae*
- *Chlamydophila pneumoniae* and
- *Legionella pneumophila*
- *Coxiella burnetii*
- *Klebsiella pneumoniae*
- *Pseudomonas aeruginosa*
- *Moraxella catarrhalis**
- *Haemophilus influenzae**
- *Staphylococcus aureus**

- **Tuberculosis/TB-Like**

- *Mycobacterium tuberculosis*
- Other acid-fast bacilli


Pneumonia:

Download

Definition:

Pneumonia is an inflammatory condition of the lung—especially affecting the **microscopic air sacs** (alveoli), and the **parenchyma** of the lung.


Lung parenchyma refers to **alveolar** tissue with respiratory **bronchioles**, **alveolar** ducts, blood **vessels** and **terminal** bronchioles.



Bacterial infections of lower respiratory tract

- Bacterial pneumonia
- Pertussis
- Inhalation anthrax
- Q fever
- Tuberculosis

Types of Pneumonia



Community acquired

Typical vs Atypical pneumonia

Typical

- Pneumonia with **typical features**
- C/F: high fever, chest pain, cough which is **usually productive, shortness of breath, lobar consolidation on Chest Xray**
- **Bacteria:**
 - ☐ *Streptococcus pneumoniae*
 - ☐ *Hemophilus influenzae*
 - ☐ *Klebsiella pneumoniae*
 - ☐ *Moraxella catarrhalis*
 - ☐ *Staphylococcus aureus*

Atypical

- Pneumonia with **atypical features**
- C/F: Less severe illness, dry cough, headache, and other systemic complaints, patchy broncho-pneumonia
- **Bacteria:**
 - ☐ *Chlamydophila species*
 - ☐ *Mycoplasma pneumoniae*
 - ☐ *Legionella pneumoniae*
 - ☐ *Coxiella burnetii*
- **Viruses**



- **Community-acquired pneumonia (CAP)** is defined as an acute infection of the pulmonary parenchyma in a patient who has acquired the infection in the community.
- **Hospital-acquired (or nosocomial) pneumonia (HAP)** is pneumonia that occurs 48 hours or more after admission and did not appear to be incubating at the time of admission .
- **Ventilator-associated pneumonia (VAP)** is a type of HAP that develops more than 48 to 72 hours after endotracheal intubation .
 - Endotracheal intubation is a **medical procedure in which a tube is placed into the windpipe (trachea) through the mouth or nose**
 - EI is often an emergency procedure that's performed on people who are unconscious or who can't breathe on their own

- **Healthcare-associated pneumonia (HCAP)** is defined as pneumonia that occurs in a non-hospitalized patient with extensive healthcare contact, as defined by one or more of the following:
 - Intravenous therapy, wound care, or intravenous chemotherapy within the prior **30 days**
 - Residence in a nursing home or other long-term care facility
 - Hospitalization in an acute care hospital for two or more days within the prior **90 days**
 - Attendance at a hospital or hemodialysis clinic within the prior **30 days**

Aspiration Pneumonia

Inflammation of the lungs from inhalation of foreign material (food, liquids, vomit) or from fluids from the mouth.

☐ Predisposing conditions

- ☐ Any condition associated with altered or reduced consciousness (e.g. after a stroke, while sleeping)
- ☐ While vomiting
- ☐ Condition causing abnormal gag and swallowing reflexes (e.g. neurological dysfunction)

☐ Etiologic agents

☐ Normal flora of oropharynx

- ☐ Oral anaerobes: *Prevotella*, *Fusobacterium* spp., *Peptostreptococcus**
- ☐ Streptococci

* Anaerobes will
covered in GED

Pathogenesis of Bacterial Pneumonia

Invasion

- **Pathogen enters alveoli** from throat, nose, airborne droplets, blood
- Multiplies and invades alveolar epithelium
- Avoids phagocytosis (e.g. capsule or intracellular growth)

Damage

- **Tissue injury** via exotoxin production, host cell lysis, death of alveolar macrophages, etc.

Immune Response

- **Inflammation** → fever, chills, fatigue
- ↑ PMNs, serum components, alveolar macrophages

Spread

- Fluid accumulation and bacterial multiplication in alveoli
- **Spread of pathogens to nearby alveoli**
- Radiologic evidence of infection → consolidation

Healthy Airway



Normal airway:
Alveoli are open

Airway with Pneumonia



Mucus:
Traps cells,
reducing air space

Swelling:
Narrows the airway,
decreasing air flow

CCF
fluids

Pneumococcal Pneumonia

- Pneumonia caused by *Streptococcus pneumoniae*
- s the most common type of pneumonia, accounting for most hospitalizations for pneumonia




- **80% of all bacterial pneumonias due to SP**
- **Normal flora of URT**
 - **General Characteristics**
 - Gram (+) cocci occur singly or paired (diplococci)
 - Lanceolate shaped
 - Culture – blood agar or chocolate agar
 - **α – Hemolysis** (partial hemolysis displayed; **green** color results)
 - Facultative anaerobe – grows best at elevated CO₂

Streptococcus pneumoniae

Virulence Factors


- ☐ **Polysaccharide capsule** – essential for virulence
 - ☐ Antiphagocytic
 - ☐ 90 serotypes, 23 responsible for most infections
- ☐ **IgA protease**
 - ☐ Colonization
- ☐ **Autolysin**
 - ☐ Release of intracellular virulence factors and cell wall fragments
- ☐ **Pneumolysin** – pore-forming toxin
 - ☐ Released from autolysed *S. pneumoniae*
 - ☐ Damages respiratory epithelium
- ☐ **Peptidoglycan/teichoic acids**
 - ☐ Inflammatory



Damage of epithelium stimulates outpouring of fluid, red blood cells, and leukocytes from alveoli →
productive cough with blood

Progression of Pneumococcal Disease

Capsule prevents phagocytosis;
Toxins damage lung cells



The outcome of colonization depends on the virulence of the specific serotype and on the host immune system


a. Pneumonia

Develops when the bacteria multiplies in the alveolar space

- Sudden onset with fever, chills and sharp chest pain.
- Bloody or rusty sputum.
- Empyema is a rare but significant complication.
- Complications caused by spreading of pneumococci to other organs: sinusitis, middle ear infection, meningitis, endocarditis, septic arthritis, bacteremia.

Pneumococcal Pneumonia - Diagnosis

- ☐ Microscopy – Gram Stain
 - ☐ **GPDC, lancet-shaped**
- ☐ Culture – blood agar
 - ☐ α -hemolytic
 - ☐ Optochin (P disk) sensitivity
- ☐ Catalase negative
- ☐ Bile soluble
 - ☐ Susceptible to lysis
- ☐ Antimicrobial sensitivity test
- ☐ PCR



Optochin disc test



Res Sus
Bile solubility

Pneumococcal Infections Treatment

- Treatment:
 - Beta-lactams = penicillin or 3rd generation cephalosporin, clindamycin, vancomycin
- Resistance to penicillin is now common
altered penicillin-binding proteins - all *S. pneumoniae* isolates should be tested for susceptibility to beta-lactams

Pneumococcal Infections Prevention

- **PPSV**: pneumococcal polysaccharide vaccine = 23-valent polysaccharide vaccine
- **PCV**: pneumococcal conjugate vaccine = 13-valent conjugated vaccine for children < 60 months

Pulmonary Tuberculosis

- Commonly caused by *Mycobacterium Tuberculosis*

General characteristics

- Are acid fast bacteria
- Contain **mycolic acid** in their cell wall (Contains 60 to 90 carbons)
- Lipid rich cell wall, 60% of the cell wall is lipid (mycolic acids, cord factor, wax-D)
- Cell wall is responsible for:-

Acid fastness, Slow growth, resistance to detergents, resistance to many common antibiotics

A person may contract pulmonary tuberculosis from inhaling droplets from a cough or sneeze by an infected person

Granuloma in lung tissue




Pathogenesis

Transmission


- Inhalation of air borne droplet nuclei that contain *M.tb* bacilli (Particle of 1 - 5 μm in diameter)
 - One droplet nuclei contains not more than **3 bacilli**
 - Coughing generates about **3000** droplet nuclei
 - Talking for 5 minutes generates **3000** droplet nuclei
 - Singing generates **3000** droplet nuclei in one minute
 - Sneezing generates the most droplet nuclei by far, which can spread to individuals up to 10 feet away

- After being inhaled, mycobacteria reach the alveoli, where they multiply in the pulmonary epithelium or macrophages.
- Within two to four weeks, many bacilli are destroyed by the immune system, but some survive and are spread by the blood to extrapulmonary sites.
- The virulence of *M. tuberculosis* rests with its ability to survive and grow within host cells.
- The organism produces no demonstrable toxins; however, when engulfed by macrophages, bacterial sulfolipids inhibit the fusion of phagocytic vesicles with lysosomes.



(Richard A. Harvey, Pamella C. Champ, Microbiology, Lippincott's illustrated reviews, 2nd ed.)

Figure: Progression of active tuberculosis infection.



3

After a few weeks, many of the macrophages die, releasing tubercle bacilli and forming a caseous center in the tubercle, which is surrounded by a mass of macrophages and lymphocytes. The disease may become dormant after this stage.

4


In some individuals, a mature tubercle is formed, as a firm outer layer containing fibroblasts surrounds the mass of macrophages and lymphocytes. The caseous center enlarges by the process of liquefaction, forming a tuberculous cavity in which the bacilli multiply.

5

Liquefaction continues until the tubercle ruptures, allowing bacilli to spill into the bronchiole and be disseminated throughout the respiratory system and other systems of the body.

(Richard A. Harvey, Pamella C. Champ, Microbiology, Lippincott's illustrated reviews, 2nd ed.)

Figure: Progression of active tuberculosis infection.



(Richard A. Harvey, Pamela C. Champ, Microbiology, Lippincott's illustrated reviews, 2nd ed.)

Figure: Stages in the pathogenesis of tuberculosis.

Clinical Presentation

- Weight loss
- Productive cough (For more than a month)
- Chest pain
- High fever
- Night sweating
- Haemoptysis

Extrapulmonary TB

- During secondary TB, bacilli disseminate to regional lymph nodes, kidneys, long bones, genital tract, brain, and meninges.
- These complications are grave.
- Common immunocompromised individuals such as elderly, malnourished, or HIV-infected individuals.

Laboratory identification:

Specimens: Sputum, CSF, Biopsy or FNA

1. Ziehl Neelson staining / AFS

- Specific & rapid
- Cheap
- Low sensitivity (43 - 70% with repeated smear) ----- HIV??



Acid-fast stain (Ziehl-Neelsen stain): *Mycobacterium* + *Nocardia* spp.

- | | |
|-------------------------------------|----------------------|
| 1. Carbol-fuchsin with flame | Primary Stain |
| 2. 3.5% HCl + 90% alcohol | Decolorizer |
| 3. Methylene blue | Counter stain |


**AFB Positive
(Red Bacilli)
bacilli)**

**AFB Negative
(No red)**

ZN Stain



Acid fast
bacilli



Inflammatory
cells

Treatment

1. **Prolonged treatment:** slow response of TB treatment and should be continued for 6-12 mon (DOTs=6-8 mon)
 - a. Most bacilli are found intracellularly
 - b. Caseous material interfere with the drug
 - c. In chronic TB, bacilli are not dividing, i.e. "metabolically inactive", hence resistant to drugs.
2. **Combination of drugs:** to reduce toxicity due to prolonged course of treatment and rapid emergence of resistant strains.
 - ✓ Drug of choice: isoniazide (INH), rifampicin, ethambutol, streptomycin & para-amino-salicylic acid

Problems with treatment

- Resistance to anti-TB drugs can occur When the drugs are misused
 - Patients do not complete their full course of treatment (
 - Health-care providers prescribe the wrong treatment, the wrong dose, or length of time for taking the drugs
 - Supply of drugs is not always available
 - The drugs are of poor quality

Types of drug Resistance

- Mono drug resistance: resistance for one drug among the group
- Poly drug resistance: resistance to more than one drug with the exception of INH and Rifampin together
- Multidrug resistance (MDR): is resistant to at least two of the best anti-TB drugs, isoniazid and rifampicin

Types of drug Resistance cont'd



- Extensively Drug-Resistant (XDR): is resistant to isoniazid and rifampin
 - Plus resistant to any fluoroquinolone and at least one of three injectable second-line drugs
 - i.e., Amikacin, Kanamycin, or Capreomycin

Prevention & control

- Early case detection & treatment
- Decreasing of over crowding
- Pasteurization of milk --- ↓ *M. bovis* infection
- Health education
- Immunization
 - A vaccine against tuberculosis has been available since early in the twentieth century.
 - It is produced from Bacille Calmette-Gurin (BCG), an attenuated strain of *M. bovis*.
 - When injected intradermally, it can confer tuberculin hypersensitivity and an enhanced ability to activate macrophages that kill the pathogen.
 - is about eighty percent protective against serious forms of tuberculosis, such as meningitis in children, and has been used in mass immunization campaigns.

TUBERCULOSIS (TB)

- fatigue
- malaise
- anorexia
- wt. loss



chronic cough
(productive)

night sweats

hemoptysis
(advanced state)

low grade temp.
(late afternoon)


Treatment:

TB medications 6 to 12 months
bedrest until symptoms
resp isolation until
negative sputum
frequently out-pt basis

Diagnostic:

TB skin test
chest x-ray
bacteriologic
sputum studies

Pulmonary Anthrax



Bacillus anthracis:

- The agent of Anthrax
- An agent used in biological warfare

Bacillus anthracis


- ☐ Aerobic GPR
- ☐ Spore-forming
- ☐ Non-motile
- ☐ Polypeptide capsule
- ☐ Zoonotic – from herbivores



- Classification of anthrax infections
 - Dermal anthrax ~95% of cases
 - Respiratory anthrax (wool-sorter's disease) – inhalation of spores
 - ☐ Bioterrorism threat
 - Gastrointestinal anthrax
 - ☐ Rare in humans, common in animals
 - Sepsis - from primary infection


Bacillus anthracis

- ☐ Ubiquitous / zoonotic, lives in soil
 - ☐ Causes anthrax in animals and humans
 - Large **Gram positive bacilli**
 - Arranged in **long serpentine chains** with a '**jointed bamboo stick**' appearance or '**Box car shaped**' morphology.
 - Forms **endospores** (not seen in clinical samples)
 - Non-motile
 - **Has a polypeptide capsule made up of poly-d glutamic acid.**
 - ☐ Culture- Blood agar: non hemolytic, large, irregular colonies with raised edge, '**MEDUSA HEAD**' appearance.
 - ☐ Exotoxins- Anthrax toxin
 - ☐ Tripartite toxin
- } Protective antigen
Edema factor
Lethal factor



Epidemiology

- Cutaneous anthrax is common in leather industry workers and butchers.
- Pulmonary anthrax- Bioterrorism- Classified as a 'Category A' biological agent



Exposure to spores present in contaminated soil or in infected animal products (hide, hair and wool)

AKA Hide Porter Disease/ Malignant pustule

Inhalation of dust, filaments of wool
AKA Wool Sorter's Disease

Consumption of infected meat

All types may lead to **septicaemia and meningitis**


Respiratory (Inhalation) Anthrax Symptoms

- Variable incubation (average 5 days)
- Prodromal phase
 - Nonspecific, flu-like syndrome
- Fulminant stage
 - Rapidly worsening course of fever, edema, enlargement of mediastinal lymph nodes
 - Respiratory distress, cyanosis, shock
 - Almost uniformly fatal within days

Pulmonary Anthrax – Pathogenesis



- **Inhalation of anthrax spores (infectious form)** → pulmonary alveoli
- Spores are **phagocytized by alveolar macrophages** where they germinate and **replicate**
- **Transported** through pulmonary lymphatics to **mediastinal lymph nodes**



B. anthracis - spore and vegetative stage



B. anthracis spores in lung bronchiole



■ Endospore

- Protective, dormant form of a bacterial cell; resistant to environment
- Major component is calcium dipicolinate
- Stain with Malachite green
- *B. anthracis* spores can live in the soil for many years

Anthrax - Diagnosis


□ Microscopy

- Gram stain = **GPR, boxcars**
- **Capsule**: stained by M'Fadyean methylene blue stain or DFA
- **Spores**: seen in malachite green stain

□ Culture on blood agar plate

- Large, **non-hemolytic adherent colonies**

□ Molecular assays



Anthrax –

Treatment and Prevention



- ☐ Treatment

- ☐ Ciprofloxacin

- ☐ Prevention

- ☐ Toxoid, cell-free **vaccine** used for high-risk persons

- ☐ Vaccination of animals to control disease

- ☐ Eradication unlikely because of spore production

Coxiella burnetii

POTENTIAL BIOTERRORISM AGENT

☐ Obligate intracellular GNR

☐ Stain poorly

- ☐ Forms spore-like stage for environmental survival

☐ Reservoirs

☐ Mammals (especially farm animals)

- ☐ High concentration in placenta


☐ Can survive in soil and milk for years

☐ Transmission

☐ Inhalation of airborne particles containing spores

☐ Zoonotic

- ☐ Risk factor = occupational exposure to animals



Coxiella burnetii – Pathology, Diagnosis and Treatment

□ Q fever

- Asymptomatic
- Self-limiting flu-like syndrome
- Atypical pneumonia
 - Non-productive cough
- Hepatitis

□ Pathogenesis

- Intracellular survival

□ Diagnosis

- Serology

□ Treatment

- Doxycycline


B. pertussis


- Causes **Pertussis or whooping cough** which is a serious, highly communicable acute tracheobronchitis
- **Pertussis** is a highly contagious disease that occurs primarily in **infants and young children** and has a worldwide distribution
 - Colonize **respiratory epithelium**
 - Spread by aerosol/direct contact
 - The bacteria is **Gram negative aerobic coccobacillus**
 - Oxidize **only aas** not CHO
 - **Nutritionally fastidious, normally cultivated on medium containing blood**

Some virulence factors of *B. pertussis*

- **Non-invasive bacterium-does not normally invade cells**
- **Attachment**
 - Filamentous hemagglutinin (FHA) and pertactin
 - Pertussis toxin
- **Toxins**
 - Tracheal cytotoxin – cell death
 - Endotoxin – cytokines, fever
 - Adenylate cyclase/hemolysin – inhibits lymphocyte function
 - Lipooligosaccharide (endotoxin)
 - **Pertussis toxin (PT)* - inhibits immune cells chemotaxis**
and acts as an adhesin

Pathogenesis of *Bordetella pertussis*





Pathophysiology of *Bordetella pertussis* causing whooping cough

Clinical features (2or 3 Stages)

a. **Catarrhal stage**

seen after 7-10 days

resemble common cold(rhinorrhea, sneezing, malaise, anorexia, low grade fever)

b. **Paroxysmal stage**

Ciliated cells are removed from the RT

mucus clearance is impaired

Whooping cough paroxysmal (repetitive cough, exhaustion, vomiting, cyanosis and convulsion)

c. **Convalescent stage**

- paroxysm diminishes in no. and severity
- 2^o complication may develop (pneumonia, encephalopathy)

Pertussis

Bordetella pertussis

Stage	Length	Clinical Features
Stage 1: Catarrhal	Usually 7-10 days; range = 4-21 days	Runny nose, fever, occasional cough
Stage 2: Paroxysmal	1-6 weeks; maybe up to 10 weeks	Paroxysms of rapid cough (thick mucus in resp tree) Whoop at end of cough 15 attacks/24 hours
Stage 3: Convalescent	7-10 days; range = 4- 21	Cough severity lessens

PERTUSSIS (WHOOPING COUGH)



Whooping cough: whooping sound made when gasping for air after a fit of coughing

Whooping cough

Diagnosis:

- Isolation of bacteria: positive during the 1st week only
- Culture from mucus or droplets: **Mercury drops colony**
- Serology: using specific anti-sera or specific antigens

Prophylaxis:


- Heat killed vaccine to children during 1st year of life in combination with diphtheria and tetanus (DPT)

Treatment:

- Antibiotics: erythromycin effective if given early
- Recovery is followed by long lasting immunity



Bordetella pertussis growing in Regan-Lowe Agar Incubated in CO₂ for 72 hours



"mercury drop" colonies on Bordet' Gengou

L. pneumophila

- Is slender, pleomorphic, gram-negative rods
- obligatively aerobic and nutritionally fastidious..
- 90% of all infections is by serotypes 1 (of the 12 serotypes)
- Commonly cause Respiratory tract disease
- Transmission is through inhalation of infectious aerosols.
- Legionellae are facultative intracellular bacteria, multiply in free-living amoebae in nature, and in alveolar macrophages, monocytes, and alveolar epithelial cells in infected hosts

- The bacteria are commonly present in **natural bodies of water**, such as lakes and streams, as well as in **air conditioning cooling towers** and condensers and in water systems (e.g., shower heads).
- Human infections are most commonly associated with **exposure to contaminated aerosols**
- **The organism survive in moist environment**, for a long time
- bacteria parasitize **amoebae** in the water and replicate
- bacteria also survive in **biofilms** that develop in the **pipes** of water systems

Clinical Diseases

(1) an influenza-like illness (**Pontiac fever**)

- Is a **self-limited**, febrile
- The name "**Pontiac**" is derived from the city in USA where the first outbreak appeared

2) a severe form of pneumonia (**Legionnaires'disease**).

Pontiac Fever

❑ Mild form of *Legionella* infection


❑ Flu-like symptoms

❑ Usually self-limiting

Not this →




This one ↓



Legionnaires' disease


Infection

Caused by bacteria which thrives in warm water and damp places like:



Source: WHO, UK NHS

Symptoms



Treatment

Antibiotics



No vaccine is currently available




Prevention



Water supply systems should be cooled below 20°C or heated above 60°C

Legionnaires' Disease Pathogenesis

- 1) Adheres to respiratory epithelium
- 2) Attaches to alveolar macrophages → **coiling phagocytosis**
- 3) Prevents phagosome-lysosome fusion
- 4) **Replication inside the macrophage**
- 5) Lysis of phagocytes
Hydrolytic enzymes → Lung damage and inflammatory response




Reproduced from: Swanson MS, Hammer BK, 2000. *Legionella pneumoniae* pathogenesis: a fateful journey from amoebae to macrophages. *Annu Rev Microbiol* 54:567-613.

Legionella is a **facultative intracellular pathogen**
Major role in **immunity—cell mediated**; Minor role: Humoral immunity

Legionnaires' Disease – Diagnosis

□ Microscopy

- Poorly staining GNR
- **Direct fluorescent antibody (DFA)**
- **Silver stain**



Fluorescein-labeled antibody attached to *Legionella* bacilli

□ Culture

- Fastidious
 - **Requires cysteine and iron**
 - **Buffered charcoal yeast extract (BCYE)** (*Legionella* rides its BiCYcLE)



BCYE

□ Urinary antigen test

- Serogroup 1 only



Legionella urinary antigen card



Legionnaires' Disease

Treatment and Prevention



- Treatment
 - Azithromycin
 - Levofloxacin
- Prevention
 - Disinfection of water systems

Klebsiella pneumoniae

- *K. pneumoniae* is the most frequent cause of gram negative bacterial pneumonia.
- *Klebsiella*-related pneumonia frequently afflicts debilitated and malnourished persons, particularly *chronic alcoholics*.
- Thick and gelatinous sputum is characteristic, because the organism produces an abundant viscid capsular polysaccharide, which the patient may have difficulty coughing up.

Pseudomonas aeruginosa

- *Pseudomonas* pneumonia is common in persons who are neutropenic, usually secondary to chemotherapy; in victims of extensive burns; and in patients requiring mechanical ventilation.
- *P. aeruginosa* has a propensity to invade blood vessels at the site of infection, with consequent extra-pulmonary spread; *Pseudomonas* bacteremia is a fulminant disease, with death often occurring within a matter of days.

Orthomyxoviruses (Influenza virus)


- Orthomyxoviruses are spherical, **enveloped** viruses containing a **segmented**, **negative** strand RNA genome.
- Viruses in this family infect **humans, horses, and pigs, as well as nondomestic animals**
- Orthomyxoviruses have one genus and three types

Genus:- **Influenzae**

types:- **influenzae A, B, and C.**

-Only influenza virus types **A** and **B** are of medical importance.

- Type A influenza viruses differ from type B viruses in that they have an animal reservoir (humans, animals, birds) and are
- Type A divided into subtypes.
- B infects only human beings
- .



1. Schematic diagram of structure of an influenza virus.

- The envelope is covered with two different types of **spikes**

1. **Hemagglutinin (HA)** – **15** different subtypes; most important

virulence factor; Assist virus to bind to host cells receptors

(**neuraminic acid/ sialic acid**)

- hemagglutinin agglutinates red blood cells, which is the basis of a **diagnostic test**
- important in determining the **subtype** of influenza virus isolates (**classification**)

2. Neuraminidase (N) – 9 types

- The neuraminidase cleaves **neuraminic acid (sialic acid)** to release progeny virus from the infected cell (budding)
- **degrades** the protective **layer of mucus** in the respiratory tract (virulence role)
- important in determining the **subtype** of influenza virus isolates
- HA and NA are used to sub type the viruses

Classification


- HA and NA, are used to **subtype** the viruses
- Only type **A** possesses subtypes

Nomenclature of subtypes of influenza A virus

Strains of influenza **A virus** are classified to subtypes by the following four characteristics:

1. Type (A, B, and C)
2. Place of original isolation
4. strain number
3. Date of original isolation
4. Antigen (HA and NA)

Influenza virus nomenclature



E.g.,

- **A/Bangkok/15/79 (H3N2)**, meaning that it is an influenza A virus that was first isolated in Bangkok in 1979 and contains HA (H3) and NA (N2) antigens.
- Antigenic subtypes of HA and NA are given in **(brackets)**
- When it is isolated from animals the host is mentioned


E.g., **A/swine/Bangkok/15/79 (H3N2)**

- 15 subtypes of HA (H1–H15) and nine subtypes of NA (N1–N9), in many different combinations, have been recovered from birds, animals, or humans


Antigenic shift

Major antigenic changes (antigenic shift) result from the **reassortment** (genome exchange of two strain) of genomes among different strains, including animal strains.

- When this occurs, there will be **worldwide epidemics**
/pandemic may be the consequence since the entire population is susceptible to the virus (**occurs only in influenzae A**)
- There is formation/appearance of **new strain**



REASSORTMENT OF INFLUENZA VIRUS



Antigenic Drift

- This term denotes MINOR changes in hemagglutinin and neuraminidase of influenza virus.
- This involves no change in serotype; there is merely an alteration in amino acid sequence of HA or NA leading to change in antigenicity.
- **No formation of new strain**

Influenza Virus: Antigenic Changes


Antigenic drift

- minor, continual changes, same subtype
- caused by point mutations in gene
- may result in epidemic

Antigenic shift

- major, abrupt changes, new subtype
- caused by exchange of gene segments
- may result in pandemic

Antigenic Drift and Shift of Influenza Strains



Antigenic drift vs. shift. Antigenic drift creates influenza viruses with slightly modified antigens, while antigenic shift generates viruses with entirely new antigens (shown in red). Source: Wikipedia Commons and USDA.

Pathogenesis


The virus initially cause upper respiratory tract infection

- It target and kill mucus secreting ciliated and cause the loss of primary defense system.
- In LRT the virus causes sever desquamation (shedding) of bronchial or alveolar epithelium
- It also promotes bacterial adhesion to the epithelial cells (secondary bacterial infection)

The most susceptible population are

- Children, immuno-suppressed people (including pregnant women), elderly people, people with heart and lung ailments(including smokers).

Symptoms of **Influenza**



Complications

- bacterial pneumonia, myositis, and Reye syndrome (an acute encephalitis that affects children occurs after a variety of acute febrile viral infections)
- Secondary bacterial infection usually involves *Streptococcus pneumoniae*, *Haemophilus influenzae*, or *Staphylococcus aureus*.

Mode of transmission

- Virus is spread by inhalation of small aerosol droplets expelled during talking, breathing, and coughing.
Virus likes cool, less humid atmosphere

	TYPE A	TYPE B	TYPE C
severity of illness	++++	++	+
animal reservoir	yes	no	no
human pandemics	yes	no	no
human epidemics	yes	yes	no (sporadic)
antigenic changes	shift, drift	drift	drift
segmented genome	Yes (8)	Yes (8)	Yes(7)
amantadine, rimantidine	sensitive	no effect	no effect
zanamivir	sensitive	sensitive	

Lab diagnosis:


Diagnosis:

- Cell or tissue culture
- Hemadsorption
- Viral antigen detection

Treatment

- Two basic approaches
 - Symptomatic care & anticipation of potential complications
- Best treatments include:
 - Rest and fluid intake, cough suppressants
 - Conservative use of analgesics for myalgia and headache
- Amantidine & rimantadine (diagnosed within 12-24 hrs) (M2)
- Zanamivir and tamiflu (oseltamivir) target neuraminidase

Prevention



- Get the flu vaccine each year due to high mutation rate of the virus.
- Practice good hygiene and personal health habits.
- Cover your mouth when while sneezing and wash your hands regularly as the virus spreads through aerosols.
- Since the flu is a virus, antibiotics won't work unless there is a secondary bacterial infection.

Hantavirus Pulmonary Syndrome

Cause: *Hantavirus*
(enveloped, segmented,
–ssRNA virus).

Portal of entry:
Inhalation of infected
rodent excrement or
saliva.

Signs and symptoms:
Sudden onset of fever,
fatigue, muscle aches,
headache, chills,
abdominal pain;
followed by coughing
and severe difficulty in
breathing.



▲ The deer mouse, *Peromyscus maniculatus*.

Incubation period: 14–30 days.

Susceptibility: Those exposed to rodents in endemic areas.

Treatment: Supportive therapy for symptoms must occur under hospital supervision.

Prevention: Avoid rodent droppings, camp on ground sheet, carefully clean out potential rodent nesting sites at home, and keep home clean.

Respiratory Syncytial Virus Infection

- is the most common childhood respiratory disease in newborns and young Children

RSV is the leading cause of bronchiolitis (inflammation of the bronchioles) and pneumonia among children less than one year of age and the leading respiratory killer of infants worldwide

As its name indicates, the virus causes **syncytia** to form in the lungs.

- Plugs of mucus, fibrin, and dead cells in the bronchioles make it difficult to breathe. The action of cytotoxic T cells and other specific immune responses to RSV infection further damages the lungs

Treatment—Aerosolized ribavirin for very sick infants.

Prevention—Passive immunization with palivizumab (monoclonal antibody) or immune globulins in infants who have been exposed is effective. Handwashing and the use of gloves may prevent nosocomial outbreaks in the newborn nursery.

Fungal infections of the Lower respiratory tract

Fungal infections of lower respiratory tract



- Fungal spores are easily inhaled and may germinate in LRT
- Two major factors govern incidence and spread of fungal infection
 - Ubiquity of the infectious organisms
 - Found in soil
 - Resident flora
 - Adaptive immune response
 - Usually keeps these infections under control
 - Immunocompromised patients at much greater risk
- Incidence is being increased in recent years. Why?

1. *Pneumocystis pneumonia* (pcp)

- A lethal pneumonia
 - Common in AIDS patients
- Caused by the fungus *Pneumocystis (carinii) jiroveci*
 - Never been grown in culture
 - Most information comes from clinical information of patients


Pneumocystis Pneumonia

- *Pneumocystis jirovecii* (formerly *P. carinii*) is sometimes found in healthy human lung
- Pneumonia; a common serious complication of AIDS



2. Blastomycosis

- Caused by *Blastomyces dermatitidis*
- Spores of the fungi enter through the respiratory system
- Primarily affect the lungs
 - Can spread through bloodstream and affect other parts
- Men between ages of 20 and 40 years are the most commonly infected



Pathogenesis...Blastomycosis


- Infection of the lungs is gradual
 - Fever, chills, and drenching sweats develop
 - Chest pain, difficulty breathing, and cough may also develop
 - Can sometimes heal without treatment
 - Skin: warty patches develop surrounded by tiny painless abscesses
 - Bones: painful swellings
 - GUT: prostatitis or painful swelling of epididymis

3. Histoplasmosis

- Caused by *Histoplasma capsulatum*
 - Occurs in soil contaminated with bat or bird droppings
 - Commonly found in temperate, subtropical and tropical zones
 - 50% - 90% of residents in these areas test positive for exposure
 - People who live and work in the vicinity of bat or bird droppings are at increased risk of infection
 - Amphotericin B is the treatment of choice if necessary

...Histoplasmosis

- *Histoplasma capsulatum*: subclinical respiratory infection that only occasionally progresses to severe, generalized disease
- Acquired by inhalation of airborne conidia
- Isolation of the fungus or identification in tissues sample is necessary for diagnosis
- Treatment : Amphotericin B



Pathogenesis...Histoplasmosis



- **Transmission** is through inhalation of conidia
 - Small enough to reach bronchioles and alveoli
- After inhalation:
 - Microconidia convert to yeast form, phagocytosis, tubercle formation
 - Severe cases may develop chills, malaise, chest pain, and extensive pulmonary infiltration

5. Coccidioidomycosis

- Caused by *Coccidioides immitis*
- Infection can be symptomatic or asymptomatic
 - Symptomatic form known as Valley Fever
- Arthroconidia of the fungus are inhaled.
 - Small enough to bypass defenses of the upper tract.
 - Lodge directly in bronchioles.
- Fungal outer wall has antiphagocytic properties
 - Prevents elimination

...*Coccidioidomycosis*

- Inhalation of the airborne arthroconidia of *Coccidioides immitis*
- Most cases are subclinical, but when there are predisposing factors such as fatigue and poor nutrition, a progressive disease resembling tuberculosis can result → fever, coughing, weight loss; occasionally fatal



Pathogenesis...coccidioidomycosis

- **Arthroconidia** convert to spherules which grow slowly
 - Completely inhibit phagocytosis
- Disseminated coccidioidomycosis: AIDS patients and on immunosuppressive therapy
- Can also cause a form of coccidioidal **meningitis**
 - Can be fatal if not treated aggressively
- Usually self-limiting and no treatment is required
- Progressive pulmonary infection or infection of central nervous system is treated with amphotericin B

6. Aspergillosis

- Invasive aspergillosis shows a rapid progression to death
- Typically seen in the immunocompromised
 - Particularly patients with leukemia or AIDS
 - Patients undergoing bone marrow transplantation
- Also seen in individuals with preexisting pulmonary disease
 - Chronic bronchitis, asthma, and tuberculosis
 - Fungus produces extracellular proteases, phospholipases, and toxic metabolites

...*Aspergillosis*

- Caused by the fungus *Aspergillus*
 - Widely distributed and found throughout world
 - Dispersal is through inhalation of resistant conidia.
 - Seen more and more in nosocomial infections associated with air-conditioning systems.

...Aspergillosis

Many other opportunistic fungi may cause respiratory d/se

Aspergillus fumigatus can cause:

- Allergic bronchopulmonary aspergillosis
- Aspergilloma: in patients with pre-existing lung disease →
mass of hyphae produce fungus ball
- Disseminated aspergillosis

...Aspergillosis

