

MTH251 & MTH252

Mathematical Analysis I & II

Course Textbook

Department of Mathematical Sciences
UNIST

Mathematical Analysis I & II

Course Textbook

Author

Muho Choi (Department of Computer Science and Engineering)

Email: sihyeon8240@unist.ac.kr

GitHub: <https://github.com/sihyeon8240>

Source Code & Contributions

GitHub: <https://github.com/sihyeon8240/Math>

Feel free to open issues or pull requests for corrections.

© 2026. Muho Choi. All rights reserved.

Developed for the Department of Mathematical Sciences, UNIST.

This textbook is licensed under a **Creative Commons Attribution-NonCommercial-ShareAlike 4.0 International License** (CC BY-NC-SA 4.0).

You are free to: Share, copy, and redistribute the material in any medium or format; adapt, remix, transform, and build upon the material.

Under the following terms: You must give appropriate credit; you may not use the material for commercial purposes; if you remix or transform the material, you must distribute your contributions under the same license.

Contents

1	The Number Systems	1
1.1	The Natural, Integer, and Rational Number Systems	1
1.2	Ordered Sets and Ordered Fields	2
1.3	The Real Field	6
1.4	The Extended Real Number System	14
1.5	The Complex Field	15
1.6	Euclidean Spaces	18
2	Basic Topology	20
2.1	Relations	20
2.2	Functions	21
2.3	Finite Sets	24
2.4	Countable and Uncountable Sets	32
2.5	Metric Spaces	40
2.6	Compact Sets	42
2.7	Perfect Sets	42
2.8	Connected Sets	42
3	Numerical Sequences and Series	43
3.1	Convergent Sequences	43
3.2	Subsequences	43
3.3	Cauchy Sequences	43
3.4	Upper and Lower Limits	43
3.5	Some Special Sequences	43
3.6	Series	43
3.7	Series of Nonnegative Terms	43
3.8	The Number e	43
3.9	The Root and Ratio Tests	43
3.10	Power Series	43
3.11	Summation by Parts	44
3.12	Absolute Convergence	44
3.13	Addition and Multiplication of Series	44
3.14	Rearrangements	44

4	Continuity	45
4.1	Limits of Functions	45
4.2	Continuous Functions	45
4.3	Continuity and Compactness	45
4.4	Continuity and Connectedness	45
4.5	Discontinuities	45
4.6	Monotonic Functions	45
4.7	Infinite Limits and Limits at Infinity	45
5	Differentiation	46
5.1	The Derivative of a Real Function	46
5.2	Mean Value Theorems	46
5.3	The Continuity of Derivatives	46
5.4	L'Hospital's Rule	46
5.5	Derivatives of Higher Order	46
5.6	Taylor's Theorem	46
5.7	Differentiation of Vector-valued Functions	46
6	The Riemann-Stieltjes Integral	47
6.1	Definition and Existence of the Integral	47
6.2	Properties of the Integral	47
6.3	Integration and Differentiation	47
6.4	Integration of Vector-valued Functions	47
6.5	Rectifiable Curves	47
7	Sequences and Series of Functions	48
7.1	Discussion of Main Problem	48
7.2	Uniform Convergence	48
7.3	Uniform Convergence and Continuity	48
7.4	Uniform Convergence and Integration	48
7.5	Uniform Convergence and Differentiation	48
7.6	Equicontinuous Families of Functions	48
7.7	Equicontinuous Families of Functions	48
7.8	The Stone-Weierstrass Theorem	48
8	Some Special Functions	49
8.1	Power Series	49
8.2	The Exponential and Logarithmic Functions	49
8.3	The Trigonometric Functions	49
8.4	The Algebraic Completeness of the Complex Field	49
8.5	Fourier Series	49
8.6	The Gamma Function	49

9	Functions of Several Variables	50
9.1	Linear Transformations	50
9.2	Differentiation	50
9.3	The Contraction Principle	50
9.4	The Inverse Function Theorem	50
9.5	The Implicit Function Theorem	50
9.6	The Rank Theorem	50
9.7	Determinants	50
9.8	Derivatives of Higher Order	50
9.9	Differentiation of Integrals	50
10	Integration of Differential Forms	51
10.1	Integration	51
10.2	Primitive Mappings	51
10.3	Partitions of Unity	51
10.4	Change of Variables	51
10.5	Differential Forms	51
10.6	Simplexes and Chains	51
10.7	Stokes' Theorem	51
10.8	Closed Forms and Exact Forms	51
10.9	Vector Analysis	51
11	The Lebesgue Theory	52
11.1	Set Functions	52
11.2	Construction of the Lebesgue Measure	52
11.3	Measure Spaces	52
11.4	Measurable Functions	52
11.5	Simple Functions	52
11.6	Integration	52
11.7	Comparison with the Riemann Integral	52
11.8	Integration of Complex Functions	52
11.9	Functions of Class $\mathcal{L}^2$	52

Chapter 1

The Number Systems

1.1 The Natural, Integer, and Rational Number Systems

We denote the set $\{1, 2, 3, \dots\}$ of natural numbers by $\mathbb{N}$. The set $\mathbb{N}$ is defined by the following axioms:

- $1 \in \mathbb{N}$.
- If $n \in \mathbb{N}$, then $s(n) \in \mathbb{N}$ where $s(n) = n + 1$ is called the **successor** of n .
- 1 is not the successor of any element of $\mathbb{N}$.
- If $m, n \in \mathbb{N}$ and $s(m) = s(n)$, then $m = n$.
- Any subset of $\mathbb{N}$ which contains 1, and contains $s(n)$ whenever it contains n , must be equal to $\mathbb{N}$.

The axioms above are called the **Peano axioms** (or *Peano Postulates*).

Remark. Let $P(n)$ be a statement about $n \in \mathbb{N}$, and let $S := \{n \in \mathbb{N} \mid P(n) \text{ is true}\}$. Suppose $P(n)$ satisfies the following conditions:

- $P(1)$ is true.
- $P(n + 1)$ is true whenever $P(n)$ is true.

Then, we have the following:

- $1 \in S$.
- $n + 1 \in S$ whenever $n \in S$.

Thus, we have $S = \mathbb{N}$, i.e., $P(n)$ is true for all $n \in \mathbb{N}$. This method of proof is called the **principle of mathematical induction**.

Theorem 1.1.1. *Every nonempty subset of $\mathbb{N}$ has a least element. This theorem is called the **well-ordering property** of $\mathbb{N}$.*

Proof. Let S be a nonempty subset of $\mathbb{N}$. Suppose S does not have a least element. Let T and L be defined as follows:

- $T := \mathbb{N} \setminus S$.
- $L := \{n \in \mathbb{N} \mid \{1, 2, \dots, n\} \subseteq T\}$.

We will show that $L = \mathbb{N}$, which implies that $S = \emptyset$, a contradiction.

- **Base case:** $1 \in L$ since $\{1\} \subseteq T$. Otherwise, $1 \in S$ and 1 is the least element of S .
- **Inductive step:** Suppose $n \in L$, i.e., $\{1, 2, \dots, n\} \subseteq T$. Then $n + 1 \notin S$. Otherwise, $n + 1$ is the least element of S . Therefore, we have $n + 1 \in T$ and $\{1, 2, \dots, n + 1\} \subseteq T$. Thus, we have $n + 1 \in L$.

By the principle of mathematical induction, we have $L = \mathbb{N}$. □

We define the set $\mathbb{Z}$ of integers and the set $\mathbb{Q}$ of rational numbers as follows:

- $\mathbb{Z} := \{m - n \mid m, n \in \mathbb{N}\}$.
- $\mathbb{Q} := \left\{ \frac{m}{n} \mid m \in \mathbb{Z}, n \in \mathbb{N} \right\}$.

Also, we denote the set of positive numbers or negative numbers as follows:

- $\mathbb{Z}^+ := \{n \mid n \in \mathbb{N}\}, \mathbb{Z}^- := \{-n \mid n \in \mathbb{N}\}$.
- $\mathbb{Q}^+ := \left\{ \frac{m}{n} \mid m \in \mathbb{N}, n \in \mathbb{N} \right\}, \mathbb{Q}^- := \left\{ \frac{m}{n} \mid m \in \mathbb{Z}^-, n \in \mathbb{N} \right\}$.

1.2 Ordered Sets and Ordered Fields

A set S is called an **ordered set** if it is equipped with a relation $<$ such that for any $x, y, z \in S$, we have the following properties:

- (A) $x < y$ and $y < z$ implies $x < z$.
- (B) Only one of the relations $x < y$, $x = y$, or $x > y$ holds.

$x > y$ means $y < x$, $x \leq y$ means $(x < y \text{ or } x = y)$ and $x \geq y$ means $y \leq x$.

Remark. In an ordered set S , the relation $<$ is called an **order** (or a **total order**) on S . If a relation $<$ on T satisfies (A) only, then the relation $<$ is called a **partial order** on T and the set T is called **partially ordered**.

A set F is called a **field** if it is equipped with two operations, *addition* and *multiplication*, and satisfies the following axioms:

(A) Axioms for addition

- $x + y \in F$ for all $x, y \in F$.
- $x + y = y + x$ for all $x, y \in F$.
- $(x + y) + z = x + (y + z)$ for all $x, y, z \in F$.
- There exists an element $0 \in F$ such that $x + 0 = x$ for all $x \in F$.
- For each $x \in F$, there exists an element $-x \in F$ such that $x + (-x) = 0$.

(B) Axioms for multiplication

- $x \cdot y \in F$ for all $x, y \in F$.
- $x \cdot y = y \cdot x$ for all $x, y \in F$.
- $(x \cdot y) \cdot z = x \cdot (y \cdot z)$ for all $x, y, z \in F$.
- There exists an element $1 \in F$ with $1 \neq 0$ such that $x \cdot 1 = x$ for all $x \in F$.
- For each $x \in F$ with $x \neq 0$, there exists an element $x^{-1} \in F$ such that $x \cdot x^{-1} = 1$.

(C) The distributive law

- $x \cdot (y + z) = x \cdot y + x \cdot z$ for all $x, y, z \in F$.

A set F is called an **ordered field** if it is both an ordered set and a field, and satisfies the following axioms:

- If $x, y, z \in F$ and $x < y$, then $x + z < y + z$.
- If $x, y \in F$ and $x > 0$ and $y > 0$, then $x \cdot y > 0$.

Definition 1.2.1. Let S be an ordered set and $E \subseteq S$.

- E is **bounded above** if there exists some $M \in S$ such that $x \leq M$ for all $x \in E$. Such an M is called an **upper bound** of E .
- E is **bounded below** if there exists some $m \in S$ such that $m \leq x$ for all $x \in E$. Such an m is called a **lower bound** of E .

Theorem 1.2.2. Any nonempty subset of $\mathbb{Z}$ which is bounded below in $\mathbb{Z}$ has a unique least element, and any nonempty subset of $\mathbb{Z}$ which is bounded above in $\mathbb{Z}$ has a unique greatest element.

Proof. Let E be a nonempty subset of $\mathbb{Z}$ which is bounded below in $\mathbb{Z}$. Then there exists a lower bound $m \in \mathbb{Z}$ of E . Let F be defined as follows:

$$F := \{x - m + 1 \mid x \in E\} \subseteq \mathbb{N}.$$

By [Theorem 1.1.1](#), F has a least element n . Then we have

$$n \leq x - m + 1 \text{ for all } x \in E \implies n + m - 1 \leq x \text{ for all } x \in E.$$

Since $n \in F$, there exists a unique $x_0 \in E$ such that $n = x_0 - m + 1$, so $x_0 = n + m - 1$ is the unique least element of E .

Let S be a nonempty subset of $\mathbb{Z}$ which is bounded above in $\mathbb{Z}$. Then there exists an upper bound $M \in \mathbb{Z}$ of S . Let T be defined as follows:

$$T := \{-x + M + 1 \mid x \in S\} \subseteq \mathbb{N}.$$

By [Theorem 1.1.1](#), T has a least element n . Then we have

$$n \leq -x + M + 1 \text{ for all } x \in S \implies M - n + 1 \geq x \text{ for all } x \in S.$$

Since $n \in T$, there exists a unique $x_0 \in S$ such that $n = -x_0 + M + 1$, so $x_0 = M - n + 1$ is the unique greatest element of S . $\square$

Definition 1.2.3. Let S be an ordered set. Suppose $E \subseteq S$ is bounded above. If there exists an $\alpha \in S$ such that

- α is an upper bound of E .
- If $\beta < \alpha$, then β is not an upper bound of E .

then α is called the **supremum** (or *least upper bound*) of E , denoted by

$$\alpha = \sup E.$$

Similarly, suppose $F \subseteq S$ is bounded below. If there exists an $\gamma \in F$ such that

- γ is a lower bound of F .
- If $\beta > \gamma$, then β is not a lower bound of F .

then γ is called the **infimum** (or *greatest lower bound*) of F , denoted by

$$\gamma = \inf F.$$

Example 1.2.4. Let $S \subseteq \mathbb{Q}^+$ be defined as

$$S := \{p \in \mathbb{Q}^+ \mid p^2 < 2\}.$$

For any $p \in S$, let

$$q := p - \frac{p^2 - 2}{p + 2} = \frac{2p + 2}{p + 2} > p.$$

Then we have $q \in \mathbb{Q}^+$ and

$$q^2 - 2 = \frac{2(p^2 - 2)}{(p + 2)^2} < 0 \implies q \in S.$$

This implies that any $p \in S$ is not an upper bound of S .

For any $p \in \mathbb{Q}^+$ with $p^2 > 2$, p is an upper bound of S . Let

$$q := p - \frac{p^2 - 2}{p + 2} = \frac{2p + 2}{p + 2} < p.$$

Then we have $q \in \mathbb{Q}^+$ and

$$q^2 - 2 = \frac{2(p^2 - 2)}{(p + 2)^2} > 0 \implies q \text{ is a lower bound of } S.$$

This implies that such p is not the least upper bound of S . Since there exists no $p \in \mathbb{Q}^+$ such that $p^2 = 2$, S has no least upper bound in $\mathbb{Q}^+$.

Definition 1.2.5. An ordered set S is said to have the **least upper bound property** if any nonempty subset of S which is bounded above has a least upper bound in S .

Theorem 1.2.6. Let S be an ordered set with the least upper bound property. Then, any nonempty subset of S which is bounded below has a greatest lower bound in S .

Proof. Let E be a nonempty subset of S which is bounded below. Let L be defined as follows:

$$L := \{\lambda \mid \lambda \text{ is a lower bound of } E\}.$$

Then, L is nonempty and bounded above. By the least upper bound property, there exists $\alpha = \sup L \in S$. We will show that $\alpha = \inf E$.

Suppose $\alpha \notin L$. Then we have the following:

- α is not a lower bound of $E \implies$ There exists $\beta \in E$ such that $\beta < \alpha$.
- $\alpha = \sup L \implies \beta$ is not an upper bound of $L \implies$ There exists $\gamma \in L$ such that $\beta < \gamma \leq \alpha$.
- $\gamma \in L \implies \gamma$ is a lower bound of $E \implies \gamma \leq \beta$.

This is a contradiction. Therefore, $\alpha \in L$, i.e., α is a lower bound of E . Since α is an upper bound of L , for any $\delta > \alpha$, we have $\delta \notin L \implies \delta$ is not a lower bound of E . Therefore, $\alpha = \inf E$. $\square$

1.3 The Real Field

There exists an ordered field $\mathbb{R}$ which has the [least upper bound property](#). Moreover, $\mathbb{R}$ contains $\mathbb{Q}$ as a subfield, i.e., $\mathbb{Q} \subseteq \mathbb{R}$ and the operations of addition and multiplication on $\mathbb{R}$ coincide with those on $\mathbb{Q}$.

We can construct $\mathbb{R}$ from $\mathbb{Q}$. But for now, we will assume the existence of $\mathbb{R}$ and its properties.

Theorem 1.3.1. *For any $x \in \mathbb{R}^+$ and $y \in \mathbb{R}$, there exists some $n \in \mathbb{N}$ such that $nx > y$. This theorem is called the **Archimedean property** of $\mathbb{R}$.*

Proof. Suppose $x, y \in \mathbb{R}$ and $x > 0$. Let E be defined as follows:

$$E := \{nx \mid nx \leq y, n \in \mathbb{N}\}.$$

If $E = \emptyset$, which implies that $x > y$, then $n = 1$ satisfies $nx > y$. Otherwise, E is nonempty and bounded above by y . By the least upper bound property, there exists $\alpha = \sup E \in \mathbb{R}$. Then we have the following:

- $\alpha - x$ is not an upper bound of $E \implies$ There exists some $m \in \mathbb{N}$ such that $mx > \alpha - x$.
- $mx > \alpha - x \implies (m+1)x > \alpha \implies (m+1)x \notin E \implies (m+1)x > y$.

Therefore, we can take $n = m + 1$ to complete the proof. $\square$

Corollary 1.3.2. *For any $\varepsilon \in \mathbb{R}^+$ and $b \in \mathbb{R}$ with $b > 1$, there exists some $n \in \mathbb{N}$ such that $b^{-n} < \varepsilon$.*

Proof. For any $n \in \mathbb{N}$ and $x \in \mathbb{R}$ with $x \geq -1$, we have

$$(1+x)^n \geq 1+nx.$$

Let $a \in \mathbb{R}^+$ and $\varepsilon \in \mathbb{R}^+$ be given. By [Theorem 1.3.1](#), there exists some $n \in \mathbb{N}$ such that $na > \frac{1}{\varepsilon}$. Then we have

$$(1+a)^n \geq 1+na > na > \frac{1}{\varepsilon} \implies (1+a)^{-n} < \varepsilon.$$

Since $b > 1$, we can take $a = b - 1 > 0$ to complete the proof. $\square$

Theorem 1.3.3. *Any nonempty subset of $\mathbb{Z}$ which is bounded below in $\mathbb{R}$ has a unique least element, and any nonempty subset of $\mathbb{Z}$ which is bounded above in $\mathbb{R}$ has a unique greatest element.*

Proof. Let E be a nonempty subset of $\mathbb{Z}$ which is bounded below in $\mathbb{R}$. Then there exists a lower bound $\alpha \in \mathbb{R}$ of E . Then we have the following:

- If $\alpha \geq 0$, then $-1 \in \mathbb{Z}$ is a lower bound of E .
- If $\alpha < 0$, then by [Theorem 1.3.1](#), there exists some $n \in \mathbb{N}$ such that $n > -\alpha$, which implies that $-n \in \mathbb{Z}$ is a lower bound of E .

Therefore, E is bounded below in $\mathbb{Z}$. By [Theorem 1.2.2](#), E has a unique least element.

Let F be a nonempty subset of $\mathbb{Z}$ which is bounded above in $\mathbb{R}$. Then there exists an upper bound $\beta \in \mathbb{R}$ of F . Then we have the following:

- By [Theorem 1.3.1](#), there exists some $n \in \mathbb{N}$ such that $n > \beta$, which implies that $n \in \mathbb{Z}$ is an upper bound of F .

Therefore, F is bounded above in $\mathbb{Z}$. By [Theorem 1.2.2](#), F has a unique greatest element. $\square$

Definition 1.3.4. For any $x \in \mathbb{R}$, the **floor function** $\lfloor x \rfloor$ and **ceiling function** $\lceil x \rceil$ are defined as follows:

$$\lfloor x \rfloor := \max\{k \in \mathbb{Z} \mid k \leq x\}, \quad \lceil x \rceil := \min\{k \in \mathbb{Z} \mid k \geq x\}$$

[Theorem 1.3.3](#) guarantees the *existence* and *uniqueness* of $\lfloor x \rfloor$ and $\lceil x \rceil$.

Theorem 1.3.5. For any $x, y \in \mathbb{R}$ with $x < y$, there exists some $p \in \mathbb{Q}$ such that $x < p < y$. This theorem states that $\mathbb{Q}$ is **dense** in $\mathbb{R}$.

Proof. Let $\varepsilon := y - x > 0$. By [Theorem 1.3.1](#), there exists some $n \in \mathbb{N}$ such that $\frac{1}{n} < \varepsilon$. Let S be defined as follows:

$$S := \left\{ k \in \mathbb{Z} \mid \frac{k}{n} < x \right\}.$$

S is nonempty since $n\lfloor x \rfloor - 1 \in S$, and is bounded above by $nx \in \mathbb{R}$. By [Theorem 1.3.3](#), S has a greatest element, say m .

$$x < \frac{m+1}{n} < x + \frac{1}{n} < x + (y - x) = y.$$

Therefore, we can take $p = \frac{m+1}{n} \in \mathbb{Q}$ to complete the proof. $\square$

Theorem 1.3.6. *For each $x \in \mathbb{R}^+$ and $n \in \mathbb{N}$, there exists a unique $y \in \mathbb{R}^+$ such that $y^n = x$. We denote y by $\sqrt[n]{x}$ or $x^{\frac{1}{n}}$.*

Proof. Let E be defined as follows:

$$E := \{t \in \mathbb{R}^+ \mid t^n < x\}.$$

E is nonempty since $2^{-1} \in E$ if $x \geq 1$, and $x^2 \in E$ otherwise. Also, E is bounded above by $\max\{x, 1\}$. By the least upper bound property, there exists $\alpha = \sup E \in \mathbb{R}^+$. We will show that $\alpha^n = x$.

For any $a, b \in \mathbb{R}^+$ with $a < b$, we have

$$b^n - a^n = (b - a) \sum_{k=0}^{n-1} b^{n-1-k} a^k < (b - a) n b^{n-1}.$$

Assume that $\alpha^n < x$. By Theorem 1.3.5, there exists some $h \in \mathbb{R}^+$ such that

$$h < \min\left\{1, \frac{x - \alpha^n}{n(\alpha + 1)^{n-1}}\right\}.$$

Put $a = \alpha$ and $b = \alpha + h$. Then we have

$$\begin{aligned} (\alpha + h)^n - \alpha^n &< hn(\alpha + h)^{n-1} < hn(\alpha + 1)^{n-1} \\ \implies (\alpha + h)^n &< \alpha^n + hn(\alpha + 1)^{n-1} < \alpha^n + (x - \alpha^n) = x. \end{aligned}$$

Therefore, we have $\alpha + h \in E$, which is a contradiction.

Assume that $\alpha^n > x$. By Theorem 1.3.5, there exists some $h \in \mathbb{R}^+$ such that

$$h < \min\left\{1, \alpha, \frac{\alpha^n - x}{n(\alpha + 1)^{n-1}}\right\}.$$

Put $a = \alpha - h$ and $b = \alpha$. Then we have

$$\begin{aligned} \alpha^n - (\alpha - h)^n &< hn\alpha^{n-1} < hn(\alpha + 1)^{n-1} \\ \implies (\alpha - h)^n &> \alpha^n - hn(\alpha + 1)^{n-1} > \alpha^n - (\alpha^n - x) = x. \end{aligned}$$

Therefore, we have $\alpha - h$ is an upper bound of E , which is a contradiction.

Therefore, we have $\alpha^n = x$, which guarantees the *existence* of y . For any $\alpha, \beta \in \mathbb{R}^+$, we have $\alpha \neq \beta \implies \alpha^n \neq \beta^n$, which guarantees the *uniqueness* of y . $\square$

Definition 1.3.7. Let $y \in [0, 1)$ and $b \in \mathbb{N}$ be given with $b \geq 2$. A sequence $\{a_n\}_{n=1}^\infty$ is called the **greedy base b expansion** of y if for each $n \in \mathbb{N}$, we have

$$0 \leq y - \sum_{k=1}^n \frac{a_k}{b^k} < \frac{1}{b^n}$$

where $a_k \in S_b := \{0, 1, 2, \dots, b-1\}$ for all $k \in \mathbb{N}$. Especially, if $b = 10$, then the sequence is called the **greedy decimal expansion** of y .

Theorem 1.3.8. For any $y \in [0, 1)$, the greedy base b expansion of y exists and is unique.

Proof. We define sequences $\{t_n\}_{n=1}^\infty$, $\{T_n\}_{n=1}^\infty$ and $\{a_n\}_{n=1}^\infty$ as follows:

- $t_1 = by$, $T_1 := \{t \in S_b \mid t \leq t_1\}$, $a_1 = \max T_1$.
 - $t_2 = b^2y - ba_1$, $T_2 := \{t \in S_b \mid t \leq t_2\}$, $a_2 = \max T_2$.
 - $\vdots$
 - $t_n = b^n y - \sum_{k=1}^{n-1} a_k b^{n-k}$, $T_n := \{t \in S_b \mid t \leq t_n\}$, $a_n = \max T_n$
- for all $n \in \mathbb{N}$ with $n \geq 2$,

Suppose that $t_n \geq 0$ for some $n \in \mathbb{N}$. Since $0 \in T_n$ and T_n is bounded above, T_n has a greatest element by [Theorem 1.2.2](#), which implies that $\{a_n\}_{n=1}^\infty$ is well-defined. Let $P(n) : t_n \geq 0$ and a_n is well-defined.

- **Base case:** $P(1)$ is true since $t_1 = by \geq 0$.
- **Inductive step:** Suppose that $P(n)$ is true for some $n \in \mathbb{N}$. Then we have

$$\begin{aligned} t_{n+1} &= b^{n+1}y - \sum_{k=1}^n a_k b^{n+1-k} \\ &= b \left(b^n y - \sum_{k=1}^n a_k b^{n-k} \right) \\ &= b \left(b^n y - \sum_{k=1}^{n-1} a_k b^{n-k} - a_n \right) \\ &= b(t_n - a_n) \geq 0. \end{aligned}$$

Therefore, $P(n+1)$ is true whenever $P(n)$ is true.

By the principle of mathematical induction, $P(n)$ is true for all $n \in \mathbb{N}$. Then, by the definition of a_n , we have

$$\begin{aligned} a_n &\leq b^n y - \sum_{k=1}^{n-1} a_k b^{n-k} < a_n + 1 \\ \implies 0 &\leq b^n y - \sum_{k=1}^n a_k b^{n-k} < 1 \\ \implies 0 &\leq y - \sum_{k=1}^n \frac{a_k}{b^k} < \frac{1}{b^n} \end{aligned}$$

for each $n \in \mathbb{N}$. Therefore, the *existence* of the *greedy base b expansion* of y is guaranteed.

Next, suppose that there exist two sequences $\{a_n\}_{n=1}^{\infty}$ and $\{c_n\}_{n=1}^{\infty}$ such that for each $n \in \mathbb{N}$, we have

$$\begin{aligned} \bullet \quad 0 &\leq y - \sum_{k=1}^n \frac{a_k}{b^k} < \frac{1}{b^n} \\ \bullet \quad 0 &\leq y - \sum_{k=1}^n \frac{c_k}{b^k} < \frac{1}{b^n} \end{aligned}$$

where $a_k \in S_b = \{0, 1, 2, \dots, b-1\}$ for all $k = 1, 2, \dots, n$. We will show that for each $n \in \mathbb{N}$, we have $a_n = c_n$. Let $Q(n) : a_k = c_k$ for all $k = 1, 2, \dots, n$.

• **Base case:** We have

$$\begin{aligned} - \quad 0 &\leq y - \frac{a_1}{b} < \frac{1}{b} \implies \frac{a_1}{b} \leq y < \frac{a_1 + 1}{b}. \\ - \quad 0 &\leq y - \frac{c_1}{b} < \frac{1}{b} \implies \frac{c_1}{b} \leq y < \frac{c_1 + 1}{b}. \end{aligned}$$

By combining the two inequalities, we have

$$\begin{aligned} - \quad \frac{a_1}{b} &\leq y < \frac{c_1 + 1}{b} \implies a_1 < c_1 + 1 \implies a_1 \leq c_1. \\ - \quad \frac{c_1}{b} &\leq y < \frac{a_1 + 1}{b} \implies c_1 < a_1 + 1 \implies c_1 \leq a_1. \end{aligned}$$

Therefore, $a_1 = c_1$, which shows that $Q(1)$ is true.

• **Inductive step:** Suppose that $Q(n)$ is true for some $n \in \mathbb{N}$. We have

$$- \quad 0 \leq y - \sum_{k=1}^{n+1} \frac{a_k}{b^k} < \frac{1}{b^{n+1}} \implies \sum_{k=1}^{n+1} \frac{a_k}{b^k} \leq y < \sum_{k=1}^{n+1} \frac{a_k}{b^k} + \frac{1}{b^{n+1}}.$$

$$- 0 \leq y - \sum_{k=1}^{n+1} \frac{c_k}{b^k} < \frac{1}{b^{n+1}} \implies \sum_{k=1}^{n+1} \frac{c_k}{b^k} \leq y < \sum_{k=1}^{n+1} \frac{c_k}{b^k} + \frac{1}{b^{n+1}}.$$

By combining the two inequalities, we have

$$\begin{aligned} - \sum_{k=1}^{n+1} \frac{a_k}{b^k} \leq y < \sum_{k=1}^{n+1} \frac{c_k}{b^k} + \frac{1}{b^{n+1}} &\implies \sum_{k=1}^{n+1} \frac{a_k - c_k}{b^k} < \frac{1}{b^{n+1}}. \\ - \sum_{k=1}^{n+1} \frac{c_k}{b^k} \leq y < \sum_{k=1}^{n+1} \frac{a_k}{b^k} + \frac{1}{b^{n+1}} &\implies \sum_{k=1}^{n+1} \frac{c_k - a_k}{b^k} < \frac{1}{b^{n+1}}. \end{aligned}$$

By the inductive hypothesis, we have $a_k = c_k$ for all $k = 1, 2, \dots, n$. Therefore, we have

$$\begin{aligned} - \frac{a_{n+1} - c_{n+1}}{b^{n+1}} < \frac{1}{b^{n+1}} &\implies a_{n+1} - c_{n+1} < 1 \implies a_{n+1} \leq c_{n+1}. \\ - \frac{c_{n+1} - a_{n+1}}{b^{n+1}} < \frac{1}{b^{n+1}} &\implies c_{n+1} - a_{n+1} < 1 \implies c_{n+1} \leq a_{n+1}. \end{aligned}$$

Therefore, $a_{n+1} = c_{n+1}$, and thus $Q(n+1)$ is true whenever $Q(n)$ is true.

By the principle of mathematical induction, $Q(n)$ is true for all $n \in \mathbb{N}$. Therefore, the *uniqueness* of the *greedy base b expansion* of y is guaranteed. $\square$

Theorem 1.3.9. *Let $\{a_n\}_{n=1}^{\infty}$ be the greedy base b expansion of $a \in [0, 1)$, and let $\{c_n\}_{n=1}^{\infty}$ be the greedy base b expansion of $c \in [0, 1)$ for some $b \in \mathbb{N}$ with $b \geq 2$. If $a_n = c_n$ for all $n \in \mathbb{N}$, then $a = c$.*

Proof. Suppose that $a \neq c$. Without loss of generality, assume that $a < c$. Then, by [Corollary 1.3.2](#), there exists some $n \in \mathbb{N}$ such that

$$b^n > \frac{1}{c - a} \implies \frac{1}{b^n} < c - a.$$

By the definition of the *greedy base b expansion*, we have

$$\begin{aligned} \bullet \quad 0 \leq a - \sum_{k=1}^n \frac{a_k}{b^k} < \frac{1}{b^n} &\implies \sum_{k=1}^n \frac{a_k}{b^k} + \frac{1}{b^n} \leq a + \frac{1}{b^n}. \\ \bullet \quad 0 \leq c - \sum_{k=1}^n \frac{c_k}{b^k} < \frac{1}{b^n} &\implies c < \sum_{k=1}^n \frac{c_k}{b^k} + \frac{1}{b^n}. \end{aligned}$$

By the assumption that $a_n = c_n$ for all $n \in \mathbb{N}$, we have

$$\sum_{k=1}^n \frac{a_k}{b^k} = \sum_{k=1}^n \frac{c_k}{b^k}.$$

Therefore, we have a contradiction:

$$\sum_{k=1}^n \frac{a_k}{b^k} + \frac{1}{b^n} \leq a + \frac{1}{b^n} < c < \sum_{k=1}^n \frac{c_k}{b^k} + \frac{1}{b^n}.$$

□

Theorem 1.3.10. *Let $b \in \mathbb{N}$ with $b \geq 2$ and $\{a_n\}_{n=1}^\infty$ be given with $a_n \in S_b := \{0, 1, 2, \dots, b-1\}$ for all $n \in \mathbb{N}$. If there exists no $y \in [0, 1)$ such that $\{a_n\}_{n=1}^\infty$ is the greedy base b expansion of y , then there exists some $N \in \mathbb{N}$ such that*

$$a_n = b - 1 \quad \text{for all } n \geq N.$$

Proof. We will prove the contrapositive of the statement: For all $N \in \mathbb{N}$, if there exists some $n \in \mathbb{N}$ such that

$$n \geq N \text{ and } a_n \neq b - 1,$$

then there exists $y \in [0, 1)$ such that $\{a_n\}_{n=1}^\infty$ is the greedy base b expansion of y .

Assume that the condition of the above claim holds. Let $x_n = \sum_{k=1}^n \frac{a_k}{b^k}$ for each $n \in \mathbb{N}$. Since $a_k \in S_b$ for all $k \in \mathbb{N}$, we have

$$0 \leq x_n \leq \sum_{k=1}^n \frac{b-1}{b^k} = 1 - \frac{1}{b^n} < 1 \quad \text{for all } n \in \mathbb{N}.$$

Then $X := \{x_n \mid n \in \mathbb{N}\}$ is bounded above. By the [least upper bound property](#), there exists $y = \sup X \geq 0$. First, we will show that $y < 1$. By the assumption, there exists $m \in \mathbb{N}$ such that $a_m \neq b - 1$. Then, for all $n > m$, we have

$$\begin{aligned} x_n &= x_m + \sum_{k=m+1}^n \frac{a_k}{b^k} \leq x_m + \sum_{k=m+1}^n \frac{b-1}{b^k} \\ \implies x_n &\leq x_m + \frac{1}{b^m} - \frac{1}{b^n} < x_m + \frac{1}{b^m}. \end{aligned}$$

Since $a_m \leq b - 2$, we have

$$\begin{aligned} x_m + \frac{1}{b^m} &= \sum_{k=1}^{m-1} \frac{a_k}{b^k} + \frac{a_m + 1}{b^m} \\ &\leq \sum_{k=1}^{m-1} \frac{a_k}{b^k} + \frac{b-1}{b^m} \\ &\leq \sum_{k=1}^m \frac{b-1}{b^k} = 1 - \frac{1}{b^m}. \end{aligned}$$

We have $x_n \leq x_m$ for all $n \leq m$, and $x_n < x_m + \frac{1}{b^m} \leq 1 - \frac{1}{b^m}$ for all $n > m$. Thus, $1 - \frac{1}{b^m}$ is an upper bound of X . Since $y = \sup X$, we have

$$y \leq 1 - \frac{1}{b^m} < 1 \implies y \in [0, 1).$$

Next, we will show that $\{a_n\}_{n=1}^\infty$ is the *greedy base b expansion* of y . Let $n \in \mathbb{N}$ be fixed. By the assumption, there exists $m \in \mathbb{N}$ such that $m \geq n$ and $a_m \leq b - 2$.

$$\begin{aligned} x_m &= x_n + \sum_{k=n+1}^m \frac{a_k}{b^k} < x_n + \sum_{k=n+1}^m \frac{b-1}{b^k} \\ \implies x_m &< x_n + \frac{1}{b^n} - \frac{1}{b^m} \implies x_m + \frac{1}{b^m} < x_n + \frac{1}{b^n}. \end{aligned}$$

Since $a_m \leq b - 2$, $x_m + \frac{1}{b^m}$ is an upper bound of X by the same argument as above. Since $y = \sup X$, we have

$$y \leq x_m + \frac{1}{b^m} < x_n + \frac{1}{b^n} \implies 0 \leq y - x_n < \frac{1}{b^n}.$$

Thus, $\{a_n\}_{n=1}^\infty$ is the *greedy base b expansion* of y . □

Theorem 1.3.11. *Let $b \in \mathbb{N}$ with $b \geq 2$ and $\{a_n\}_{n=1}^\infty$ be given with $a_n \in S_b := \{0, 1, 2, \dots, b-1\}$ for all $n \in \mathbb{N}$. If there exists some $N \in \mathbb{N}$ such that*

$$a_n = b - 1 \quad \text{for all } n \geq N,$$

then there exists no $y \in [0, 1)$ such that $\{a_n\}_{n=1}^\infty$ is the greedy base b expansion of y . This theorem is the converse of [Theorem 1.3.10](#).

Proof. Suppose that there exists some $y \in [0, 1)$ such that $\{a_n\}_{n=1}^\infty$ is the *greedy base b expansion* of y . Then we have

$$0 \leq y - \sum_{k=1}^n \frac{a_k}{b^k} < \frac{1}{b^n} \quad \text{for all } n \in \mathbb{N}.$$

Let $N \in \mathbb{N}$ be such that $a_n = b - 1$ for all $n \geq N$. Let $x_n = \sum_{k=1}^n \frac{a_k}{b^k}$ for each $n \in \mathbb{N}$.

Then, for all $n \geq N$, we have

$$\begin{aligned} x_n &= x_{N-1} + \sum_{k=N}^n \frac{b-1}{b^k} = x_{N-1} + \frac{1}{b^{N-1}} - \frac{1}{b^n} \\ \implies y - x_n &= y - x_{N-1} - \frac{1}{b^{N-1}} + \frac{1}{b^n} \geq 0 \\ \implies \frac{1}{b^n} &\geq x_{N-1} + \frac{1}{b^{N-1}} - y. \end{aligned}$$

Since $0 \leq y - x_{N-1} < \frac{1}{b^{N-1}}$, we have $x_{N-1} + \frac{1}{b^{N-1}} - y > 0$. Then, by [Corollary 1.3.2](#), there exists some $n \in \mathbb{N}$ such that

$$\frac{1}{b^n} < x_{N-1} + \frac{1}{b^{N-1}} - y,$$

which is a contradiction. Therefore, there exists no $y \in [0, 1)$ such that $\{a_n\}_{n=1}^\infty$ is the greedy base b expansion of y . $\square$

1.4 The Extended Real Number System

The Extended real number system $\overline{\mathbb{R}}$ is defined as

$$\overline{\mathbb{R}} := \mathbb{R} \cup \{-\infty, +\infty\}.$$

We define the following operations on $\overline{\mathbb{R}}$:

- $x + (+\infty) = +\infty$ for all $x \in \mathbb{R} \cup \{\infty\}$.
- $x + (-\infty) = -\infty$ for all $x \in \mathbb{R} \cup \{-\infty\}$.
- $x \cdot (\pm\infty) = \pm\infty$ for all $x \in \mathbb{R}^+ \cup \{\infty\}$.
- $x \cdot (\pm\infty) = \mp\infty$ for all $x \in \mathbb{R}^- \cup \{-\infty\}$.
- $\frac{x}{\pm\infty} = 0$ for all $x \in \mathbb{R}$.

$\overline{\mathbb{R}}$ is not a field, but it is an ordered set with the following order relation:

$$-\infty < x < +\infty \quad \text{for all } x \in \mathbb{R}.$$

Theorem 1.4.1. *Every subset of $\overline{\mathbb{R}}$ has a least upper bound and a greatest lower bound in $\overline{\mathbb{R}}$.*

Proof. Let $E \subseteq \overline{\mathbb{R}}$. If $E = \emptyset$, then $\sup E = -\infty$ and $\inf E = +\infty$. If $E \neq \emptyset$, then we have the following cases:

- If E is bounded above in $\mathbb{R}$, then $\sup E$ exists in $\mathbb{R}$ by the least upper bound property of $\mathbb{R}$.
- If E is not bounded above in $\mathbb{R}$, then $\sup E = +\infty$.
- If E is bounded below in $\mathbb{R}$, then $\inf E$ exists in $\mathbb{R}$ by the greatest lower bound property of $\mathbb{R}$.
- If E is not bounded below in $\mathbb{R}$, then $\inf E = -\infty$.

$\square$

1.5 The Complex Field

The complex field $\mathbb{C}$ is defined as

$$\mathbb{C} := \{(x, y) \mid x, y \in \mathbb{R}\}.$$

We define the following operations on $\mathbb{C}$:

- $(x, y) + (u, v) = (x + u, y + v)$ for all $(x, y), (u, v) \in \mathbb{C}$.
- $(x, y) \cdot (u, v) = (xu - yv, xv + yu)$ for all $(x, y), (u, v) \in \mathbb{C}$.

$\mathbb{C}$ is a field with the following properties:

- The additive identity is $(0, 0)$ and the multiplicative identity is $(1, 0)$.
- The additive inverse of (x, y) is $(-x, -y)$ and the multiplicative inverse of (x, y) is

$$\left(\frac{x}{x^2 + y^2}, -\frac{y}{x^2 + y^2} \right)$$

for all $(x, y) \in \mathbb{C}$ with $(x, y) \neq (0, 0)$.

Remark. The function

$$f : \{(x, 0) \mid x \in \mathbb{R}\} \subseteq \mathbb{C} \rightarrow \mathbb{R}$$

defined by $f((x, 0)) = x$ is an *isomorphism*. Then we can identify $\mathbb{R}$ with the subfield $\{(x, 0) \mid x \in \mathbb{R}\}$ of $\mathbb{C}$. Now we will denote $(x, 0) \in \mathbb{C}$ by $x \in \mathbb{R}$.

Remark. We define $i = (0, 1) \in \mathbb{C}$. Then we have $i^2 = -1$, and any complex number $(x, y) \in \mathbb{C}$ can be written as

$$(x, y) = x + yi.$$

Definition 1.5.1. Let $z = x + yi \in \mathbb{C}$ for some $x, y \in \mathbb{R}$. The **real part** of z is defined as

$$\Re(z) = x \text{ or } \operatorname{Re}(z) = x,$$

the **imaginary part** of z is defined as

$$\Im(z) = y \text{ or } \operatorname{Im}(z) = y,$$

and the **complex conjugate** of z is defined as

$$\bar{z} = x - yi.$$

Theorem 1.5.2. Let $z, w \in \mathbb{C}$. Then we have the following properties:

- $\overline{(\overline{z})} = z$.
- $z = \overline{z} \iff z \in \mathbb{R}$.
- $\overline{z + w} = \overline{z} + \overline{w}$.
- $\overline{zw} = \overline{z} \cdot \overline{w}$.
- $z + \overline{z} = 2\operatorname{Re}(z)$ and $z - \overline{z} = 2i\operatorname{Im}(z)$.
- $z\overline{z} = x^2 + y^2 \in \mathbb{R}^+ \cup \{0\}$.

Proof. The proof follows directly from the definitions and properties of the operations on $\mathbb{C}$. $\square$

Definition 1.5.3. Let $z \in \mathbb{C}$. the **modulus** of z is defined as

$$|z| = \sqrt{z\overline{z}} = \sqrt{x^2 + y^2} \in \mathbb{R}^+ \cup \{0\}.$$

The existence and uniqueness of $|z|$ follows from the last part of [Theorem 1.5.2](#) and [Theorem 1.3.6](#).

Theorem 1.5.4. Let $a_1, a_2, \dots, a_n \in \mathbb{C}$ and $b_1, b_2, \dots, b_n \in \mathbb{C}$. Then we have

$$\left| \sum_{k=1}^n a_k \overline{b_k} \right|^2 \leq \sum_{k=1}^n |a_k|^2 \cdot \sum_{k=1}^n |b_k|^2.$$

This theorem is called the **Cauchy-Schwarz inequality**.

Proof. Let $i, j \in \mathbb{N}$ with $1 \leq i < j \leq n$. Then we have

$$\begin{aligned} |a_i b_j - a_j b_i|^2 &= (a_i b_j - a_j b_i) \overline{(a_i b_j - a_j b_i)} \\ &= |a_i|^2 |b_j|^2 + |a_j|^2 |b_i|^2 - (a_i \overline{b_i} \cdot \overline{a_j} b_j + \overline{a_i} b_i \cdot a_j \overline{b_j}). \end{aligned}$$

Consider the following sums:

$$\begin{aligned} \sum_{i < j} (|a_i|^2 |b_j|^2 + |a_j|^2 |b_i|^2) &= \sum_{i < j} (|a_i|^2 |b_j|^2) + \sum_{j < i} (|a_i|^2 |b_j|^2) \\ &= \sum_{i \neq j} (|a_i|^2 |b_j|^2) \\ &= \sum_{k=1}^n |a_k|^2 \cdot \sum_{k=1}^n |b_k|^2 - \sum_{k=1}^n |a_k|^2 |b_k|^2. \end{aligned}$$

$$\begin{aligned}
 \sum_{i < j} (a_i \bar{b}_i \cdot \bar{a}_j b_j + \bar{a}_i b_i \cdot a_j \bar{b}_j) &= \sum_{i \neq j} (a_i \bar{b}_i \cdot \bar{a}_j b_j) \\
 &= \sum_{i=1}^n a_i \bar{b}_i \cdot \sum_{j=1}^n \bar{a}_j b_j - \sum_{k=1}^n a_k \bar{b}_k \cdot \bar{a}_k b_k \\
 &= \left| \sum_{k=1}^n a_k \bar{b}_k \right|^2 - \sum_{k=1}^n |a_k|^2 |b_k|^2.
 \end{aligned}$$

Therefore, we have

$$\sum_{i < j} |a_i b_j - a_j b_i|^2 = \sum_{k=1}^n |a_k|^2 \cdot \sum_{k=1}^n |b_k|^2 - \left| \sum_{k=1}^n a_k \bar{b}_k \right|^2 \geq 0.$$

□

Corollary 1.5.5. *Let $a_1, a_2, \dots, a_n \in \mathbb{R}$ and $b_1, b_2, \dots, b_n \in \mathbb{R}$. Then we have*

$$\left(\sum_{k=1}^n a_k b_k \right)^2 \leq \sum_{k=1}^n a_k^2 \cdot \sum_{k=1}^n b_k^2.$$

Theorem 1.5.6. *Let $z, w \in \mathbb{C}$. Then we have the following properties:*

- (A) $|zw| = |z| \cdot |w|$.
- (B) $|z + w| \leq |z| + |w|$.
- (C) $||z| - |w|| \leq |z - w|$.

Proof. Let $z = x + yi$ and $w = u + vi$ for some $x, y, u, v \in \mathbb{R}$. Then we have

- (A) $|zw|^2 = (xu - yv)^2 + (xv + yu)^2 = (x^2 + y^2)(u^2 + v^2) = |z|^2 \cdot |w|^2$.
- (B) By [Corollary 1.5.5](#), we have $(xu + yv)^2 \leq (x^2 + y^2)(u^2 + v^2)$.

$$\begin{aligned}
 |z + w|^2 &= (x + u)^2 + (y + v)^2 \\
 &= x^2 + y^2 + u^2 + v^2 + 2(xu + yv) \\
 &\leq x^2 + y^2 + u^2 + v^2 + 2\sqrt{x^2 + y^2} \cdot \sqrt{u^2 + v^2} \\
 &= (|z| + |w|)^2.
 \end{aligned}$$

- (C) Again, by [Corollary 1.5.5](#), we have

$$\begin{aligned}
 |z - w|^2 &= (x - u)^2 + (y - v)^2 \\
 &= x^2 + y^2 + u^2 + v^2 - 2(xu + yv) \\
 &\geq x^2 + y^2 + u^2 + v^2 - 2\sqrt{x^2 + y^2} \cdot \sqrt{u^2 + v^2} \\
 &= (|z| - |w|)^2.
 \end{aligned}$$

□

1.6 Euclidean Spaces

For $n \in \mathbb{N}$, we define the **Euclidean n -space** $\mathbb{R}^n$ as the set of all ordered n -tuples of real numbers, i.e.,

$$\mathbb{R}^n := \{ \mathbf{x} = (x_1, x_2, \dots, x_n) \mid x_k \in \mathbb{R} \text{ for all } k = 1, 2, \dots, n \}$$

where $x_1, x_2, \dots, x_n$ are called the **coordinates** of $\mathbf{x}$. The elements of $\mathbb{R}^n$ with $n > 1$ are called **vectors** or *points*, which are denoted by boldface letters.

Let $\mathbf{x}, \mathbf{y} \in \mathbb{R}^n$ and $\alpha \in \mathbb{R}$. We define the following operations on $\mathbb{R}^n$:

- $\mathbf{x} + \mathbf{y} = (x_1 + y_1, x_2 + y_2, \dots, x_n + y_n)$.
- $\alpha \mathbf{x} = (\alpha x_1, \alpha x_2, \dots, \alpha x_n)$.

Then $\mathbb{R}^n$ is a *vector space* over $\mathbb{R}$ with the zero vector $\mathbf{0} = (0, 0, \dots, 0)$.

Definition 1.6.1. Let $\mathbf{x}, \mathbf{y} \in \mathbb{R}^n$. The **inner product** of $\mathbf{x}$ and $\mathbf{y}$ is defined as

$$\mathbf{x} \cdot \mathbf{y} = \sum_{k=1}^n x_k y_k.$$

The **norm** of $\mathbf{x}$ is defined as

$$\|\mathbf{x}\| = \sqrt{\mathbf{x} \cdot \mathbf{x}} = \sqrt{\sum_{k=1}^n x_k^2}.$$

The **distance** between $\mathbf{x}$ and $\mathbf{y}$ is defined as

$$d(\mathbf{x}, \mathbf{y}) = \|\mathbf{x} - \mathbf{y}\| = \sqrt{\sum_{k=1}^n (x_k - y_k)^2}.$$

We can confirm that $d(x, y) = |x - y|$ in the case of $n = 1$.

Theorem 1.6.2. Let $\mathbf{x}, \mathbf{y}, \mathbf{z} \in \mathbb{R}^n$ and $\alpha \in \mathbb{R}$. Then we have the following properties:

- (A) $|\mathbf{x} \cdot \mathbf{y}| \leq \|\mathbf{x}\| \cdot \|\mathbf{y}\|$.
- (B) $\|\mathbf{x} + \mathbf{y}\| \leq \|\mathbf{x}\| + \|\mathbf{y}\|$.
- (C) $\|\mathbf{x} - \mathbf{y}\| \geq \|\mathbf{x}\| - \|\mathbf{y}\|$.

Proof. (A) By [Corollary 1.5.5](#), we have

$$|\mathbf{x} \cdot \mathbf{y}|^2 = \left| \sum_{k=1}^n x_k y_k \right|^2 \leq \sum_{k=1}^n x_k^2 \cdot \sum_{k=1}^n y_k^2 = \|\mathbf{x}\|^2 \cdot \|\mathbf{y}\|^2.$$

(B) By part (A), we have

$$\begin{aligned} \|\mathbf{x} + \mathbf{y}\|^2 &= (\mathbf{x} + \mathbf{y}) \cdot (\mathbf{x} + \mathbf{y}) = \|\mathbf{x}\|^2 + 2\mathbf{x} \cdot \mathbf{y} + \|\mathbf{y}\|^2 \\ &\leq \|\mathbf{x}\|^2 + 2|\mathbf{x} \cdot \mathbf{y}| + \|\mathbf{y}\|^2 \leq \|\mathbf{x}\|^2 + 2\|\mathbf{x}\| \cdot \|\mathbf{y}\| + \|\mathbf{y}\|^2 \\ &= (\|\mathbf{x}\| + \|\mathbf{y}\|)^2. \end{aligned}$$

(C) By part (B), we have

$$\begin{aligned} \|\mathbf{x}\| &= \|(\mathbf{x} - \mathbf{y}) + \mathbf{y}\| \leq \|\mathbf{x} - \mathbf{y}\| + \|\mathbf{y}\| \implies \|\mathbf{x}\| - \|\mathbf{y}\| \leq \|\mathbf{x} - \mathbf{y}\|. \\ \|\mathbf{y}\| &= \|(\mathbf{y} - \mathbf{x}) + \mathbf{x}\| \leq \|\mathbf{y} - \mathbf{x}\| + \|\mathbf{x}\| \implies \|\mathbf{y}\| - \|\mathbf{x}\| \leq \|\mathbf{x} - \mathbf{y}\|. \end{aligned}$$

□

Chapter 2

Basic Topology

2.1 Relations

A **relation** $\sim$ on a set S defines a subset R of $S \times S$, and vice versa.

$$R := \{(x, y) \in S \times S \mid x \sim y\}, \quad x \sim y \iff (x, y) \in R.$$

Definition 2.1.1. A relation $\sim$ on a set S is called an **equivalence relation** if it satisfies the following properties:

- Reflexivity: $x \sim x$ for all $x \in S$.
- Symmetry: If $x \sim y$, then $y \sim x$ for all $x, y \in S$.
- Transitivity: If $x \sim y$ and $y \sim z$, then $x \sim z$ for all $x, y, z \in S$.

Definition 2.1.2. A **partition** of a set S is a collection $\{A_\alpha\}_{\alpha \in I}$ of nonempty subsets of S such that

- $A_\alpha \cap A_\beta = \emptyset$ for all $\alpha, \beta \in I$ with $\alpha \neq \beta$.
- $\bigcup_{\alpha \in I} A_\alpha = S$.

Definition 2.1.3. Let $\sim$ be an equivalence relation on a set S . For each $x \in S$, the **equivalence class** of x is defined as

$$[x] := \{y \in S \mid y \sim x\}.$$

We denote the set of all equivalence classes of S by $S/\sim := \{[x] \mid x \in S\}$.

Theorem 2.1.4. *Let $\sim$ be an equivalence relation on a set S . For any $x, y \in S$, we have either*

$$[x] = [y] \quad \text{or} \quad [x] \cap [y] = \emptyset.$$

Therefore, $S/\sim$ is a partition of S .

Proof. Suppose $[x] \cap [y] \neq \emptyset$. Then there exists $z \in S$ such that $z \in [x]$ and $z \in [y]$. This implies that $z \sim x$ and $z \sim y$. By the symmetry and transitivity of $\sim$, we have $x \sim z$ and $z \sim y$, which implies that $x \sim y$.

Therefore, for any $w \in [x]$, we have $w \sim x$ and $x \sim y$, which implies that $w \sim y$, so $w \in [y]$. This shows that $[x] \subseteq [y]$. Similarly, we can show that $[y] \subseteq [x]$, so $[x] = [y]$. $\square$

Theorem 2.1.5. *Let $\{A_\alpha\}_{\alpha \in I}$ be a partition of a set S . Define a relation $\sim$ on S by*

$$x \sim y \iff x, y \in A_\alpha \text{ for some } \alpha \in I.$$

Then $\sim$ is an equivalence relation on S , and the equivalence classes of $\sim$ are precisely the sets A_α .

Proof. We need to show that $\sim$ is reflexive, symmetric, and transitive.

- Reflexivity: For any $x \in S$, there exists $\alpha \in I$ such that $x \in A_\alpha$, so $x \sim x$.
- Symmetry: If $x \sim y$, then there exists $\alpha \in I$ such that $x, y \in A_\alpha$, so $y \sim x$.
- Transitivity: If $x \sim y$ and $y \sim z$, then there exist $\alpha, \beta \in I$ such that $x, y \in A_\alpha$ and $y, z \in A_\beta$. Since $y \in A_\alpha \cap A_\beta$ and the sets $\{A_\alpha\}_{\alpha \in I}$ are disjoint, we must have $\alpha = \beta$, so $x, z \in A_\alpha$ and $x \sim z$.

Therefore, $\sim$ is an equivalence relation on S . The equivalence class of any $x \in S$ is precisely the set A_α containing x , so the equivalence classes of $\sim$ are exactly the sets A_α . $\square$

2.2 Functions

A *function* (or a *mapping*) is defined with 3 components.

- A **domain** X .
- A **codomain** Y .
- A **rule** f which associates each element $x \in X$ to a *unique* element $y \in Y$.

f is said to be a *function* of X into Y , which is denoted by $f : X \rightarrow Y$. For each $x \in X$, the *unique* element $y \in Y$ associated with x is denoted by $f(x)$.

Remark. Let $f : X \rightarrow Y$ be a function. For any subset $E \subseteq X$, we define the **image** of E under f as

$$f(E) := \{f(x) \mid x \in E\} \subseteq Y.$$

Especially, if $E = X$, then $f(X) \subseteq Y$ is called the **range** of f .

Remark. Let $f : X \rightarrow Y$ be a function. For any subset $E \subseteq Y$, we define the **inverse image** of E under f as

$$f^{-1}(E) := \{x \in X \mid f(x) \in E\} \subseteq X.$$

Definition 2.2.1. Let $f : X \rightarrow Y$ be a function. We define the properties of f as follows:

- f is **injective** (or *one-to-one*) if for any $x_1, x_2 \in X$,

$$f(x_1) = f(x_2) \implies x_1 = x_2.$$

- f is **surjective** (or *onto*) if $f(X) = Y$.
- f is **bijective** (or a *one-to-one correspondence*) if it is both injective and surjective.

Definition 2.2.2. Let $f : X \rightarrow Y$ be a bijective function. Then for any $y \in Y$, there exists a *unique* $x \in X$ such that $f(x) = y$. Therefore, we can define a function $f^{-1} : Y \rightarrow X$ by $f^{-1}(y) = x$. f^{-1} is called the **inverse function** of f .

Theorem 2.2.3. Let $f : X \rightarrow Y$ be a bijective function. Then the inverse function $f^{-1} : Y \rightarrow X$ is bijective.

Proof. Let $y_1, y_2 \in Y$. Suppose that $f^{-1}(y_1) = f^{-1}(y_2) = x$. Then, by [Definition 2.2.2](#), we have both $f(x) = y_1$ and $f(x) = y_2$. Since f is a function, we have $y_1 = y_2$, which implies that f^{-1} is injective.

By the definition of inverse image, we have $f^{-1}(Y) \subseteq X$. Also, for any $x \in X$, we have $f(x) \in Y$, which implies that $x \in f^{-1}(Y)$. Therefore, we have $f^{-1}(Y) \supseteq X \implies f^{-1}(Y) = X$, which implies that f^{-1} is surjective. $\square$

Definition 2.2.4. Let $f : X \rightarrow Y$ be a function. Let $E \subseteq X$ and $F \subseteq Y$ such that $f(E) \subseteq F$. Then we define the restriction functions as follows:

- The **restriction** of f to E is defined by

$$f|_E : E \rightarrow Y, \quad f|_E(x) = f(x) \quad \text{for all } x \in E.$$

- The **Corestriction** of f to F is defined by

$$f|_F : X \rightarrow F, \quad f|_F(x) = f(x) \quad \text{for all } x \in X.$$

- The **Bi-restriction** of f to E and F is defined by

$$f|_E^F : E \rightarrow F, \quad f|_E^F(x) = f(x) \quad \text{for all } x \in E.$$

We will simply call all of them as just *restriction*, if there is no ambiguity.

Theorem 2.2.5. *Let $f : X \rightarrow Y$ be a function. Let $E \subseteq X$ and $F \subseteq Y$ such that $f(E) \subseteq F$. Then we have the following properties:*

- (A) *If f is injective, then $f|_E^F$ is also injective.*
- (B) *If $f(E) = F$, then $f|_E^F$ is surjective.*
- (C) *If f is injective and $f(E) = F$, then $f|_E^F$ is bijective.*

Proof. (C) follows from (A) and (B). Let $x_1, x_2 \in E$.

- (A) Assume that f is injective. Then we have

$$f|_E^F(x_1) = f|_E^F(x_2) \implies f(x_1) = f(x_2) \implies x_1 = x_2.$$

- (B) Assume that $f(E) = F$. Then for any $y \in F$, there exists $x \in E$ such that $f(x) = y$. Therefore, we have $f|_E^F(E) = F$.

□

Definition 2.2.6. Let X, Y, Z be sets and $f : X \rightarrow Y, g : Y \rightarrow Z$ be functions. Then we define the **composition** of f and g as the function $g \circ f : X \rightarrow Z$ defined by

$$(g \circ f)(x) = g(f(x)) \quad \text{for all } x \in X.$$

Theorem 2.2.7. *Let X, Y, Z be sets and $f : X \rightarrow Y, g : Y \rightarrow Z$ be functions. Then we have the following properties:*

- (A) *If f and g are injective, then $g \circ f$ is also injective.*

(B) If f and g are surjective, then $g \circ f$ is also surjective.

(C) If f and g are bijective, then $g \circ f$ is also bijective.

Proof. (C) follows from (A) and (B). Let $x_1, x_2 \in X$.

(A) Assume that f and g are injective. Then we have

$$\begin{aligned} (g \circ f)(x_1) = (g \circ f)(x_2) &\implies g(f(x_1)) = g(f(x_2)) \\ &\implies f(x_1) = f(x_2) \implies x_1 = x_2. \end{aligned}$$

(B) Assume that f and g are surjective. Then for any $z \in Z$, there exists $y \in Y$ such that $g(y) = z$. Since f is surjective, there exists $x \in X$ such that $f(x) = y$. Therefore, we have $(g \circ f)(x) = g(f(x)) = g(y) = z$.

□

Theorem 2.2.8. Let S be a set of sets. For [Sections 2.3](#) and [2.4](#), we consider the relation $\sim$ defined on S by

$$X \sim Y \iff \text{There exists a bijection } f : X \rightarrow Y.$$

The relation $\sim$ is an equivalence relation on S .

Proof. Let $X, Y, Z \in S$.

- **Reflexivity:** The identity function $id_X : X \rightarrow X$ defined by $id_X(x) = x$ for all $x \in X$ is bijective. Therefore, we have $X \sim X$.
- **Symmetry:** Suppose that $X \sim Y$. Then there exists a bijection $f : X \rightarrow Y$. By [Theorem 2.2.3](#), the inverse function $f^{-1} : Y \rightarrow X$ is also bijective. Therefore, we have $Y \sim X$.
- **Transitivity:** Suppose that $X \sim Y$ and $Y \sim Z$. Then there exist bijections $f : X \rightarrow Y$ and $g : Y \rightarrow Z$. By [Theorem 2.2.7](#), the composition $g \circ f : X \rightarrow Z$ is also bijective. Therefore, we have $X \sim Z$.

□

2.3 Finite Sets

Let $J_n := \{1, 2, \dots, n\}$ for each $n \in \mathbb{N}$. A set X is said to be **finite** if $J_n \sim X$ for some $n \in \mathbb{N}$. In this case, n is called the **cardinality** of X , denoted by $|X|$. Especially, if $X = \emptyset$, we define $|X| = 0$. X is said to be **infinite** if X is not finite.

Lemma 2.3.1. Every subset of J_n is finite for each $n \in \mathbb{N}$.

Proof. We will prove the lemma by mathematical induction on n . Let $P(n)$: Every subset of J_n is finite.

- **Base case:** Let $n = 1$. Then $J_1 = \{1\}$, whose only subsets are $\emptyset$ and J_1 . $J_1 \sim J_1$ by $f := id_{J_1}$, and $\emptyset$ is finite by definition. Thus $P(1)$ is true.
- **Inductive step:** Assume that $P(n)$ is true for some $n \in \mathbb{N}$. Let $A \subseteq J_{n+1}$. We have two cases:
 - If $n+1 \notin A$, then $A \subseteq J_n$, which is finite by the inductive hypothesis.
 - If $n+1 \in A$, then $A = B \cup \{n+1\}$ for some $B \subseteq J_n$. By the inductive hypothesis, B is finite, so there exists a bijection $f : J_m \rightarrow B$ for some $m \in \mathbb{N}$. Let $g : J_{m+1} \rightarrow A$ be defined by

$$g(k) = \begin{cases} f(k) & \text{if } k \leq m \\ n+1 & \text{if } k = m+1 \end{cases}.$$

Then g is bijective, which implies that A is finite.

Thus $P(n+1)$ is true whenever $P(n)$ is true.

By the principle of mathematical induction, $P(n)$ is true for all $n \in \mathbb{N}$. □

Theorem 2.3.2. *Every subset of a finite set is finite.*

Proof. Let X be a finite set and $Y \subseteq X$. Then there exists a bijection $f : J_n \rightarrow X$ for some $n \in \mathbb{N}$. Let $Z = f^{-1}(Y) \subseteq J_n$. Since $Z \subseteq J_n$, Z is finite by [Lemma 2.3.1](#), so there exists a bijection $g : J_m \rightarrow Z$ for some $m \in \mathbb{N}$.

By [Theorem 2.2.5](#), $f|_Z^Y$ is bijective. Then, by [Theorem 2.2.7](#), the composition $f|_Z^Y \circ g : J_m \rightarrow Y$ is bijective. Thus Y is finite. □

Theorem 2.3.3. *Let X be a finite ordered set. Then X has both a least element and a greatest element.*

Proof. Since X is finite, there exists a bijection $f : J_n \rightarrow X$ for some $n \in \mathbb{N}$. The elements of X can be listed as $X = \{a_1, a_2, \dots, a_n\}$, where $a_k = f(k)$ for each $k = 1, 2, \dots, n$.

First, we will show that X has a least element by mathematical induction on n . Let $P(n)$: Any finite ordered set of cardinality n has a least element.

- **Base case:** Let $n = 1$. Then $X = \{a_1\}$, which has a least element a_1 .

- **Inductive step:** Assume that $P(n)$ is true for some $n \in \mathbb{N}$. Let $X = \{a_1, a_2, \dots, a_{n+1}\}$. By the inductive hypothesis, the subset $\{a_1, a_2, \dots, a_n\}$ has a least element, say a_k for some $k = 1, 2, \dots, n$. We have two cases:
 - If $a_k \leq a_{n+1}$, then a_k is the least element of X .
 - If $a_{n+1} < a_k$, then a_{n+1} is the least element of X .

Thus $P(n+1)$ is true whenever $P(n)$ is true.

By the principle of mathematical induction, $P(n)$ is true for all $n \in \mathbb{N}$.

Next, we will show that X has a greatest element by mathematical induction on n . Let $Q(n)$: Any finite ordered set of cardinality n has a greatest element.

- **Base case:** Let $n = 1$. Then $X = \{a_1\}$, which has a greatest element a_1 .
- **Inductive step:** Assume that $Q(n)$ is true for some $n \in \mathbb{N}$. Let $X = \{a_1, a_2, \dots, a_{n+1}\}$. By the inductive hypothesis, the subset $\{a_1, a_2, \dots, a_n\}$ has a greatest element, say a_k for some $k = 1, 2, \dots, n$. We have two cases:
 - If $a_k \geq a_{n+1}$, then a_k is the greatest element of X .
 - If $a_{n+1} > a_k$, then a_{n+1} is the greatest element of X .

Thus $Q(n+1)$ is true whenever $Q(n)$ is true.

By the principle of mathematical induction, $Q(n)$ is true for all $n \in \mathbb{N}$. □

Lemma 2.3.4. Let J_m, J_n be given for some $m, n \in \mathbb{N}$. Then we have the following properties:

- (A) If there exists an injection $f : J_m \rightarrow J_n$, then $m \leq n$.
- (B) If there exists a surjection $f : J_m \rightarrow J_n$, then $m \geq n$.
- (C) If there exists a bijection $f : J_m \rightarrow J_n$, then $m = n$.

This lemma guarantees the uniqueness of the cardinality of a finite set. If $J_m \sim X$ and $J_n \sim X$, then $J_m \sim J_n$, which implies that $m = n$ by (C).

Proof. We will prove (A) by mathematical induction on m , (B) by mathematical induction on n , and (C) follows from (A) and (B).

- (A) Let $n \in \mathbb{N}$ be fixed. Let $P(m)$: If there exists an injective function $f_m : J_m \rightarrow J_n$, then $m \leq n$.
 - **Base case:** Let $m = 1$. Then $m \leq n$ holds.

- **Inductive step:** Assume that $P(m)$ is true for some $m \in \mathbb{N}$. If $n = 1$, there does not exist an injective function $f_{m+1} : J_{m+1} \rightarrow J_n$ since $f_{m+1}(1) \neq f_{m+1}(m+1)$, where $P(m+1)$ is true.

Otherwise, suppose that such a function f_{m+1} exists. We have two cases:

- $f_{m+1}(m+1) = n$. Let $f_m \equiv f_{m+1}|_{J_m}^{J_{n-1}}$. Then $f_m : J_m \rightarrow J_{n-1}$ is injective by [Theorem 2.2.5](#). By the inductive hypothesis, we have $m \leq n-1$, which implies that $m+1 \leq n$.
- $f_{m+1}(m+1) = x \neq n$. Let $g_n : J_n \rightarrow J_n$ be defined by

$$g_n(y) = \begin{cases} x, & \text{if } y = n \\ n, & \text{if } y = x \\ y, & \text{otherwise} \end{cases}.$$

Then g_n is bijective. Let $f_m \equiv (g_n \circ f_{m+1})|_{J_m}^{J_{n-1}}$. Then $f_m : J_m \rightarrow J_{n-1}$ is injective by [Theorems 2.2.5](#) and [2.2.7](#). By the inductive hypothesis, we have $m \leq n-1$, which implies that $m+1 \leq n$.

Thus $P(m+1)$ is true whenever $P(m)$ is true.

By the principle of mathematical induction, $P(m)$ is true for all $m \in \mathbb{N}$.

- (B) Let $m \in \mathbb{N}$ be fixed. Let $Q(n)$: If there exists a surjective function $f_n : J_m \rightarrow J_n$, then $m \geq n$.

- **Base case:** Let $n = 1$. Then $m \geq n$ holds.
- **Inductive step:** Assume that $Q(n)$ is true for some $n \in \mathbb{N}$. If $m = 1$, there does not exist a surjective function $f_{n+1} : J_m \rightarrow J_{n+1}$ since $f_{n+1}(1)$ is unique, where $Q(n+1)$ is true.

Otherwise, suppose that such a function f_{n+1} exists. We have two cases:

- $f_{n+1}(m) = n+1$. Let $K_{n+1} := (f_{n+1})^{-1}(\{n+1\}) \subseteq J_m$. Let $f_n : J_m \rightarrow J_n$ be defined by

$$f_n(y) = \begin{cases} f_{n+1}(y), & \text{if } y \in J_m \setminus K_{n+1} \\ n, & \text{if } y \in K_{n+1} \end{cases}.$$

Then f_n is surjective. By the inductive hypothesis, we have $m-1 \geq n$, which implies that $m \geq n+1$.

- $f_{n+1}(m) = x \neq n+1$. Let $g_{n+1} : J_{n+1} \rightarrow J_{n+1}$ be defined by

$$g_{n+1}(y) = \begin{cases} x, & \text{if } y = n+1 \\ n+1, & \text{if } y = x \\ y, & \text{otherwise} \end{cases}.$$

Then g_{n+1} is bijective. Let $K_{n+1} := (g_{n+1})^{-1}(\{n+1\}) \subseteq J_m$. Let $f_n : J_m \rightarrow J_n$ be defined by

$$f_n(y) = \begin{cases} (g_{n+1} \circ f_{n+1})(y), & \text{if } y \in J_m \setminus K_{n+1} \\ n, & \text{if } y \in K_{n+1} \end{cases}.$$

Then f_n is surjective by [Theorem 2.2.7](#). By the inductive hypothesis, we have $m-1 \geq n$, which implies that $m \geq n+1$.

Thus $Q(n+1)$ is true whenever $Q(n)$ is true.

By the principle of mathematical induction, $Q(n)$ is true for all $n \in \mathbb{N}$.

□

Lemma 2.3.5. *Let J_m, J_n be given for some $m, n \in \mathbb{N}$. Then we have the following properties:*

- (A) *If $m \leq n$, then there exists an injection $f : J_m \rightarrow J_n$.*
- (B) *If $m \geq n$, then there exists a surjection $f : J_m \rightarrow J_n$.*
- (C) *If $m = n$, then there exists a bijection $f : J_m \rightarrow J_n$.*

This lemma is the converse of [Lemma 2.3.4](#).

Proof. We will prove the lemma by explicitly constructing the functions.

- (A) If $m \leq n$, define $f : J_m \rightarrow J_n$ by $f(i) = i$ for all $i \in J_m$. Then f is an injection.
- (B) If $m \geq n$, define $f : J_m \rightarrow J_n$ by $f(i) = i$ for all $i \in J_n \subseteq J_m$ and $f(i) = n$ for all $i \in J_m \setminus J_n$. Then f is a surjection.
- (C) If $m = n$, define $f : J_m \rightarrow J_n$ by $f(i) = i$ for all $i \in J_m$. Then f is a bijection.

□

Theorem 2.3.6. *Let X, Y be finite sets. Then we have the following properties:*

- (A) *If there exists an injection $f : X \rightarrow Y$, then $|X| \leq |Y|$.*
- (B) *If there exists a surjection $f : X \rightarrow Y$, then $|X| \geq |Y|$.*
- (C) *If there exists a bijection $f : X \rightarrow Y$, then $|X| = |Y|$.*

Moreover, the converses of (A), (B), and (C) are also true.

The contraposition of (A) is called the **pigeonhole principle**, which states that if $|X| > |Y|$, then there exists no injection $f : X \rightarrow Y$.

Proof. Let $|X| = m$ and $|Y| = n$ for some $m, n \in \mathbb{N}$. Then there exist bijections $f_X : J_m \rightarrow X$ and $f_Y : J_n \rightarrow Y$. By Theorems 2.2.3 and 2.2.7, the composition

$$F \equiv (f_Y)^{-1} \circ f \circ f_X : J_m \rightarrow J_n$$

is injective, surjective, or bijective if f is injective, surjective, or bijective, respectively. Then, by Lemma 2.3.4, we have the desired results:

- (A) If f is injective, then F is injective, which implies that $m \leq n$.
- (B) If f is surjective, then F is surjective, which implies that $m \geq n$.
- (C) If f is bijective, then F is bijective, which implies that $m = n$.

Also, by Theorems 2.2.3 and 2.2.7, the composition

$$f \equiv f_Y \circ G \circ (f_X)^{-1} : X \rightarrow Y$$

is injective, surjective, or bijective if $G : J_m \rightarrow J_n$ is injective, surjective, or bijective, respectively. Then, by Lemma 2.3.5, we have the desired converses:

- (A) If $m \leq n$, then there exists an injective function $G : J_m \rightarrow J_n$, which implies that there exists an injective function $f : X \rightarrow Y$.
- (B) If $m \geq n$, then there exists a surjective function $G : J_m \rightarrow J_n$, which implies that there exists a surjective function $f : X \rightarrow Y$.
- (C) If $m = n$, then there exists a bijective function $G : J_m \rightarrow J_n$, which implies that there exists a bijective function $f : X \rightarrow Y$.

□

Lemma 2.3.7. Let X, Y be finite sets with $X \cap Y = \emptyset$. Then the union $X \cup Y$ is finite.

Proof. Let $m = |X|$ and $n = |Y|$. There exist bijections $f : J_m \rightarrow X$ and $g : J_n \rightarrow Y$. Let $h : J_{m+n} \rightarrow X \cup Y$ be defined by

$$h(i) = \begin{cases} f(i), & \text{if } i \leq m \\ g(i - m), & \text{if } m < i \leq m + n \end{cases}.$$

Then h is bijective, which implies that $X \cup Y$ is finite.

□

Theorem 2.3.8. *Let X, Y be finite sets. Then the union $X \cup Y$ and intersection $X \cap Y$ are finite.*

Proof. Let A, B, C be defined by

- $A = X \setminus Y \subseteq X$.
- $B = Y \setminus X \subseteq Y$.
- $C = X \cap Y \subseteq X$.

Then A, B, C are finite by [Theorem 2.3.2](#) and are disjoint each other. We have the followings:

- $X = (X \setminus Y) \cup (X \cap Y) = A \cup C$.
- $Y = (Y \setminus X) \cup (X \cap Y) = B \cup C$.
- $X \cup Y = ((X \setminus Y) \cup (Y \setminus X)) \cup (X \cap Y) = (A \cup B) \cup C$.

By [Lemma 2.3.7](#), $X, Y, X \cup Y$ are finite. □

Theorem 2.3.9. *Let $X_1, X_2, \dots, X_n$ be finite sets for some $n \in \mathbb{N}$. Then the union $\bigcup_{k=1}^n X_k$ and intersection $\bigcap_{k=1}^n X_k$ are finite.*

Proof. Since $\bigcap_{k=1}^n X_k \subseteq X_1$, $\bigcap_{k=1}^n X_k$ is finite by [Theorem 2.3.2](#) for any $n \in \mathbb{N}$. Let

$P(n) : \bigcup_{k=1}^n X_k$ is finite for some $n \in \mathbb{N}$, given that $X_1, X_2, \dots, X_n$ are finite sets.

- **Base case:** Let X_1 be a finite set. Then, $\bigcup_{k=1}^1 X_k = X_1$ is finite. Thus, $P(1)$ is true.

- **Inductive step:** Assume that $P(n)$ is true for some $n \in \mathbb{N}$.

Let $X_1, X_2, \dots, X_{n+1}$ be finite sets. Then we have

$$\bigcup_{k=1}^{n+1} X_k = \left(\bigcup_{k=1}^n X_k \right) \cup X_{n+1}.$$

By the inductive hypothesis, $\bigcup_{k=1}^n X_k$ is finite. Then, by [Theorem 2.3.8](#), $\bigcup_{k=1}^{n+1} X_k$ is finite. Thus $P(n+1)$ is true whenever $P(n)$ is true.

By the principle of mathematical induction, $P(n)$ is true for all $n \in \mathbb{N}$. $\square$

Lemma 2.3.10. *Let X, Y be finite sets. Then the cartesian product $X \times Y$ is finite.*

Proof. Let $m = |X|$ and $n = |Y|$. Then there exist bijections $f : J_m \rightarrow X$ and $g : J_n \rightarrow Y$. Let $h : J_{mn} \rightarrow X \times Y$ be defined by

$$h(i) = \left(f\left(\left\lceil \frac{i}{n} \right\rceil\right), g\left(i - n\left(\left\lceil \frac{i}{n} \right\rceil - 1\right)\right) \right) \text{ for all } i \in J_{mn}.$$

Then h is bijective, which implies that $X \times Y$ is finite. $\square$

Theorem 2.3.11. *Let $X_1, X_2, \dots, X_n$ be finite sets for some $n \in \mathbb{N}$. Then the cartesian product $\prod_{k=1}^n X_k$ is finite.*

Proof. Let $P(n) : \prod_{k=1}^n X_k$ is finite for some $n \in \mathbb{N}$, given that $X_1, X_2, \dots, X_n$ are finite sets.

- **Base case:** Let X_1 be a finite set. Then, $\prod_{k=1}^1 X_k = X_1$ is finite. Thus, $P(1)$ is true.

- **Inductive step:** Assume that $P(n)$ is true for some $n \in \mathbb{N}$.

Let $X_1, X_2, \dots, X_{n+1}$ be finite sets. Then we have

$$\prod_{k=1}^{n+1} X_k = \left(\prod_{k=1}^n X_k \right) \times X_{n+1}.$$

By the inductive hypothesis, $\prod_{k=1}^n X_k$ is finite. Then, by [Lemma 2.3.10](#), $\prod_{k=1}^{n+1} X_k$ is finite. Thus $P(n+1)$ is true whenever $P(n)$ is true.

By the principle of mathematical induction, $P(n)$ is true for all $n \in \mathbb{N}$. $\square$

Theorem 2.3.12. *Let X be a nonempty set. Then there exists no surjection $f : X \rightarrow \mathcal{P}(X)$, where*

$$\mathcal{P}(X) = \{E \mid E \subseteq X\}$$

*is the **power set** of X . In particular, if X is finite, then $|\mathcal{P}(X)| > |X|$. This theorem is called **Cantor's theorem**.*

Proof. Let $S = \{x \in X \mid x \notin f(x)\}$. Then $S \subseteq X$, which implies that $S \in \mathcal{P}(X)$. Suppose that there exists a surjection $f : X \rightarrow \mathcal{P}(X)$. Then there exists $x \in X$ such that $f(x) = S$.

If $x \in S$, then $x \notin f(x) = S$, which is a contradiction. If $x \notin S$, then $x \in f(x) = S$, which is also a contradiction. Therefore, there exists no surjection $f : X \rightarrow \mathcal{P}(X)$.

If X is finite, by [Theorem 2.3.6](#), there exists no surjection $f : X \rightarrow \mathcal{P}(X)$ if and only if $|X| < |\mathcal{P}(X)|$. $\square$

2.4 Countable and Uncountable Sets

A set X is said to be **countable** if $\mathbb{N} \sim X$. A set X is said to be **uncountable** if it is neither finite nor countable. A set X is said to be **at most countable** if X is either finite or countable.

Theorem 2.4.1. *Let X be an infinite set and Y be a finite set. Then $X \setminus Y$ is infinite.*

Proof. Suppose $X \setminus Y$ is finite. Since Y is finite, by [Theorem 2.3.8](#), $(X \setminus Y) \cup Y$ is finite. Then, by [Theorem 2.3.2](#), $X \subseteq (X \setminus Y) \cup Y$ is finite, which contradicts the assumption. Thus, $X \setminus Y$ is infinite. $\square$

Theorem 2.4.2. *Let X be an infinite set. If $X \sim Y$, then Y is infinite.*

Proof. Suppose Y is finite. Then there exists a bijection $f : J_n \rightarrow Y$ for some $n \in \mathbb{N}$. Since $X \sim Y$, there exists a bijection $g : X \rightarrow Y$.

Then the composition $g^{-1} \circ f : J_n \rightarrow X$ is a bijection, which implies that X is finite. This contradicts the assumption that X is infinite. Thus, Y is infinite. $\square$

Lemma 2.4.3. *$\mathbb{N}$ is infinite.*

Proof. Suppose $\mathbb{N}$ is finite. Then there exists a bijection $f : J_n \rightarrow \mathbb{N}$ for some $n \in \mathbb{N}$. Then, by [Theorem 2.3.2](#), $f(J_n)$ is finite. By [Theorem 2.3.3](#), $f(J_n)$ has a greatest element, say m .

Then we have $m+1 \in \mathbb{N}$ and $m < m+1 \notin f(J_n)$, which contradicts the assumption that f is surjective. Thus, $\mathbb{N}$ is infinite. $\square$

Theorem 2.4.4. *Every countable set is infinite.*

Proof. Let X be a countable set. Then there exists a bijection $f : \mathbb{N} \rightarrow X$. By Theorem 2.4.2 and lemma 2.4.3, X is infinite. $\square$

Remark. Let X be a countable set. Then there exists a bijection $f : \mathbb{N} \rightarrow X$. The elements of X can be listed as $X = \{x_1, x_2, x_3, \dots\}$, where $x_n = f(n)$ for all $n \in \mathbb{N}$. In other words, the elements of X can be arranged in a sequence.

Theorem 2.4.5. $\mathbb{Z}$ is countable.

Proof. Let $f : \mathbb{N} \rightarrow \mathbb{Z}$ be defined by

$$f(n) = \begin{cases} -\frac{n}{2}, & \text{if } n \text{ is even} \\ \frac{n-1}{2}, & \text{if } n \text{ is odd} \end{cases}.$$

We will show that f is bijective, which implies that $\mathbb{Z}$ is countable.

- **Injectivity:** Suppose $f(n_1) = f(n_2)$ for some $n_1, n_2 \in \mathbb{N}$. Since $f(n) \geq 0$ for odd n and $f(n) < 0$ for even n , we have n_1 and n_2 are either both even or both odd.

- If both n_1 and n_2 are even, then $-\frac{n_1}{2} = -\frac{n_2}{2}$, which implies $n_1 = n_2$.
- If both n_1 and n_2 are odd, then $\frac{n_1-1}{2} = \frac{n_2-1}{2}$, which implies $n_1 = n_2$.

Thus, f is injective.

- **Surjectivity:** Let $k \in \mathbb{Z}$. We need to find an $n \in \mathbb{N}$ such that $f(n) = k$.

- If $k \geq 0$, let $n = 2k + 1$. Then $f(n) = \frac{n-1}{2} = \frac{2k}{2} = k$.
- If $k < 0$, let $n = -2k$. Then $f(n) = -\frac{n}{2} = -\frac{-2k}{2} = k$.

Thus, f is surjective.

Since f is both injective and surjective, we have f is bijective. $\square$

Lemma 2.4.6. Every infinite subset of $\mathbb{N}$ is countable.

Proof. Let $X \subseteq \mathbb{N}$ be an infinite subset. Let $X_1 := X$ and define a function $f : \mathbb{N} \rightarrow X$ as follows:

- $f(n) = \min X_n$

$$\bullet X_{n+1} := X_n \setminus \{f(n)\}$$

for each $n \in \mathbb{N}$. By [Theorem 2.4.1](#), X_n is infinite for each $n \in \mathbb{N}$, so is nonempty. Then, by [Theorem 1.1.1](#), X_n has a least element, so f is well-defined. We will show that f is bijective, which implies that X is countable.

- **Injectivity:** Suppose $n_1 \neq n_2$ for some $n_1, n_2 \in \mathbb{N}$. Without loss of generality, assume that $n_1 < n_2$. Then $f(n_1) \in X_{n_1}$ and $f(n_2) \in X_{n_2}$. Since $X_{n_2} \subseteq X_{n_1} \setminus \{f(n_1)\}$, we have $f(n_2) \neq f(n_1)$. Thus, f is injective.
- **Surjectivity:** Let $x \in X$. Let $S := \{n \in \mathbb{N} \mid n < x\}$. If $S = \emptyset$, then $f(1) = x$. Otherwise, by [Lemma 2.3.1](#), $S \subseteq \{1, 2, \dots, x-1\}$ is finite. Let $k = |S|$, $S = \{s_1, s_2, \dots, s_k\}$ with $s_1 < s_2 < \dots < s_k$. Then we have

$$\begin{aligned} - \min X_1 &= \min S \implies X_2 = X \setminus \{s_1\}. \\ - \min X_2 &= \min S \setminus \{s_1\} \implies X_3 = X \setminus \{s_1, s_2\}. \\ &\vdots \\ - \min X_k &= \min S \setminus \{s_1, s_2, \dots, s_{k-1}\} \implies X_{k+1} = X \setminus S. \end{aligned}$$

Thus, $f(k+1) = \min X_{k+1} = x$. Therefore, f is surjective.

□

Theorem 2.4.7. *Every infinite subset of a countable set is countable.*

Proof. Let X be a countable set and let $Y \subseteq X$ be an infinite subset. Since X is countable, there exists a bijection $f : \mathbb{N} \rightarrow X$. By [Theorem 2.2.5](#), $f|_{f^{-1}(Y)} : f^{-1}(Y) \rightarrow Y$ is bijective. Then, by [Theorem 2.4.2](#), $f^{-1}(Y) \subseteq \mathbb{N}$ is infinite.

By [Lemma 2.4.6](#), $f^{-1}(Y)$ is countable. Then there exists a bijection $g : \mathbb{N} \rightarrow f^{-1}(Y)$. The composition $f|_{f^{-1}(Y)} \circ g : \mathbb{N} \rightarrow Y$ is bijective, which implies that Y is countable. □

Theorem 2.4.8. *Let X be infinite. If there exists a surjection $f : \mathbb{N} \rightarrow X$, then X is countable.*

Proof. For each $k \in X$, let $S_k := f^{-1}(\{k\})$. Since f is surjective, $S_k \neq \emptyset$ and by [Theorem 1.1.1](#), S_k has a least element for all $k \in X$. Let $g : X \rightarrow \mathbb{N}$ be defined by

$$g(k) = \min S_k \text{ for all } k \in X.$$

Suppose $g(k_1) = g(k_2)$ for some $k_1, k_2 \in X \implies \min S_{k_1} = \min S_{k_2}$. Since $S_i \cap S_j = \emptyset$ for all $i, j \in X$ with $i \neq j$, we have $S_{k_1} = S_{k_2} \implies k_1 = k_2$. Thus,

g is injective.

Since g is injective, by [Theorem 2.2.5](#), $g|^{g(X)} : X \rightarrow g(X)$ is bijective. Since X is infinite, $g(X) \subseteq \mathbb{N}$ is infinite by [Theorem 2.4.2](#). Then, by [Lemma 2.4.6](#), $g(X)$ is countable, which implies that X is countable. $\square$

Theorem 2.4.9. *Let $\{X_n\}_{n \in \mathbb{N}}$ be a sequence of countable sets. Then $\bigcup_{n=1}^{\infty} X_n$ is countable, i.e., any countable union of countable sets is countable.*

Proof. Since each X_n is countable, there exists a bijection $f_n : \mathbb{N} \rightarrow X_n$ for each $n \in \mathbb{N}$. For each $n \in \mathbb{N}$, we define the followings:

- $t_n = \min T_n$, where $T_n := \left\{ k \in \mathbb{N} \mid n \leq \frac{k(k+1)}{2} \right\}$.
- $s_n = n - \frac{t_n(t_n - 1)}{2} = n - \frac{t_n(t_n + 1)}{2} + t_n \in \mathbb{N}$.

Since $2n \in T_n$, $T_n \neq \emptyset$. Then, by [Theorem 1.1.1](#), T_n has a least element. Thus, t_n is well-defined. Since $\frac{t_n(t_n - 1)}{2} < n \leq \frac{t_n(t_n + 1)}{2}$, we have

$$t_n - s_n + 1 = \frac{t_n(t_n + 1)}{2} - n + 1 \geq 1$$

so $t_n - s_n + 1 \in \mathbb{N}$. Let $g : \mathbb{N} \rightarrow \bigcup_{n=1}^{\infty} X_n$ be defined by

$$g(k) = f_{s_k}(t_k - s_k + 1) \text{ for all } k \in \mathbb{N}.$$

Let $x \in \bigcup_{n=1}^{\infty} X_n$. Then there exists an $m \in \mathbb{N}$ such that $x \in X_m$. Since f_m is surjective, there exists a $l \in \mathbb{N}$ such that $f_m(l) = x$. Let

$$k = \frac{(l + m - 1)(l + m - 2)}{2} + m \in \mathbb{N}.$$

We will show that $l + m - 2 \notin T_k$ and $l + m - 1 \in T_k$, which implies that

- $t_k = \min T_k = l + m - 1$.
- $s_k = k - \frac{t_k(t_k - 1)}{2} = m$.
- $g(k) = f_{s_k}(t_k - s_k + 1) = f_m(l + m - 1 - m + 1) = f_m(l) = x$.

Consider the following inequalities when $z = l + m - 2$ and $z = l + m - 1$:

- Let $z = l + m - 2$. Then we have

$$\begin{aligned} k - \frac{z(z+1)}{2} &= k - \frac{(l+m-2)(l+m-1)}{2} \\ &= \frac{(l+m-1)(l+m-2)}{2} + m - \frac{(l+m-2)(l+m-1)}{2} \\ &= m > 0. \end{aligned}$$

- Let $z = l + m - 1$. Then we have

$$\begin{aligned} k - \frac{z(z+1)}{2} &= k - \frac{(l+m-1)(l+m)}{2} \\ &= \frac{(l+m-1)(l+m-2)}{2} + m - \frac{(l+m-1)(l+m)}{2} \\ &= m - \frac{(l+m-1)(l+m) - (l+m-1)(l+m-2)}{2} \\ &= m - (l+m-1) = 1-l \leq 0. \end{aligned}$$

Thus g is surjective, which implies that $\bigcup_{n=1}^{\infty} X_n$ is countable by [Theorem 2.4.8](#). $\square$

Corollary 2.4.10. *Let $X_1, X_2, \dots, X_n$ be countable sets for some $n \in \mathbb{N}$. Then $\bigcup_{i=1}^n X_i$ is countable, i.e., any finite union of countable sets is countable.*

Proof. Let $X_{n+1}, X_{n+2}, \dots$ be countable sets. Then, by [Theorem 2.4.9](#), $\bigcup_{n=1}^{\infty} X_n$ is countable. Suppose that $\bigcup_{i=1}^n X_i$ is finite. Then $X_1 \subseteq \bigcup_{i=1}^n X_i$ is finite by [Theorem 2.3.2](#), which contradicts [Theorem 2.4.4](#). Thus, $\bigcup_{i=1}^n X_i$ is infinite. Since $\bigcup_{i=1}^n X_i \subseteq \bigcup_{n=1}^{\infty} X_n$, by [Theorem 2.4.7](#), $\bigcup_{i=1}^n X_i$ is countable. $\square$

Theorem 2.4.11. *Suppose that X is at most countable, and, for each $x \in X$, Y_x is at most countable. Then the union*

$$Y = \bigcup_{x \in X} Y_x$$

is at most countable, i.e., any at most countable union of at most countable sets is at most countable.

Proof. We have 4 cases to consider:

- (A) Finite union of finite sets is finite by [Theorem 2.3.9](#).
- (B) Finite union of countable sets is countable by [Corollary 2.4.10](#).
- (C) Countable union of finite sets is countable.
- (D) Countable union of countable sets is countable by [Theorem 2.4.9](#).

Let us prove (C). Let X be a countable set and Y_x be finite for each $x \in X$. Then, there exists a bijection $f : \mathbb{N} \rightarrow X$, and for each $x \in X$, there exists a bijection $g_x : J_{n_x} \rightarrow Y_x$ for some $n_x \in \mathbb{N}$. Let $h : \mathbb{N} \rightarrow Y$ be defined by

- $h(j) = g_{f(1)}(j)$ for $j = 1, 2, \dots, n_{f(1)}$.
- $h\left(\sum_{i=1}^{k-1} n_{f(i)} + m\right) = g_{f(k)}(m)$ for each $k \in \mathbb{N}$ with $k \geq 2$, and $m = 1, 2, \dots, n_{f(k)}$.

Then, h is a surjection from $\mathbb{N}$ to Y . By [Theorem 2.4.8](#), Y is countable. $\square$

Lemma 2.4.12. $\mathbb{N} \times \mathbb{N}$ is countable.

Proof. We will explicitly construct a bijection from $\mathbb{N} \times \mathbb{N} \rightarrow \mathbb{N}$. Let $f : \mathbb{N} \times \mathbb{N} \rightarrow \mathbb{N}$ be defined by

$$f(m, n) = \frac{(m+n-1)(m+n-2)}{2} + m$$

for each $(m, n) \in \mathbb{N} \times \mathbb{N}$. We will show that f is bijective.

- **Injectivity:** Suppose $(m_1, n_1) \neq (m_2, n_2)$. If $m_1 + n_1 = m_2 + n_2$, then $m_1 \neq m_2$, so $f(m_1, n_1) \neq f(m_2, n_2)$. Otherwise, $m_1 + n_1 \neq m_2 + n_2$. Without loss of generality, assume $m_1 + n_1 < m_2 + n_2$. Then, since $m_1 + n_1 + 1 \leq m_2 + n_2$, we have

$$\begin{aligned} f(m_1, n_1) &= \frac{(m_1 + n_1 - 1)(m_1 + n_1 - 2)}{2} + m_1 \\ &\leq \frac{(m_1 + n_1 - 1)(m_1 + n_1 - 2)}{2} + (m_1 + n_1 - 1) \\ &= \frac{(m_1 + n_1)(m_1 + n_1 - 1)}{2} \\ &\leq \frac{(m_2 + n_2 - 1)(m_2 + n_2 - 2)}{2} \\ &< \frac{(m_2 + n_2 - 1)(m_2 + n_2 - 2)}{2} + m_2 = f(m_2, n_2), \end{aligned}$$

Thus, $f(m_1, n_1) \neq f(m_2, n_2)$, so f is injective.

- **Surjectivity:** Let $k \in \mathbb{N}$. Let t_k, s_k be defined as in the proof of [Theorem 2.4.9](#). Then we have $f(s_k, t_k - s_k + 1) = k$. Thus, f is surjective.

□

Theorem 2.4.13. *Let X and Y be countable sets. Then $X \times Y$ is countable.*

Proof. There exist bijections $f : \mathbb{N} \rightarrow X$ and $g : \mathbb{N} \rightarrow Y$. Let $h : \mathbb{N} \times \mathbb{N} \rightarrow X \times Y$ be defined by

$$h(m, n) = (f(m), g(n)) \text{ for all } (m, n) \in \mathbb{N} \times \mathbb{N}.$$

Then, h is a bijection from $\mathbb{N} \times \mathbb{N}$ to $X \times Y$. By [Lemma 2.4.12](#), $\mathbb{N} \times \mathbb{N}$ is countable, so $X \times Y$ is countable. □

Theorem 2.4.14. *Let $X_1, X_2, \dots, X_n$ be countable sets for some $n \in \mathbb{N}$. Then $\prod_{i=1}^n X_i$ is countable, i.e., any finite product of countable sets is countable.*

Proof. We will prove the theorem by mathematical induction on n . Let $P(n) : \prod_{i=1}^n X_i$ is countable for some $n \in \mathbb{N}$, given that $X_1, X_2, \dots, X_n$ are countable sets.

- **Base case:** Let X_1 be a countable set. Then $\prod_{i=1}^1 X_i = X_1$ is countable. Thus, $P(1)$ is true.

- **Inductive step:** Assume that $P(n)$ is true for some $n \in \mathbb{N}$.

Let $X_1, X_2, \dots, X_{n+1}$ be countable sets. Then we have

$$\prod_{i=1}^{n+1} X_i = \left(\prod_{i=1}^n X_i \right) \times X_{n+1}$$

By the inductive hypothesis, $\prod_{i=1}^n X_i$ is countable. Then, by [Theorem 2.4.13](#),

$\prod_{i=1}^{n+1} X_i$ is countable. Thus, $P(n+1)$ is true when $P(n)$ is true.

By the principle of mathematical induction, $P(n)$ is true for all $n \in \mathbb{N}$. □

Theorem 2.4.15. *$\mathbb{Q}$ is countable.*

Proof. Any $q \in \mathbb{Q}$ can be expressed as $q = \frac{m}{n}$ for some $m \in \mathbb{Z}$ and $n \in \mathbb{N}$. Thus, we have

$$f : \mathbb{Z} \times \mathbb{N} \rightarrow \mathbb{Q}, \quad f(m, n) = \frac{m}{n} \text{ for all } (m, n) \in \mathbb{Z} \times \mathbb{N}.$$

is a surjection. By [Theorems 2.4.5](#) and [2.4.13](#), $\mathbb{Z} \times \mathbb{N}$ is countable. Then, there exists a bijection $g : \mathbb{N} \rightarrow \mathbb{Z} \times \mathbb{N}$. Let $h \equiv f \circ g : \mathbb{N} \rightarrow \mathbb{Q}$. By [Theorem 2.2.7](#), h is a surjection.

Since $\mathbb{N} \subseteq \mathbb{Q}$, by [Theorem 2.3.2](#) and [lemma 2.4.3](#), $\mathbb{Q}$ is infinite. Then, by [Theorem 2.4.8](#), $\mathbb{Q}$ is countable. $\square$

Theorem 2.4.16. Let $A := \{0, 1\}$. Then $X = \prod_{n=1}^{\infty} A$ is uncountable.

Proof. For each $n \in \mathbb{N}$, let $e_n \in X$ be the sequence whose n -th term is 1 and all other terms are 0. Let $f : \mathbb{N} \rightarrow X$ be defined by $f(n) = e_n$ for all $n \in \mathbb{N}$. Since $f|_{f(\mathbb{N})}$ is bijective, $f(\mathbb{N}) \subseteq X$ is countable, so is infinite by [Theorem 2.4.4](#).

Then, by [Theorem 2.3.2](#), X is infinite. Suppose that X is countable. Then, there exists a bijection $g : \mathbb{N} \rightarrow X$. Let $X = \{g(1), g(2), g(3), \dots\}$, where

- $g(1) = (a_{11}, a_{12}, a_{13}, \dots)$.
- $g(2) = (a_{21}, a_{22}, a_{23}, \dots)$.
- $\vdots$

Let $x = (b_1, b_2, b_3, \dots) \in X$ be defined by $b_n = 1 - a_{nn}$ for all $n \in \mathbb{N}$. Then, $x \neq g(n)$ for all $n \in \mathbb{N}$, which contradicts the assumption that g is surjective. Thus, X is uncountable. $\square$

Theorem 2.4.17. $\mathbb{R}$ is uncountable.

Proof. Since $\mathbb{N} \subseteq \mathbb{R}$, $\mathbb{R}$ is infinite, as like [Q is infinite](#). Let $f : \mathbb{N} \rightarrow [0, 1)$ be defined by $f(n) = \frac{1}{n+1}$ for all $n \in \mathbb{N}$. $[0, 1) \subseteq \mathbb{R}$ is infinite as like [X is infinite](#). We will prove that $[0, 1) \subseteq \mathbb{R}$ is uncountable. Then, by [Theorem 2.4.7](#), $\mathbb{R}$ is uncountable.

Let $g : [0, 1) \rightarrow \prod_{n=1}^{\infty} \{0, 1\}$ be defined by $g(x) = \{a_n^x\}_{n=1}^{\infty}$ for all $x \in [0, 1)$, where

$\{a_n^x\}_{n=1}^{\infty}$ is the greedy base 2 expansion of x . Let $Y := g([0, 1)) \subseteq \prod_{n=1}^{\infty} \{0, 1\}$.

Then, by [Theorems 1.3.9](#) and [2.2.5](#), $g|_Y$ is bijective.

Suppose that $[0, 1)$ is countable. Then there exists a bijection $h : \mathbb{N} \rightarrow [0, 1)$. By [Theorem 2.2.7](#), $g|_Y \circ h : \mathbb{N} \rightarrow Y$ is bijective, so $Y \subseteq \prod_{n=1}^{\infty} \{0, 1\}$ is countable. Let

$Z := \prod_{n=1}^{\infty} \{0, 1\} \setminus Y$. By [Theorems 1.3.10](#) and [1.3.11](#), we have

$$Z = \left\{ \{a_n\}_{n=1}^{\infty} \in \prod_{n=1}^{\infty} \{0, 1\} \mid a_n = 1 \text{ for all } n \geq N \text{ for some } N \in \mathbb{N} \right\}.$$

Let Z_N be defined by

$$Z_N = \left\{ \{a_n\}_{n=1}^{\infty} \in \prod_{n=1}^{\infty} \{0, 1\} \mid a_n = 1 \text{ for all } n \geq N \right\} \text{ for each } N \in \mathbb{N}.$$

Let $F_N : \prod_{n=1}^{N-1} \{0, 1\} \rightarrow Z_N$ be defined by

$$F_N \left(\{a_n\}_{n=1}^{N-1} \right) = \{a_n\}_{n=1}^{N-1} \cup \{1\}_{n=N}^{\infty} \text{ for all } \{a_n\}_{n=1}^{N-1} \in \prod_{n=1}^{N-1} \{0, 1\}.$$

Then F_N is bijective, so Z_N is finite by [Theorem 2.3.11](#). By [Theorem 2.4.11](#), $Z = \bigcup_{N=1}^{\infty} Z_N$ is countable. Then, by [Corollary 2.4.10](#), $\prod_{n=1}^{\infty} \{0, 1\} = Y \cup Z$ is countable, which contradicts [Theorem 2.4.16](#). Thus, $[0, 1)$ is uncountable. $\square$

2.5 Metric Spaces

Let X be a set. A function $d : X \times X \rightarrow \mathbb{R}$ is called a **metric** on X if for any $p, q \in X$, we have

(A) $d(p, q) \geq 0$ and $d(p, q) = 0 \iff p = q$.

(B) $d(p, q) = d(q, p)$.

(C) $d(p, r) \leq d(p, q) + d(q, r)$ for any $r \in X$.

(C) is called the **triangle inequality**. The pair (X, d) is called a **metric space**.

Theorem 2.5.1. We define a function $d : \mathbb{R}^n \times \mathbb{R}^n \rightarrow \mathbb{R}$ by

$$d(\mathbf{x}, \mathbf{y}) = \|\mathbf{x} - \mathbf{y}\| = \sqrt{\sum_{k=1}^n (x_k - y_k)^2}$$

for all $\mathbf{x}, \mathbf{y} \in \mathbb{R}^n$. Then d is a metric on $\mathbb{R}^n$. Especially, d is called the **standard metric** on $\mathbb{R}^n$.

Proof. Let $\mathbf{x}, \mathbf{y}, \mathbf{z} \in \mathbb{R}^n$ be given.

- $d(\mathbf{x}, \mathbf{y}) \geq 0$ is obvious. Also, we have

$$\begin{aligned} d(\mathbf{x}, \mathbf{y}) = 0 &\iff \|\mathbf{x} - \mathbf{y}\| = 0 \\ &\iff x_k = y_k \text{ for all } k = 1, 2, \dots, n \\ &\iff \mathbf{x} = \mathbf{y}. \end{aligned}$$

- $d(\mathbf{x}, \mathbf{y}) = d(\mathbf{y}, \mathbf{x})$ is obvious.
- By [Theorem 1.6.2](#), we have

$$\begin{aligned} d(\mathbf{x}, \mathbf{z}) &= \|\mathbf{x} - \mathbf{z}\| = \|\mathbf{x} - \mathbf{y} + (\mathbf{y} - \mathbf{z})\| \\ &\leq \|\mathbf{x} - \mathbf{y}\| + \|\mathbf{y} - \mathbf{z}\| = d(\mathbf{x}, \mathbf{y}) + d(\mathbf{y}, \mathbf{z}) \end{aligned}$$

Therefore, d is a metric on $\mathbb{R}^n$. □

Definition 2.5.2. Let (X, d) be a metric space. For any $p \in X$ and $r > 0$, we define the **neighborhood** of p with radius r as

$$N_r(p) := \{q \in X \mid d(p, q) < r\}.$$

Especially, if $X = \mathbb{R}^n$ and d is the standard metric on $\mathbb{R}^n$, then $N_r(p)$ is called the **open ball** of radius r centered at p , which is denoted by $B_r(p)$.

Definition 2.5.3. Let (X, d) be a metric space and $E \subseteq X$. Let $p \in X$.

- p is called an **limit point** of E if for any $r > 0$, we have $E \cap (N_r(p) \setminus \{p\}) \neq \emptyset$.
- p is called a **isolated point** of E if there exists some $r > 0$ such that $N_r(p) \cap E = \{p\}$.
- p is called an **interior point** of E if there exists some $r > 0$ such that $N_r(p) \subseteq E$.

Definition 2.5.4. Let (X, d) be a metric space and $E \subseteq X$.

- E is called **open** if every point of E is an interior point of E .
- E is called **closed** if every limit point of E belongs to E .

Theorem 2.5.5. Let (X, d) be a metric space. For any $p \in X$ and $r > 0$, the neighborhood $N_r(p)$ is open.

Proof. Let $q \in N_r(p)$ be given. Then we have $d(p, q) < r$. Put $\varepsilon = r - d(p, q) > 0$. We will show that $N_\varepsilon(q) \subseteq N_r(p)$. Let $s \in N_\varepsilon(q)$ be given. Then we have $d(q, s) < \varepsilon$. By the *triangle inequality*, we have

$$d(p, s) \leq d(p, q) + d(q, s) < d(p, q) + \varepsilon = r.$$

Therefore, we have $s \in N_r(p)$, which implies that $N_\varepsilon(q) \subseteq N_r(p)$. Since q is an arbitrary point of $N_r(p)$, every point of $N_r(p)$ is an interior point of $N_r(p)$. Thus, we conclude that $N_r(p)$ is open. $\square$

2.6 Compact Sets

2.7 Perfect Sets

2.8 Connected Sets

Chapter 3

Numerical Sequences and Series

3.1 Convergent Sequences

3.2 Subsequences

3.3 Cauchy Sequences

3.4 Upper and Lower Limits

3.5 Some Special Sequences

3.6 Series

3.7 Series of Nonnegative Terms

3.8 The Number e

3.9 The Root and Ratio Tests

3.10 Power Series

3.11 Summation by Parts

3.12 Absolute Convergence

3.13 Addition and Multiplication of Series

3.14 Rearrangements

Chapter 4

Continuity

4.1 Limits of Functions

4.2 Continuous Functions

4.3 Continuity and Compactness

4.4 Continuity and Connectedness

4.5 Discontinuities

4.6 Monotonic Functions

4.7 Infinite Limits and Limits at Infinity

Chapter 5

Differentiation

5.1 The Derivative of a Real Function

5.2 Mean Value Theorems

5.3 The Continuity of Derivatives

5.4 L'Hospital's Rule

5.5 Derivatives of Higher Order

5.6 Taylor's Theorem

5.7 Differentiation of Vector-valued Functions

Chapter 6

The Riemann-Stieltjes Integral

6.1 Definition and Existence of the Integral

6.2 Properties of the Integral

6.3 Integration and Differentiation

6.4 Integration of Vector-valued Functions

6.5 Rectifiable Curves

Chapter 7

Sequences and Series of Functions

7.1 Discussion of Main Problem

7.2 Uniform Convergence

7.3 Uniform Convergence and Continuity

7.4 Uniform Convergence and Integration

7.5 Uniform Convergence and Differentiation

7.6 Equicontinuous Families of Functions

7.7 Equicontinuous Families of Functions

7.8 The Stone-Weierstrass Theorem

Chapter 8

Some Special Functions

8.1 Power Series

8.2 The Exponential and Logarithmic Functions

8.3 The Trigonometric Functions

8.4 The Algebraic Completeness of the Complex Field

8.5 Fourier Series

8.6 The Gamma Function

Chapter 9

Functions of Several Variables

9.1 Linear Transformations

9.2 Differentiation

9.3 The Contraction Principle

9.4 The Inverse Function Theorem

9.5 The Implicit Function Theorem

9.6 The Rank Theorem

9.7 Determinants

9.8 Derivatives of Higher Order

9.9 Differentiation of Integrals

Chapter 10

Integration of Differential Forms

10.1 Integration

10.2 Primitive Mappings

10.3 Partitions of Unity

10.4 Change of Variables

10.5 Differential Forms

10.6 Simplexes and Chains

10.7 Stokes' Theorem

10.8 Closed Forms and Exact Forms

10.9 Vector Analysis

Chapter 11

The Lebesgue Theory

11.1 Set Functions

11.2 Construction of the Lebesgue Measure

11.3 Measure Spaces

11.4 Measurable Functions

11.5 Simple Functions

11.6 Integration

11.7 Comparison with the Riemann Integral

11.8 Integration of Complex Functions

11.9 Functions of Class $\mathcal{L}^2$

Bibliography

- [1] Walter Rudin, *Principles of Mathematical Analysis*, McGraw-Hill International Editions.
- [2] Kenneth Ross, *Elementary Analysis*, Springer Nature.
- [3] Steven Douglass, *Introduction to Mathematical Analysis*, Pearson.
- [4] Terence Tao, *Analysis I*, Springer Nature.