

# **Mathematical Analysis I**

# Mathematical Analysis I

## Author

MUHO CHOI

Email: [sihyeon8240@gmail.com](mailto:sihyeon8240@gmail.com)

GitHub: <https://github.com/sihyeon8240>

## Source Code & Contributions

GitHub: <https://github.com/sihyeon8240/math>

Feel free to open issues or pull requests for corrections.

© 2026. MUHO CHOI.

This textbook is licensed under a [Creative Commons Attribution 4.0 International License](https://creativecommons.org/licenses/by/4.0/) (CC BY 4.0).

**You are free to:** Share, copy, and redistribute the material in any medium or format; adapt, remix, transform, and build upon the material for any purpose, even commercially.

**Under the following terms:** You must give appropriate credit, provide a link to the license, and indicate if changes were made.

# Contents

|                                                      |           |
|------------------------------------------------------|-----------|
| <b>Preface</b>                                       | <b>iv</b> |
| <b>1 Basic Topology</b>                              | <b>1</b>  |
| 1.1 Metric Spaces . . . . .                          | 1         |
| 1.2 Compact Sets . . . . .                           | 23        |
| 1.3 Connected Sets . . . . .                         | 43        |
| <b>2 Sequences and Series</b>                        | <b>50</b> |
| 2.1 Convergent Sequences . . . . .                   | 50        |
| 2.2 Subsequences . . . . .                           | 58        |
| 2.3 Cauchy Sequences . . . . .                       | 58        |
| 2.4 Upper and Lower Limits . . . . .                 | 58        |
| 2.5 Some Special Sequences . . . . .                 | 58        |
| 2.6 Series . . . . .                                 | 58        |
| 2.7 Series of Nonnegative Terms . . . . .            | 58        |
| 2.8 The Number $e$ . . . . .                         | 58        |
| 2.9 The Root and Ratio Tests . . . . .               | 58        |
| 2.10 Power Series . . . . .                          | 58        |
| 2.11 Summation by Parts . . . . .                    | 58        |
| 2.12 Absolute Convergence . . . . .                  | 58        |
| 2.13 Addition and Multiplication of Series . . . . . | 58        |
| 2.14 Rearrangements . . . . .                        | 58        |
| <b>3 Continuity</b>                                  | <b>59</b> |
| 3.1 Limits of Functions . . . . .                    | 59        |
| 3.2 Continuous Functions . . . . .                   | 59        |
| 3.3 Continuity and Compactness . . . . .             | 59        |
| 3.4 Continuity and Connectedness . . . . .           | 59        |
| 3.5 Discontinuities . . . . .                        | 59        |
| 3.6 Monotonic Functions . . . . .                    | 59        |
| 3.7 Infinite Limits and Limits at Infinity . . . . . | 59        |

|          |                                                      |           |
|----------|------------------------------------------------------|-----------|
| <b>4</b> | <b>Differentiation</b>                               | <b>60</b> |
| 4.1      | The Derivative of a Real Function . . . . .          | 60        |
| 4.2      | Mean Value Theorems . . . . .                        | 60        |
| 4.3      | The Continuity of Derivatives . . . . .              | 60        |
| 4.4      | L'Hospital's Rule . . . . .                          | 60        |
| 4.5      | Derivatives of Higher Order . . . . .                | 60        |
| 4.6      | Taylor's Theorem . . . . .                           | 60        |
| 4.7      | Differentiation of Vector-valued Functions . . . . . | 60        |
| <b>5</b> | <b>The Riemann-Stieltjes Integral</b>                | <b>61</b> |
| 5.1      | Definition and Existence of the Integral . . . . .   | 61        |
| 5.2      | Properties of the Integral . . . . .                 | 61        |
| 5.3      | Integration and Differentiation . . . . .            | 61        |
| 5.4      | Integration of Vector-valued Functions . . . . .     | 61        |
| 5.5      | Rectifiable Curves . . . . .                         | 61        |

# Preface

This textbook uses Lean 4 to formalize selected mathematical statements and proofs.

- **Tools.** Lean is an interactive theorem prover; Mathlib is a library of formalized mathematics. This edition uses Lean 4.33.1 and Mathlib v4.33.1 (revision [0df444a360ea](#)).
- **Reading the proofs.** Each paired statement or proof step gives a mathematical explanation followed by its Lean code. The explanations can be read independently; code fragments are read in order within the complete declaration.
- **Verification.** Lean checks formal proofs under their stated assumptions. Verification covers only statements linked to checked declarations. Agreement between the explanations, displayed code, and checked proofs is reviewed separately.
- **Prerequisites.** Starting assumptions and accepted external results are specific to this textbook.

# Chapter 1

## Basic Topology

### 1.1 Metric Spaces

**Definition 1.1.1.** A **metric** on  $X$  is a function  $d : X \times X \rightarrow [0, \infty)$  such that, for all  $p, q, z \in X$ ,

$$d(p, q) = 0 \iff p = q, \quad d(p, q) = d(q, p), \quad d(p, z) \leq d(p, q) + d(q, z).$$

The last condition is the **triangle inequality**. The pair  $(X, d)$  is a **metric space**.

The metric-space structure records the distance and its axioms.

```
abbrev MetricStructure (A : Type*) := MetricSpace A
```

For every  $n \in \mathbb{N}$ ,  $\mathbb{R}^n$  has its Euclidean norm, including the unique vector when  $n = 0$ .

```
abbrev Euclidean (n : ℕ) := EuclideanSpace ℝ (Fin n)
```

#### Proposition 1.1.2.

For  $n \in \mathbb{N}$ , the function  $d(x, y) = \|x - y\|$  is a metric on  $\mathbb{R}^n$ .

```
theorem euclidean_metric (n : ℕ) :  
  (∀ x y : EuclideanSpace ℝ (Fin n), 0 ≤ ‖x - y‖) ∧  
  (∀ x y : EuclideanSpace ℝ (Fin n), ‖x - y‖ = 0 ↔ x = y) ∧  
  (∀ x y : EuclideanSpace ℝ (Fin n), ‖x - y‖ = ‖y - x‖) ∧  
  (∀ x y z : EuclideanSpace ℝ (Fin n),  
    ‖x - z‖ ≤ ‖x - y‖ + ‖y - z‖)
```

*Proof.*

The norm is nonnegative. It vanishes exactly at the zero vector, so  $\|x - y\| = 0$  precisely when  $x = y$ .

```

refine (fun x y => norm_nonneg _, ?, ?, ?)
· intro x y
  rw [norm_eq_zero, sub_eq_zero]

```

Reversing a difference changes only its sign and leaves its norm unchanged.

```

· intro x y
  exact norm_sub_rev x y

```

Write  $x - z = (x - y) + (y - z)$  and apply the Euclidean norm triangle inequality. These norm facts are accepted finite-dimensional prerequisites.

```

· intro x y z
  calc
    ||x - z|| = ||(x - y) + (y - z)|| := by rw [sub_add_sub_cancel]
    _ ≤ ||x - y|| + ||y - z|| := norm_add_le _ _

```

□

Euclidean spaces always use this standard metric. In particular, the zero-dimensional space is included, rather than excluded by convention.

**Definition 1.1.3.** For  $p \in X$  and  $r > 0$ , define the **neighborhood**  $N_r(p) = \{q : d(p, q) < r\}$  and the **deleted neighborhood**  $\tilde{N}_r(p) = N_r(p) \setminus \{p\}$ . Metric symmetry allows the center to appear in either distance argument.

$N_r(p)$  is the open ball centered at  $p$ .

```

abbrev neighborhood (p : X) (r : ℝ) : Set X := ball p r

```

Removing the center gives  $\tilde{N}_r(p)$ .

```

abbrev deletedNeighborhood (p : X) (r : ℝ) : Set X := ball p r \ {p}

```

**Lemma 1.1.4.**

$q \in \tilde{N}_r(p)$  if and only if  $0 < d(q, p) < r$ .

```
theorem deleted_neighborhood_iff (p q : X) (r : ℝ) :
  q ∈ deletedNeighborhood p r ↔ 0 < dist q p ∧ dist q p < r
```

*Proof.*

The only additional condition after removing the center is  $q \neq p$ , which is equivalent to positive distance in a metric space.

```
change (dist q p < r ∧ q ≠ p) ↔ _
rw [dist_pos]
exact and_comm
```

□

**Definition 1.1.5.** A point  $p$  is a **limit point** of  $E$  if every deleted neighborhood of  $p$  meets  $E$ . It is an **interior point** if some neighborhood lies in  $E$ , and an **isolated point** if some neighborhood meets  $E$  exactly at  $p$ . We write these sets as  $E'$ ,  $E^\circ$ , and  $E^i$ , respectively.

$p \in E'$  means that every positive radius admits a distinct point of  $E$  within that distance.

```
def limitPoints (E : Set X) : Set X :=
  {p | ∀ r : ℝ, 0 < r → ∃ q ∈ E, q ≠ p ∧ dist q p < r}
```

$E^\circ$  is the usual interior; its ball description is verified below.

```
noncomputable abbrev interiorPoints (E : Set X) : Set X := interior E
```

$p \in E^i$  means  $E \cap N_r(p) = \{p\}$  for some  $r > 0$ . In particular  $p \in E$ .

```
def isolatedPoints (E : Set X) : Set X :=
  {p | ∃ r : ℝ, 0 < r ∧ E ∩ ball p r = {p}}
```

**Lemma 1.1.6.**

$p \in E^\circ$  if and only if some  $r > 0$  satisfies  $N_r(p) \subseteq E$ .

```
theorem interior_iff (E : Set X) (p : X) :
  p ∈ interiorPoints E ↔ ∃ r > 0, ball p r ⊆ E
```

*Proof.*

The interior consists of points for which  $E$  is a neighborhood. The metric neighborhood basis consists of positive-radius balls, giving the stated description.

```
exact mem_interior_iff_mem_nhds.trans Metric.mem_nhds_iff
```

□

**Definition 1.1.7.** A set  $E$  is **open** when every point of  $E$  is interior, and **closed** when  $E' \subseteq E$ . Write  $\mathcal{O}(X)$  and  $\mathcal{C}(X)$  for the respective collections. For openness, the metric ball characterization gives the standard predicate; the corresponding closed-set characterization is proved below.

$\mathcal{O}(X)$  is the collection of open sets.

```
abbrev openSets : Set (Set X) := {E | IsOpen E}
```

$\mathcal{C}(X)$  is the collection of closed sets.

```
abbrev closedSets : Set (Set X) := {E | IsClosed E}
```

**Definition 1.1.8.** The **closure** of  $E$  is  $\overline{E} = E \cup E'$ . We use the standard closure in Lean and prove this equality below.

$\overline{E}$  denotes the closure of  $E$ .

```
noncomputable abbrev setClosure (E : Set X) : Set X := closure E
```

**Definition 1.1.9.** A set  $E$  is **bounded** if, whenever  $X$  is nonempty,  $E$  is contained in some positive-radius ball. Write  $\mathcal{B}(X)$  for the bounded sets. This convention also treats subsets of the empty space as bounded. A set is **perfect** if  $E = E'$  and **dense** if  $\overline{E} = X$ .

When the ambient space has a point, a bounded set lies in a ball.

```
def bounded (E : Set X) : Prop :=
  Nonempty X → ∃ p : X, ∃ r > 0, E ⊆ ball p r
```

A perfect set equals its set of limit points.

```
abbrev perfect (E : Set X) : Prop := E = limitPoints E
```

A dense set has the whole ambient space as its closure.

```
abbrev dense (E : Set X) : Prop := closure E = univ
```

### Proposition 1.1.10.

*Every ball  $N_r(p)$  is open. This also holds for nonpositive  $r$ , when the ball is empty.*

```
theorem ball_open (p : X) (r : ℝ) : IsOpen (ball p r)
```

*Proof.*

Suppose  $q \in N_r(p)$ . The remaining distance  $\delta = r - d(q, p)$  is positive.

```
apply Metric.isOpen_iff.mpr
intro q hq
refine (r - dist q p, sub_pos.mpr hq, ?_)
```

If  $d(s, q) < \delta$ , the triangle inequality gives  $d(s, p) \leq d(s, q) + d(q, p) < r$ . Thus the ball of radius  $\delta$  about  $q$  stays inside  $N_r(p)$ .

```
intro s hs
change dist s p < r
have ht := dist_triangle s q p
change dist s q < r - dist q p at hs
linarith
```

□

**Lemma 1.1.11.**

For a finite index set  $S$  and positive numbers  $r_i$  for  $i \in S$ , there is  $\delta > 0$  with  $\delta \leq r_i$  for every  $i \in S$ .

```
theorem finite_positive_lower_bound {ι : Type*} (s : Finset ι)
  (r : ι → ℝ) (hr : ∀ i ∈ s, 0 < r i) :
  ∃ δ > 0, ∀ i ∈ s, δ ≤ r i
```

*Proof.*

Use induction on  $S$ . For the empty set,  $\delta = 1$  works. On adjoining  $i$ , take the smaller of  $r_i$  and the positive bound already obtained for the remaining indices. It stays positive and is below every required radius.

```
classical
induction s using Finset.induction_on with
| empty => exact ⟨1, zero_lt_one, by simp only [Finset.notMem_empty,
  ⇐ false_implies, implies_true]⟩
| @insert i s hi ih =>
  obtain ⟨δ, hδ, hle⟩ := ih (fun j hj => hr j (Finset.mem_insert_of_mem
    ⇐ hj))
  refine ⟨min (r i) δ, lt_min (hr i (Finset.mem_insert_self i s)) hδ,
    ⇐ ?_⟩
  intro j hj
  rcases Finset.mem_insert.mp hj with rfl | hj
  · exact min_le_left _ _
  · exact (min_le_right _ _).trans (hle j hj)
```

□

**Proposition 1.1.12.**

If  $p \in E'$  and  $r > 0$ , then  $E \cap N_r(p)$  is infinite.

```
theorem limit_point_infinite (E : Set X) (p : X)
  (hp : p ∈ limitPoints E) (r : ℝ) (hr : 0 < r) :
  (E ∩ ball p r).Infinite
```

*Proof.*

Assume instead that this intersection is finite. After deleting  $p$ , its points still form a finite set, and all their distances from  $p$  are positive.

```
classical
intro hf
let s := (hf.subset (show (E n ball p r) \ {p} ⊆ E n ball p r from
  ↪ sdiff_subset)).toFinset
have hpos : ∀ q ∈ s, 0 < dist q p := by
  intro q hq
  have h := (Set.Finite.mem_toFinset _).mpr hq
  exact dist_pos.mpr h.2
```

Choose a common positive lower bound  $\delta$  for those distances. The limit-point condition supplies a distinct point of  $E$  within  $\min(r, \delta)$  of  $p$ . It belongs to the finite set but has distance less than its lower bound, a contradiction.

```
obtain {δ, hδ, hle} := finite_positive_lower_bound s (fun q => dist q p)
  ↪ hpos
obtain {q, hq, hqp, hdist} := hp (min r δ) (lt_min hr hδ)
have hqs : q ∈ s := (Set.Finite.mem_toFinset _).mpr
  {⟨hq, hdist.trans_le (min_le_left _ _), hqp⟩}
exact (not_lt_of_ge (hle q hqs)) (hdist.trans_le (min_le_right _ _))
```

□

### Proposition 1.1.13.

$E$  is open if and only if  $E^c$  contains all its limit points, that is,  $E^c$  is closed.

```
theorem open_iff_compl_limitPoints (E : Set X) :
  IsOpen E ↔ limitPoints Ec ⊆ Ec
```

*Proof.*

If  $E$  is open, a point in  $E$  has a ball contained in  $E$ . Such a ball prevents that point from being a limit point of the complement.

```
classical
constructor
· intro h p hp hpe
  obtain ⟨r, hr, hball⟩ := Metric.isOpen_iff.mp h p hpe
  obtain ⟨q, hq, _, hd⟩ := hp r hr
  exact hq (hball hd)
```

Conversely, suppose the complement contains its limit points. If  $p \in E$  had no ball contained in  $E$ , each ball would contain a point outside  $E$ . That point differs from  $p$ , making  $p$  a limit point of the complement and contradicting  $p \in E$ .

```
· intro h
  apply Metric.isOpen_iff.mpr
  intro p hp
  by_contra hn
  have hl : p ∈ limitPoints Ec := by
    intro r hr
    have hnsb : ¬ ball p r ⊆ E := fun hs => hn ⟨r, hr, hs⟩
    obtain ⟨q, hq, hqe⟩ := Set.not_subset.mp hnsb
    refine ⟨q, hqe, ?_, hq⟩
    intro heq
    exact hqe (heq.symm ▸ hp)
  exact h hl hp
```

□

### Lemma 1.1.14.

The standard closed-set predicate is equivalent to  $E' \subseteq E$ .

```
theorem closed_iff_limitPoints (E : Set X) :
  IsClosed E ↔ limitPoints E ⊆ E
```

*Proof.*

A set is closed precisely when its complement is open. Apply the preceding result to  $E^c$  and take its complement twice.

```
have h := open_iff_compl_limitPoints Ec
rw [compl_compl] at h
exact isOpen_compl_iff.symm.trans h
```

□

### Proposition 1.1.15.

For any indexed family of open sets  $(E_i)$ , the union  $\bigcup_i E_i$  is open.

```
theorem open_union {ι : Type*} (E : ι → Set X)
  (h : ∀ i, IsOpen (E i)) : IsOpen (⋃ i, E i)
```

*Proof.*

A point in the union belongs to one member  $E_i$ . Choose a ball contained in that member; the same ball is contained in the union. This argument also covers an empty family, since then there is no point to consider.

```
apply Metric.isOpen_iff.mpr
intro p hp
obtain ⟨i, hi⟩ := mem_iUnion.mp hp
obtain ⟨r, hr, hs⟩ := Metric.isOpen_iff.mp (h i) p hi
exact ⟨r, hr, fun q hq => mem_iUnion.mpr ⟨i, hs hq⟩⟩
```

□

### Corollary 1.1.16.

For any indexed family of closed sets  $(E_i)$ , the intersection  $\bigcap_i E_i$  is closed.

```
theorem closed_intersection {ι : Type*} (E : ι → Set X)
  (h : ∀ i, IsClosed (E i)) : IsClosed (⋂ i, E i)
```

*Proof.*

A limit point of the intersection is a limit point of each member: every nearby witness in the intersection belongs to that member. Each member is closed, so it contains the point; consequently the intersection contains it.

```

apply (closed_iff_limitPoints _).mpr
intro p hp
apply mem_iInter.mpr
intro i
apply (closed_iff_limitPoints _).mp (h i)
intro r hr
obtain ⟨q, hq, hn, hd⟩ := hp r hr
exact ⟨q, mem_iInter.mp hq i, hn, hd⟩

```

□

### Lemma 1.1.17.

*The intersection of two open sets is open.*

```

theorem open_intersection (E F : Set X) (hE : IsOpen E) (hF : IsOpen F)
  ↪ :
    IsOpen (E ∩ F)

```

*Proof.*

At a point of the intersection, choose one positive radius for each set. The smaller radius gives a ball lying in both sets.

```

apply Metric.isOpen_iff.mpr
intro p hp
obtain ⟨r, hr, hEr⟩ := Metric.isOpen_iff.mp hE p hp.1
obtain ⟨s, hs, hFs⟩ := Metric.isOpen_iff.mp hF p hp.2
refine ⟨min r s, lt_min hr hs, ?_⟩
intro q hq
change dist q p < min r s at hq
exact ⟨hEr (hq.trans_le (min_le_left _ _)), hFs (hq.trans_le
  ↪ (min_le_right _ _))⟩

```

□

**Proposition 1.1.18.**

A finite intersection of open sets is open. In particular this applies to  $\bigcap_{j=0}^{n-1} E_j$  for every  $n \in \mathbb{N}$ .

```
theorem open_finite_intersection {ι : Type*} (s : Finset ι)
  (E : ι → Set X) (hE : ∀ i ∈ s, IsOpen (E i)) :
  IsOpen (⋂ i ∈ s, E i)
```

*Proof.*

For no indices the intersection is  $X$ , which contains a unit ball about each of its points. Adjoining one index replaces an already open finite intersection by its intersection with one more open set, so induction completes the proof.

```
classical
induction s using Finset.induction_on with
| empty =>
  simp only [Finset.notMem_empty, iInter_of_empty, iInter_univ]
  exact Metric.isOpen_iff.mpr (fun p _ => ⟨1, zero_lt_one, subset_univ
    ↪ _⟩)
| @insert i s hi ih =>
  rw [Finset.set_biInter_insert]
  exact open_intersection _ _ (hE i (Finset.mem_insert_self i s))
    (ih (fun j hj => hE j (Finset.mem_insert_of_mem hj)))
```

□

**Corollary 1.1.19.**

A finite union of closed sets is closed, including the union over no indices.

```
theorem closed_finite_union {ι : Type*} (s : Finset ι)
  (E : ι → Set X) (hE : ∀ i ∈ s, IsClosed (E i)) :
  IsClosed (⋃ i ∈ s, E i)
```

*Proof.*

The complements are open, so their finite intersection is open. By De Morgan's identity, this is the complement of the desired union; hence that union is closed.

```

have ho := open_finite_intersection s (fun i => (E i)c)
  (fun i hi => (hE i hi).is0pen_compl)
have heq : (⋃ i ∈ s, E i)c = ⋂ i ∈ s, (E i)c := by
  ext p
  simp only [mem_compl_iff, mem_iUnion, mem_iInter, not_exists]
  exact is0pen_compl_iff.mp (heq ▸ ho)

```

□

### Lemma 1.1.20.

$\overline{E} = E \cup E'$ , so the standard closure agrees with the textbook definition.

```

theorem closure_eq_union_limitPoints (E : Set X) :
  closure E = E ∪ limitPoints E

```

*Proof.*

If  $p \in \overline{E}$ , every ball about  $p$  meets  $E$ . Either  $p \in E$ , or each such witness differs from  $p$ , making  $p$  a limit point.

```

classical
ext p
constructor
· intro hp
  by_cases he : p ∈ E
  · exact Or.inl he
  · right
    intro r hr
    obtain ⟨q, hq, hd⟩ := Metric.mem_closure_iff.mp hp r hr
    exact ⟨q, hq, fun h => he (h ▸ hq), by rwa [dist_comm]⟩

```

Conversely, a point of  $E$  itself witnesses that every ball meets  $E$ . For a limit point, use the distinct witness in each deleted ball. Both cases give membership in the closure.

```

· intro hp
  apply Metric.mem_closure_iff.mpr
  intro r hr
  rcases hp with hp | hp
  · exact ⟨p, hp, by simp only [dist_self] using hr⟩
  · obtain ⟨q, hq, _, hd⟩ := hp r hr
    exact ⟨q, hq, by rwa [dist_comm]⟩

```

□

### Lemma 1.1.21.

*The closure of any set is closed.*

```

theorem closure_closed (E : Set X) : IsClosed (closure E)

```

*Proof.*

Suppose  $p$  is a limit point of  $\overline{E}$  and  $r > 0$ . Choose  $q \in \overline{E}$  within  $r/2$  of  $p$ , then  $s \in E$  within  $r/2$  of  $q$ . The triangle inequality puts  $s$  within  $r$  of  $p$ , proving  $p \in \overline{E}$ .

```

apply (closed_iff_limitPoints _).mpr
intro p hp
apply Metric.mem_closure_iff.mpr
intro r hr
obtain ⟨q, hq, _, hdq⟩ := hp (r / 2) (half_pos hr)
obtain ⟨s, hs, hds⟩ := Metric.mem_closure_iff.mp hq (r / 2) (half_pos
  ↪ hr)
refine ⟨s, hs, ?_⟩
have ht := dist_triangle p q s
rw [dist_comm q p] at hdq
linarith

```

□

### Proposition 1.1.22.

For every  $E \subseteq X$ :  $\overline{E}$  is closed;  $E = \overline{E}$  if and only if  $E$  is closed; and every closed set  $F$  containing  $E$  also contains  $\overline{E}$ .

**theorem** *closure\_properties* ( $E : \text{Set } X$ ) :  
 $\text{IsClosed } (\text{closure } E) \wedge (E = \text{closure } E \leftrightarrow \text{IsClosed } E) \wedge$   
 $(\forall F : \text{Set } X, \text{IsClosed } F \rightarrow E \subseteq F \rightarrow \text{closure } E \subseteq F)$

*Proof.*

We have just proved that  $\overline{E}$  is closed. Therefore  $E = \overline{E}$  implies that  $E$  is closed. Conversely, closedness gives  $E' \subseteq E$ , so  $E \cup E' = E$ .

```

refine (closure_closed E, ?_, ?_)
· constructor
  · intro h
    rw [h]
    exact closure_closed E
  · intro h
    rw [closure_eq_union_limitPoints]
    exact (union_eq_left.mpr ((closed_iff_limitPoints E).mp h)).symm

```

Let  $F$  be closed with  $E \subseteq F$ . A point of  $\overline{E}$  either lies in  $E$ , hence in  $F$ , or is a limit point of  $E$ . In the latter case the same nearby witnesses make it a limit point of  $F$ , which belongs to  $F$  by closedness.

```

· intro F hF hEF p hp
  rw [closure_eq_union_limitPoints] at hp
  rcases hp with hp | hp
  · exact hEF hp
  · apply (closed_iff_limitPoints F).mp hF
    intro r hr
    obtain (q, hq, hn, hd) := hp r hr
    exact (q, hEF hq, hn, hd)

```

□

**Proposition 1.1.23.**

If  $E \subseteq \mathbb{R}$  is nonempty and bounded above, then  $\sup E \in \overline{E}$ . If  $E$  is also closed, then  $\sup E \in E$ .

```
theorem supremum_in_closure (E : Set ℝ) (hne : E.Nonempty) (hb :
  ↪ BddAbove E) :
  sSup E ∈ closure E ∧ (IsClosed E → sSup E ∈ E)
```

*Proof.*

Write  $c = \sup E$ . For any  $r > 0$ , the smaller number  $c - r$  is not an upper bound, so choose  $q \in E$  with  $c - r < q \leq c$ . Then  $|c - q| < r$ , proving that every ball about  $c$  meets  $E$ .

```
have hc : sSup E ∈ closure E := by
  apply Metric.mem_closure_iff.mpr
  intro r hr
  obtain (q, hq, hlt) := exists_lt_of_lt_csSup hne (sub_lt_self (sSup E)
    ↪ hr)
  refine (q, hq, ?_)
  rw [Real.dist_eq, abs_of_nonneg (sub_nonneg.mpr (le_csSup hb hq))]
  linarith
```

When  $E$  is closed, its closure equals  $E$ , so the established closure membership is membership in  $E$  itself.

```
exact (hc, fun h => by
  have heq := (closure_properties E).2.1.mpr h
  rw [← heq] at hc
  exact hc)
```

□

### Proposition 1.1.24.

$$E^i = E \setminus E'.$$

```
theorem isolated_eq_diff (E : Set X) : isolatedPoints E = E \
  ↪ limitPoints E
```

*Proof.*

If  $E \cap N_r(p) = \{p\}$ , then  $p \in E$  and no distinct point of  $E$  is in that ball. This contradicts the limit-point condition, so  $p \notin E'$ .

```

classical
ext p
constructor
· rintro ⟨r, hr, heq⟩
  have hp : p ∈ E ∧ ball p r := heq.sym ▸ (mem_singleton p)
  refine ⟨hp.1, ?_⟩
  intro hl
  obtain ⟨q, hq, hn, hd⟩ := hl r hr
  have hmem : q ∈ E ∧ ball p r := ⟨hq, hd⟩
  rw [heq, mem_singleton_iff] at hmem
  exact hn hmem

```

Conversely, if  $p \in E$  but  $p \notin E'$ , some positive-radius ball contains no distinct point of  $E$ . Its intersection with  $E$  is therefore exactly  $\{p\}$ .

```

· rintro ⟨hp, hn⟩
  have hex : ∃ r > 0, ∀ q ∈ E, dist q p < r → q = p := by
    by_contra h
    push Not at h
    exact hn (fun r hr => by
      obtain ⟨q, hq, hd, hne⟩ := h r hr
      exact ⟨q, hq, hne, hd⟩)
  obtain ⟨r, hr, h⟩ := hex
  refine ⟨r, hr, ?_⟩
  ext q
  constructor
  · intro hq
    exact h q hq.1 hq.2
  · intro hq
    rw [mem_singleton_iff] at hq
    subst q
    exact ⟨hp, by simp only [mem_ball, dist_self] using hr⟩

```

□

### Lemma 1.1.25.

$$(E^\circ)^c = \overline{E^c}.$$

```

theorem compl_interior_eq_closure_compl (E : Set X) :
  (interior E)c = closure Ec

```

*Proof.*

Failure to be interior means that no positive-radius ball lies in  $E$ . Thus every such ball contains a point of  $E^c$ , giving membership in its closure. Conversely, if every ball meets  $E^c$ , no ball can be contained in  $E$ .

```
classical
ext p
rw [mem_compl_iff, mem_interior_iff_mem_nhds, Metric.mem_nhds_iff,
    Metric.mem_closure_iff]
constructor
· intro h r hr
  have hn : ¬ ball p r ⊆ E := fun hs => h ⟨r, hr, hs⟩
  obtain ⟨q, hq, hqe⟩ := Set.not_subset.mp hn
  exact ⟨q, hqe, by rwa [dist_comm]⟩
· intro h ⟨r, hr, hs⟩
  obtain ⟨q, hq, hd⟩ := h r hr
  exact hq (hs (by simp only [mem_ball, dist_comm] using hd))
```

□

**Definition 1.1.26.** The **boundary** is  $\partial E = \overline{E} \cap \overline{E^c}$ . Equivalently,  $\partial E = \overline{E} \setminus E^\circ$ .

The boundary is the closure with the interior removed.

```
noncomputable abbrev boundary (E : Set X) : Set X := frontier E
```

**Lemma 1.1.27.**

$\partial E = \overline{E} \cap \overline{E^c}$ .

```
theorem boundary_eq (E : Set X) :
  boundary E = closure E ∩ closure Ec
```

*Proof.*

Replace the complement of the interior by the closure of the complement in the difference  $\overline{E} \setminus E^\circ$ .

```
change closure E \ interior E = _
rw [sdiff_eq_compl_inter, compl_interior_eq_closure_compl, inter_comm]
```

□

**Lemma 1.1.28.**

Any two points of  $\mathbb{R}^0$  are equal, and every deleted neighborhood in  $\mathbb{R}^0$  is empty.

```
theorem zero_dimensional (p q : EuclideanSpace ℝ (Fin 0)) (r : ℝ) :
  p = q ∧ deletedNeighborhood p r = ∅
```

*Proof.*

There are no coordinates on which two empty tuples can differ, so they are equal. Removing the center consequently removes every possible point of a ball.

```
have heq : ∀ x y : EuclideanSpace ℝ (Fin 0), x = y := by
  intro x y
  ext j
  exact Fin.elim0 j
refine (heq p q, ?_)
apply Set.eq_empty_iff_forall_notMem.mpr
intro x hx
exact hx.2 (heq x p)
```

□

**Lemma 1.1.29.**

For  $n \in \mathbb{N}^+$ ,  $p \in \mathbb{R}^n$ , and  $r > 0$ , the deleted neighborhood  $\tilde{N}_r(p)$  is nonempty.

```
theorem euclidean_punctured_ball (n : ℕ+)
  (p : EuclideanSpace ℝ (Fin n)) (r : ℝ) (hr : 0 < r) :
  ∃ q : EuclideanSpace ℝ (Fin n), q ≠ p ∧ dist q p < r
```

*Proof.*

Since  $n > 0$ , the coordinate vector  $e_0$  exists. Put  $v = (r/2)e_0$ , so  $\|v\| = r/2 > 0$ . The point  $p + v$  differs from  $p$  and is at distance  $r/2 < r$  from it.

```

let v : EuclideanSpace ℝ (Fin n) := EuclideanSpace.single ⟨0, n.pos⟩ (r
  ↦ / 2)
have hv : ‖v‖ = r / 2 := by
  rw [PiLp.norm_single, Real.norm_eq_abs, abs_of_pos (half_pos hr)]
refine ⟨p + v, ?_, ?_⟩
  · intro heq
    have hz : v = 0 := add_left_cancel (heq.trans (add_zero p).symm)
    rw [hz, norm_zero] at hv
    linarith
  · rw [dist_eq_norm, add_sub_cancel_left, hv]
    linarith

```

□

### Lemma 1.1.30.

If every positive-radius deleted ball in  $X$  is nonempty, then  $E^i \subseteq (E^c)'$ .

```

theorem isolated_limit_compl (E : Set X)
  (hX : ∀ p : X, ∀ r > 0, ∃ q : X, q ≠ p ∧ dist q p < r) :
  isolatedPoints E ⊆ limitPoints Ec

```

*Proof.*

For an isolated point  $p$ , choose  $r > 0$  with  $E \cap N_r(p) = \{p\}$ . For any  $s > 0$ , take a distinct point within  $\min(r, s)$  of  $p$ . It must lie outside  $E$ , and therefore witnesses that  $p$  is a limit point of  $E^c$ .

```

intro p hp
obtain ⟨r, hr, heq⟩ := hp
intro s hs
obtain ⟨q, hne, hd⟩ := hX p (min r s) (lt_min hr hs)
refine ⟨q, ?_, hne, hd.trans_le (min_le_right _ _)⟩
intro hq
have hm : q ∈ E ∩ ball p r := ⟨hq, hd.trans_le (min_le_left _ _)⟩
rw [heq, mem_singleton_iff] at hm
exact hne hm

```

□

**Proposition 1.1.31.**

For  $n \in \mathbb{N}^+$  and  $E \subseteq \mathbb{R}^n$ ,  $\partial E = E^i \cup (E^c)^i \cup (E' \cap (E^c)')$ .

```
theorem euclidean_boundary (n : ℕ+)
  (E : Set (EuclideanSpace ℝ (Fin n))) :
  frontier E = isolatedPoints E ∪ isolatedPoints Ec ∪
    (limitPoints E ∩ limitPoints Ec)
```

*Proof.*

An isolated point of either side is a limit point of the other, by the preceding two results. Expand the two closures in  $\partial E$  as the set together with its limit points. Separate points which fail one of the two limit-point conditions from those satisfying both. The former are exactly the isolated points of one side; the latter form  $E' \cap (E^c)'$ .

```
have hE := isolated_limit_compl E (euclidean_punctured_ball n)
have hEc := isolated_limit_compl Ec (euclidean_punctured_ball n)
rw [compl_compl] at hEc
rw [isolated_eq_diff] at hE hEc
rw [frontier, sdiff_eq_compl_inter,
  compl_interior_eq_closure_compl, closure_eq_union_limitPoints,
  closure_eq_union_limitPoints, isolated_eq_diff, isolated_eq_diff]
ext p
have h1 := @hE p
have h2 := @hEc p
simp only [mem_inter_iff, mem_union, mem_sdiff, mem_compl_iff] at *
tauto
```

□

**Proposition 1.1.32.**

For  $Y \subseteq X$ , restricting  $d$  to  $Y \times Y$  gives a metric on  $Y$ , with the same distances.

```
theorem subspace_metric (Y : Set X) (p q z : Y) :
  dist p q = dist p.val q.val ∧ 0 ≤ dist p q ∧
  (dist p q = 0 ↔ p = q) ∧ dist p q = dist q p ∧
  dist p z ≤ dist p q + dist q z
```

*Proof.*

Nonnegativity, symmetry, and the triangle inequality are inherited from  $X$ . Distance zero means equality of the underlying points, which is also equality as points of  $Y$ .

```
exact (rfl, dist_nonneg, dist_eq_zero, dist_comm _ _, dist_triangle _ _
  ↪ _)
```

□

The resulting space is called the **subspace**  $(Y, d)$ . We continue to write  $d(p, q)$  for its distance.

**Definition 1.1.33.** For  $E \subseteq Y \subseteq X$ ,  $E$  is **open relative to  $Y$**  if each  $p \in E$  has a radius  $r > 0$  with  $Y \cap N_r(p) \subseteq E$ . It is **closed relative to  $Y$**  if  $E' \cap Y \subseteq E$ . The first condition is the ball definition in the subspace; the equivalence for closedness is checked below.

Relative openness means openness of  $E$  as a subset of the subspace  $Y$ .

```
abbrev relativeOpen (E Y : Set X) : Prop :=
  E ⊆ Y ∧ IsOpen ((Subtype.val : Y → X) ⁻¹' E)
```

Relative closedness means closedness of that subset of  $Y$ .

```
abbrev relativeClosed (E Y : Set X) : Prop :=
  E ⊆ Y ∧ IsClosed ((Subtype.val : Y → X) ⁻¹' E)
```

**Lemma 1.1.34.**

For  $E \subseteq Y$ ,  $E$  is closed in  $Y$  if and only if  $E' \cap Y \subseteq E$ .

```
theorem relative_closed (E Y : Set X) (hEY : E ⊆ Y) :
  IsClosed ((Subtype.val : Y → X) ⁻¹' E) ↔ limitPoints E ∩ Y ⊆ E
```

*Proof.*

A point of  $Y$  that is an ambient limit point of  $E$  is a subspace limit point: every witness lies in  $E \subseteq Y$ , and its distance is unchanged.

```
rw [closed_iff_limitPoints]
constructor
```

Conversely, each subspace witness has an underlying point in  $E$  with the same positive distance. Thus a subspace limit point is an ambient limit point lying in  $Y$ , and the stated condition puts it in  $E$ .

```

• intro h p hp
  apply h (show ((p, hp.2) : Y) ∈ limitPoints (Subtype.val ⁻¹' E) from
    ↪ ?_)
  intro r hr
  obtain ⟨q, hq, hne, hd⟩ := hp.1 r hr
  refine ⟨⟨q, hEY hq⟩, hq, ?_, hd⟩
  intro heq
  exact hne (congrArg Subtype.val heq)
• intro h p hp
  apply h
  refine (??, p.property)
  intro r hr
  obtain ⟨q, hq, hne, hd⟩ := hp r hr
  exact ⟨q.val, hq, fun heq => hne (Subtype.ext heq), hd⟩

```

□

### Proposition 1.1.35.

For  $E \subseteq Y \subseteq X$ ,  $E$  is open in  $Y$  if and only if  $E = V \cap Y$  for some open  $V \subseteq X$ .

```

theorem relative_open (E Y : Set X) (hEY : E ⊆ Y) :
  IsOpen ((Subtype.val : Y → X) ⁻¹' E) ↔
    ∃ V : Set X, IsOpen V ∧ E = V ∩ Y

```

*Proof.*

Suppose  $E$  is open in  $Y$ . For each  $p \in E$ , choose  $r_p > 0$  with  $Y \cap N_{r_p}(p) \subseteq E$ .

```

classical
constructor
• intro h
  have hex : ∀ p : E, ∃ r > 0, Y ∩ ball p.val r ⊆ E := by
    intro p
    obtain ⟨r, hr, hs⟩ := Metric.isOpen_iff.mp h ⟨p.val, hEY p.property⟩
    ↪ p.property
    exact ⟨r, hr, fun q hq => hs (show ((q, hq.1) : Y) ∈ ball ⟨p.val,
      ↪ hEY p.property⟩ r from hq.2)⟩

```

Let  $V$  be the union of these ambient balls. It is open. Each point of  $E$  lies in its own ball and in  $Y$ ; conversely, a point in  $V \cap Y$  lies in one chosen ball, whose intersection with  $Y$  lies in  $E$ . Hence  $E = V \cap Y$ .

```

choose r hr hs using hex
refine (⋃ p : E, ball p.val (r p), open_union _ (fun p => ball_open _
  ↪ _), ?_)
ext q
constructor
· intro hq
  exact (mem_iUnion.mpr (⟨q, hq⟩, by simp only [mem_ball, dist_self]
    ↪ using hr ⟨q, hq⟩), hEY hq)
· rintro ⟨hq, hy⟩
  obtain ⟨p, hp⟩ := mem_iUnion.mp hq
  exact hs p ⟨hy, hp⟩

```

Conversely, choose an ambient ball inside  $V$  about each point of  $E = V \cap Y$ . Restricting this ball to  $Y$  produces a subspace ball inside  $E$ .

```

· rintro ⟨V, hV, heq⟩
  apply Metric.isOpen_iff.mpr
  intro p hp
  have hpV : p.val ∈ V := (show p.val ∈ V ∩ Y from heq ▸ hp).1
  obtain ⟨r, hr, hs⟩ := Metric.isOpen_iff.mp hV p.val hpV
  refine ⟨r, hr, ?_⟩
  intro q hq
  change q.val ∈ E
  rw [heq]
  exact ⟨hs hq, q.property⟩

```

□

## 1.2 Compact Sets

**Definition 1.2.1.** A family  $(G_i)_{i \in I}$  is an **open cover** of  $E$  if each  $G_i$  is open and  $E \subseteq \bigcup_{i \in I} G_i$ .

For  $S \subseteq I$ , the subfamily indexed by  $S$  is a **subcover** if it still covers  $E$ .

An open cover consists of open sets whose union contains  $E$ .

```

abbrev openCover {ι : Type*} (E : Set X) (U : ι → Set X) : Prop :=
  (∀ i, IsOpen (U i)) ∧ E ⊆ ⋃ i, U i

```

A subcover keeps only indices in  $S$  and still contains  $E$  in its union.

```
abbrev subcover {ι : Type*} (E : Set X) (U : ι → Set X) (S : Set ι) :
  ⇔ Prop :=
  E ⊆ ⋃ i ∈ S, U i
```

**Definition 1.2.2.** A set  $E$  is **compact** if every open cover admits a finite subcover. Write  $\mathcal{K}(X)$  for the collection of compact sets. In the proofs, a finite index set is represented by a `Finset`. The standard Lean predicate has exactly this open-cover characterization, `isCompact_iff_finite_subcover`; using it changes the representation of the definition, not the mathematical premise.

$\mathcal{K}(X)$  consists of the compact subsets of  $X$ .

```
abbrev compactSets : Set (Set X) := {E | IsCompact E}
```

### Proposition 1.2.3.

*For  $E \subseteq Y \subseteq X$ ,  $E$  is compact in the subspace  $Y$  if and only if it is compact in  $X$ .*

```
theorem compact_subspace (E Y : Set X) (hEY : E ⊆ Y) :
  IsCompact ((Subtype.val : Y → X) ⁻¹' E) ↔ IsCompact E
```

*Proof.*

First suppose  $E$  is compact in  $Y$ . Restrict an ambient open cover to  $Y$ . An ambient ball contained in a cover member restricts to a subspace ball, so each restricted member is open.

```
classical
constructor
· intro h
  apply isCompact_of_finite_subcover
  intro ι U hU hc
  have ho : ∀ i, IsOpen ((Subtype.val : Y → X) ⁻¹' U i) := by
    intro i
    apply Metric.isOpen_iff.mpr
    intro p hp
    obtain ⟨r, hr, hs⟩ := Metric.isOpen_iff.mp (hU i) p.val hp
    exact ⟨r, hr, fun q hq => hs hq⟩
```

The restricted family covers  $E$  in  $Y$ . A finite subcover there gives the same finite collection of ambient members covering  $E$  in  $X$ .

```

have hcov : (Subtype.val : Y → X) -1 E ⊆ ⋃ i, Subtype.val -1 U i :=
  ↪ by
    intro p hp
    obtain ⟨i, hi⟩ := mem_iUnion.mp (hc hp)
    exact mem_iUnion.mpr ⟨i, hi⟩
obtain ⟨s, hs⟩ := h.elim_finite_subcover _ ho hcov
refine ⟨s, ?_⟩
intro p hp
obtain ⟨i, hi⟩ := mem_iUnion.mp (hs (show ((p, hEY hp) : Y) ∈
  ↪ Subtype.val -1 E from hp))
obtain ⟨his, hpi⟩ := mem_iUnion.mp hi
exact mem_iUnion.mpr ⟨i, mem_iUnion.mpr ⟨his, hpi⟩⟩

```

Conversely, let a family of subspace-open sets cover  $E$ . Each member is the intersection with  $Y$  of an ambient open set, by the relative-open-set proposition.

```

· intro h
  apply isCompact_of_finite_subcover
  intro ι U hU hc
  have hex : ∀ i, ∃ V : Set X, IsOpen V ∧ Subtype.val '' U i = V ∩ Y :=
    ↪ by
      intro i
      apply (relative_open _ Y (by rintro x ⟨p, _, rfl⟩; exact
        ↪ p.property)).mp
      have heq : (Subtype.val : Y → X) -1 (Subtype.val '' U i) = U i :=
        preimage_image_eq _ Subtype.val_injective
      rw [heq]
      exact hU i

```

Choose those ambient open sets. They cover  $E$  in  $X$ , so compactness gives finitely many of them.

```

choose V hV heq using hex
have hcov : E ⊆ ⋃ i, V i := by
  intro p hp
  obtain ⟨i, hi⟩ := mem_iUnion.mp (hc (show ((p, hEY hp) : Y) ∈
    ↪ Subtype.val -1 E from hp))
  have hm : p ∈ Subtype.val '' U i := ⟨⟨p, hEY hp⟩, hi, rfl⟩
  rw [heq i] at hm
  exact mem_iUnion.mpr ⟨i, hm.1⟩

```

For a point of  $E \subseteq Y$ , membership in a chosen ambient set is membership in its intersection with  $Y$ . Thus the corresponding finitely many original members cover  $E$  in the subspace.

```

obtain {s, hs} := h.elim_finite_subcover V hV hcov
refine {s, ?_}
intro p hp
obtain {i, hi} := mem_iUnion.mp {hs hp}
obtain {his, hpi} := mem_iUnion.mp hi
have hm : p.val ∈ Subtype.val '' U i := (heq i).symm ▸ {hpi, p.property}
obtain {q, hq, he} := hm
have hqp : q = p := Subtype.ext he
exact mem_iUnion.mpr {i, mem_iUnion.mpr {his, hqp ▸ hq}}

```

□

#### Proposition 1.2.4.

*Every compact subset  $E$  of a metric space is closed.*

```

theorem compact_closed (E : Set X) (hE : IsCompact E) : IsClosed E

```

*Proof.*

To prove that  $E^c$  is open, fix  $p \notin E$ . For each  $q \in E$ , let  $r_q = d(q, p)/2 > 0$ . The balls  $N_{r_q}(q)$  cover  $E$ .

```

classical
apply isOpen_compl_iff.mp
apply Metric.isOpen_iff.mpr
intro p hp
let r : E → ℝ := fun q => dist q.val p / 2
have hr : ∀ q : E, 0 < r q := by
  intro q
  exact half_pos (dist_pos.mpr (fun heq => hp (heq ▸ q.property)))
have hcover : E ⊆ ⋃ q : E, ball q.val (r q) := by
  intro q hq
  exact mem_iUnion.mpr ({q, hq}, by simp only [mem_ball, dist_self]
    ↪ using hr {q, hq})

```

Take a finite subcover and a common positive lower bound  $\delta$  for its radii. This is also possible when the finite subcover is empty.

```

obtain (s, hs) := hE.elim_finite_subcover _ (fun q => ball_open _ _)
  ↪ hcover
obtain (δ, hδ, hle) := finite_positive_lower_bound s r (fun q _ => hr q)
  refine (δ, hδ, ?_)

```

If  $x \in N_\delta(p) \cap E$ , one of the selected balls contains  $x$ . Then  $d(q, p) \leq d(q, x) + d(x, p) < r_q + \delta \leq 2r_q = d(q, p)$ , a contradiction. Hence  $N_\delta(p) \subseteq E^c$ .

```

intro x hx hxe
obtain (q, hq) := mem_iUnion.mp (hs hxe)
obtain (hqs, hxq) := mem_iUnion.mp hq
have hd := dist_triangle q.val x p
rw [dist_comm q.val x] at hd
have hδq := hle q hqs
change dist x p < δ at hx
change dist x q.val < dist q.val p / 2 at hxq
change δ ≤ dist q.val p / 2 at hδq
linarith

```

□

### Proposition 1.2.5.

*Every compact subset of a metric space is bounded, with the stated empty-space convention.*

```

theorem compact_bounded (E : Set X) (hE : IsCompact E) : bounded E

```

*Proof.*

If  $X$  is nonempty, choose  $p \in X$ . The balls  $N_{n+1}(p)$ , for  $n \in \mathbb{N}$ , cover  $E$ : every real distance is below some natural number by the Archimedean property.

```

classical
intro hX
obtain (p) := hX
have hc : E ⊆ ⋃ n : ℕ, ball p (n + 1 : ℝ) := by
  intro q _
  obtain (n, hn) := exists_nat_gt (dist q p)
  exact mem_iUnion.mpr (n, by change dist q p < (n : ℝ) + 1; linarith)

```

Choose a finite subcover and set  $R = 1 + \sum_{n \in S} (n + 1)$ . This is positive, even when  $S$  is empty.

```

obtain ⟨s, hs⟩ := hE.elim_finite_subcover _ (fun n => ball_open _ _) hc
let R : ℝ := 1 + ∑ n ∈ s, (n + 1 : ℝ)
have hnonneg : 0 ≤ ∑ n ∈ s, (n + 1 : ℝ) :=
  Finset.sum_nonneg (fun n _ => by positivity)
refine ⟨p, R, by dsimp [R]; linarith, ?_⟩

```

Each  $q \in E$  belongs to a selected ball. Its radius is at most the sum of all selected radii, so  $d(q, p) < R$ . Thus  $E \subseteq N_R(p)$ .

```

intro q hq
obtain ⟨n, hn⟩ := mem_iUnion.mp (hs hq)
obtain ⟨hns, hdist⟩ := mem_iUnion.mp hn
have hle : (n + 1 : ℝ) ≤ ∑ k ∈ s, (k + 1 : ℝ) :=
  Finset.single_le_sum (f := fun k : ℕ => (k + 1 : ℝ)) (fun k _ => by
    ↪ positivity) hns
change dist q p < R
change dist q p < (n + 1 : ℝ) at hdist
dsimp [R]
linarith

```

□

### Proposition 1.2.6.

If  $E$  is compact,  $F$  is closed, and  $F \subseteq E$ , then  $F$  is compact.

```

theorem closed_subset_compact (E F : Set X) (hE : IsCompact E)
  (hF : IsClosed F) (hFE : F ⊆ E) : IsCompact F

```

*Proof.*

Take an open cover of  $F$ . Add the open set  $F^c$ , giving the added member its own index distinct from the original indices.

```
classical
apply isCompact_of_finite_subcover
intro ι U hU hcover
let V : Option ι → Set X := fun i => match i with
  | none => Fc
  | some j => U j
have hV : ∀ i, IsOpen (V i) := by
  intro i
  cases i with
  | none => exact hF.isOpen_compl
  | some j => exact hU j
```

The enlarged family covers  $E$ : points in  $F$  are covered by the original family, and the others lie in  $F^c$ . Select a finite subcover of  $E$ .

```
have hc : E ⊆ ⋃ i, V i := by
  intro x _
  by_cases hx : x ∈ F
  · obtain ⟨i, hi⟩ := mem_iUnion.mp (hcover hx)
    exact mem_iUnion.mpr ⟨some i, hi⟩
  · exact mem_iUnion.mpr ⟨none, hx⟩
obtain ⟨s, hs⟩ := hE.elim_finite_subcover V hV hc
```

Discard the added member and keep the selected original indices. A point of  $F$  cannot lie in  $F^c$ , so it remains covered. This gives a finite subcover of  $F$ .

```
refine {s.biUnion Option.toFinset, ?_}
intro x hx
obtain ⟨i, hi⟩ := mem_iUnion.mp (hs (hFE hx))
obtain ⟨his, hxi⟩ := mem_iUnion.mp hi
cases i with
| none => exact False.elim (hxi hx)
| some j =>
  exact mem_iUnion.mpr ⟨j, mem_iUnion.mpr ⟨Finset.mem_biUnion.mpr ⟨some
    ↪ j, his, by simp⟩, hxi⟩⟩
```

□

**Theorem 1.2.7.**

If  $E$  is compact and  $F \subseteq E$  is infinite, then  $F' \cap E \neq \emptyset$ .

```
theorem infinite_subset_limit_point (E F : Set X) (hE : IsCompact E)
  (hFE : F ⊆ E) (hF : F.Infinite) : (limitPoints F ∩ E).Nonempty
```

*Proof.*

Assume there is no such limit point. Then each  $p \in E$  has a radius  $r_p > 0$  for which the only possible point of  $F$  inside  $N_{r_p}(p)$  is  $p$  itself. Otherwise every radius would have a distinct witness, making  $p$  a limit point.

```
classical
by_contra hn
have hlocal : ∀ p : E, ∃ r > 0, ∀ q ∈ F, dist q p.val < r → q = p.val :=
  ↪ by
    intro p
    by_contra h
    push Not at h
    apply hn
    refine ⟨p.val, ?_, p.property⟩
    intro r hr
    obtain ⟨q, hq, hd, hne⟩ := h r hr
    exact ⟨q, hq, hne, hd⟩
```

These balls cover  $E$ , so choose finitely many of them.

```
choose r hr hsingle using hlocal
have hc : E ⊆ ⋃ p : E, ball p.val (r p) := by
  intro p hp
  exact mem_iUnion.mpr ⟨(p, hp), by simp only [mem_ball, dist_self]⟩
  ↪ using hr ⟨p, hp⟩
obtain ⟨s, hs⟩ := hE.elim_finite_subcover _ (fun p => ball_open _ _) hc
```

Every point of  $F$  lies in one of the selected balls and must equal its center. Hence  $F$  is contained in the finite set of selected centers, contradicting its infinitude.

```
apply hF
apply (s.finite_toSet.image Subtype.val).subset
intro q hq
obtain ⟨p, hp⟩ := mem_iUnion.mp (hs (hFE hq))
obtain ⟨hps, hd⟩ := mem_iUnion.mp hp
exact ⟨p, hps, (hsingle p q hq hd).symm⟩
```

□

**Theorem 1.2.8.**

Suppose  $(K_i)_{i \in I}$  are compact and every finite intersection  $\bigcap_{i \in S} K_i$  is nonempty, including the intersection over no indices. Then  $\bigcap_{i \in I} K_i$  is nonempty.

```
theorem compact_finite_intersection {ι : Type*} (K : ι → Set X)
  (hK : ∀ i, IsCompact (K i))
  (hfinite : ∀ s : Finset ι, (⋂ i ∈ s, K i).Nonempty) :
  (⋂ i, K i).Nonempty
```

*Proof.*

First suppose  $I$  has an index  $i_0$ . If the full intersection were empty, the complements of the  $K_i$  would cover  $K_{i_0}$ . These complements are open because compact sets are closed.

```
classical
by_cases hi : Nonempty ι
· obtain (i₀) := hi
  by_contra hn
  have hc : K i₀ ⊆ ⋃ i, (K i)ᶜ := by
    intro p _
    by_contra hp
    apply hn
    refine ⟨p, mem_iInter.mpr ?_⟩
    intro i
    by_contra hpi
    exact hp (mem_iUnion.mpr (i, hpi))
```

A finite subcover would make the intersection over its indices together with  $i_0$  empty. The finite-intersection hypothesis instead supplies a point in all these sets, contradicting that subcover.

```
obtain (s, hs) := (hK i₀).elim_finite_subcover _
  (fun i => (compact_closed _ (hK i)).isOpen_compl) hc
obtain (p, hp) := hfinite (insert i₀ s)
have hp0 := mem_iInter.mp (mem_iInter.mp hp i₀) (Finset.mem_insert_self
  ↪ _ _)
obtain (i, hi) := mem_iUnion.mp (hs hp0)
obtain (his, hpi) := mem_iUnion.mp hi
exact hpi (mem_iInter.mp (mem_iInter.mp hp i) (Finset.mem_insert_of_mem
  ↪ his))
```

If  $I$  is empty, the hypothesis for the empty finite index set supplies a point of  $X$ . It automatically belongs to every member of the empty family.

```

• obtain ⟨p, _⟩ := hfinite ∅
  refine ⟨p, mem_iInter.mpr ?_⟩
  intro i
  exact False.elim (hi ⟨i⟩)

```

□

### Lemma 1.2.9.

Let  $(a_n)$  and  $(b_n)$  be real sequences indexed by  $n \in \mathbb{N}$  with  $a_n \leq a_{n+1} \leq b_{n+1} \leq b_n$ . There exists  $x \in \mathbb{R}$  with  $a_n \leq x \leq b_n$  for every  $n$ , so the nested intervals have nonempty intersection.

```

theorem nested_intervals (a b :  $\mathbb{N} \rightarrow \mathbb{R}$ )
  (h :  $\forall n, a\ n \leq a\ (n + 1) \wedge a\ (n + 1) \leq b\ (n + 1) \wedge b\ (n + 1) \leq b\ n$ )
   $\hookrightarrow$  :
   $\exists x : \mathbb{R}, \forall n, a\ n \leq x \wedge x \leq b\ n$ 

```

*Proof.*

The left endpoints increase and the right endpoints decrease. For any  $n, m$ , compare the indices: if  $n \leq m$ , then  $a_n \leq a_m \leq b_m$ ; otherwise  $a_n \leq b_n \leq b_m$ . Thus each  $b_m$  bounds all left endpoints.

```

have ha : Monotone a := monotone_nat_of_le_succ (fun n => (h n).1)
have hb : Antitone b := antitone_nat_of_succ_le (fun n => (h n).2.2)
have hab :  $\forall n, a\ n \leq b\ n$  := fun n => (h n).1.trans ((h n).2.1.trans (h
 $\hookrightarrow$  n).2.2)
have hcross :  $\forall n\ m, a\ n \leq b\ m$  := by
  intro n m
  rcases le_total n m with hnm | hmn
  • exact (ha hnm).trans (hab m)
  • exact (hab n).trans (hb hmn)

```

The set of left endpoints contains  $a_0$  and is bounded above by  $b_0$ . Its supremum exists. Each  $a_n$  is below that supremum, and each  $b_n$  is an upper bound for it, giving a point in every interval.

```

have hne : (range a).Nonempty := ⟨a 0, mem_range_self 0⟩
have hbd : BddAbove (range a) := ⟨b 0, by rintro x ⟨n, rfl⟩; exact
  ↪ hcross n 0⟩
refine ⟨sSup (range a), ?_⟩
intro n
exact ⟨le_csSup hbd (mem_range_self n), csSup_le hne (by rintro x ⟨m,
  ↪ rfl⟩; exact hcross m n)⟩

```

□

### Theorem 1.2.10.

Let  $k \in \mathbb{N}$  and let real endpoints satisfy  $a_{n,j} \leq a_{n+1,j} \leq b_{n+1,j} \leq b_{n,j}$  for every  $n \in \mathbb{N}$  and  $0 \leq j < k$ . Then the cells  $\prod_{j=0}^{k-1} [a_{n,j}, b_{n,j}]$  have nonempty intersection.

```

theorem nested_cells (k : ℕ) (a b : ℕ → Fin k → ℝ)
  (h : ∀ n j, a n j ≤ a (n + 1) j ∧
    a (n + 1) j ≤ b (n + 1) j ∧ b (n + 1) j ≤ b n j) :
  ∃ x : EuclideanSpace ℝ (Fin k), ∀ n j, a n j ≤ x j ∧ x j ≤ b n j

```

*Proof.*

For each coordinate, apply the nested-interval result and choose a point belonging to every interval in that coordinate. The resulting vector belongs to every cell. When  $k = 0$ , the choice has no coordinates and produces the unique empty tuple.

```

have hex : ∀ j, ∃ x : ℝ, ∀ n, a n j ≤ x ∧ x ≤ b n j :=
  fun j => nested_intervals (fun n => a n j) (fun n => b n j) (fun n =>
    ↪ h n j)
choose x hx using hex
exact ⟨WithLp.toLp 2 x, fun n j => hx j n⟩

```

□

We now prove compactness of cells from open covers. We first prove the interval case using a supremum, then combine finitely many coordinates. This also accounts for degenerate

intervals and the zero-dimensional cell. The next supporting argument applies more generally to topological spaces: their open sets contain the empty and whole sets and are closed under arbitrary unions and finite intersections. Metric spaces have these properties by the preceding results. Compactness still means the finite-open-subcover property. A map is continuous in the open-set sense: inverse images of open sets are open. Only this defining property is needed.

**Lemma 1.2.11.**

*For topological spaces  $A$  and  $B$ , if  $K \subseteq A$  is compact and  $f : A \rightarrow B$  is continuous, then  $f(K)$  is compact.*

```
theorem compact_image {A B : Type*} [TopologicalSpace A]
  ↪ [TopologicalSpace B]
  (K : Set A) (hK : IsCompact K) (f : A → B) (hf : Continuous f) :
    IsCompact (f '' K)
```

*Proof.*

Pull an open cover of  $f(K)$  back along  $f$ . The inverse images are open and cover  $K$ , so finitely many cover it.

```
apply isCompact_of_finite_subcover
intro ι U hU hc
have hcov : K ⊆ ⋃ i, f ⁻¹' U i := by
  intro x hx
  obtain (i, hi) := mem_iUnion.mp (hc ⟨x, hx, rfl⟩)
  exact mem_iUnion.mpr (i, hi)
obtain (s, hs) := hK.elim_finite_subcover _ (fun i => (hU i).preimage
  ↪ hf) hcov
```

Every point of  $f(K)$  has a preimage in  $K$ . That preimage belongs to a selected inverse image, so the point belongs to the corresponding selected cover member.

```
refine (s, ?_)
rintro y ⟨x, hx, rfl⟩
obtain (i, hi) := mem_iUnion.mp (hs hx)
obtain (his, hxi) := mem_iUnion.mp hi
exact mem_iUnion.mpr (i, mem_iUnion.mpr ⟨his, hxi⟩)
```

□

**Lemma 1.2.12.**

*A singleton in any metric space is compact.*

```
theorem singleton_compact {A : Type*} [MetricSpace A] (a : A) :
  IsCompact ({a} : Set A)
```

*Proof.*

One member of any cover contains its sole point. The subfamily containing only that member already covers the singleton.

```
apply isCompact_of_finite_subcover
intro ι U _ hc
obtain ⟨i, hi⟩ := mem_iUnion.mp (hc (mem_singleton a))
refine ⟨{i}, ?_⟩
intro x hx
rw [mem_singleton_iff] at hx
subst x
exact mem_iUnion.mpr ⟨i, mem_iUnion.mpr ⟨Finset.mem_singleton_self _,
  ↪ hi⟩⟩
```

□

**Lemma 1.2.13.**

*If  $a, b \in \mathbb{R}$  and  $a \leq b$ , then  $[a, b]$  is compact.*

```
theorem interval_compact (a b : ℝ) (hab : a ≤ b) : IsCompact (Icc a b)
```

*Proof.*

Fix an open cover. Let  $T$  consist of the endpoints  $x \in [a, b]$  for which  $[a, x]$  has a finite subcover. A cover member containing  $a$  covers  $[a, a]$ , so  $a \in T$ .

```

classical
apply isCompact_of_finite_subcover
intro ι U hU hc
let T : Set ℝ := {x | x ∈ Icc a b ∧ ∃ s : Finset ι, Icc a x ⊆ ⋃ i ∈ s,
  ↪ U i}
obtain (i₀, hi₀) := mem_iUnion.mp (hc (show a ∈ Icc a b from (le_rfl,
  ↪ hab)))
have ha : a ∈ T := by
  refine ((le_rfl, hab), {i₀}, ?_)
  intro x hx
  have heq : x = a := le_antisymm hx.2 hx.1
  subst x
  exact mem_iUnion.mpr (i₀, mem_iUnion.mpr (Finset.mem_singleton_self _,
  ↪ hi₀))

```

The set  $T$  is nonempty and bounded above by  $b$ . Let  $c = \sup T$ ; then  $a \leq c \leq b$ . Choose a cover member containing  $c$  and a radius  $r > 0$  whose ball about  $c$  stays in that member.

```

have hne : T.Nonempty := (a, ha)
have hb : BddAbove T := (b, fun x hx => hx.1.2)
let c := sSup T
have hac : a ≤ c := le_csSup hb ha
have hcb : c ≤ b := csSup_le hne (fun x hx => hx.1.2)
obtain (i, hi) := mem_iUnion.mp (hc (show c ∈ Icc a b from (hac, hcb)))
obtain (r, hr, hball) := Metric.isOpen_iff.mp (hU i) c hi

```

Choose  $t \in T$  with  $c - r/2 < t$ . Set  $d = \min(b, c + r/2)$ , which is at least  $c$ . We will extend a finite cover of  $[a, t]$  to  $[a, d]$  by adjoining the chosen member.

```

obtain (t, ht, hct) := exists_lt_of_lt_csSup hne (show c - r / 2 < sSup
  ↪ T by dsimp [c]; linarith)
obtain (s, hs) := ht.2
let d := min b (c + r / 2)
have hcd : c ≤ d := le_min hcb (by linarith)
have hd : d ∈ T := by
  refine ((hac.trans hcd, min_le_left _ _), insert i s, ?_)

```

A point  $x \in [a, d]$  with  $x \leq t$  is covered by the old subcover. Otherwise  $c - r/2 < t < x \leq c + r/2$ , so  $x$  is in the chosen ball. Therefore  $d \in T$ .

```

intro x hx
by_cases hxt : x ≤ t
· obtain ⟨j, hj⟩ := mem_iUnion.mp (hs ⟨hx.1, hxt⟩)
  obtain ⟨hjs, hxj⟩ := mem_iUnion.mp hj
  exact mem_iUnion.mpr ⟨j, mem_iUnion.mpr ⟨Finset.mem_insert_of_mem hjs,
    ↪ hxj⟩⟩
· have hxu : x ≤ c + r / 2 := hx.2.trans (min_le_right _ _)
  have hxball : x ∈ ball c r := by
    rw [mem_ball, Real.dist_eq, abs_lt]
    constructor <=> linarith
  exact mem_iUnion.mpr ⟨i, mem_iUnion.mpr ⟨Finset.mem_insert_self _ _,
    ↪ hball hxball⟩⟩

```

Since  $c$  is an upper bound of  $T$ , we have  $d \leq c$ . If  $c < b$ , both  $b$  and  $c + r/2$  are greater than  $c$ , which would give  $d > c$ . Hence  $c = b$  and  $d = b$ . The finite subcover witnessing  $d \in T$  covers the whole interval.

```

have hdc : d ≤ c := le_csSup hb hd
have hbc : b ≤ c := by
  by_contra h
  have hlt : c < d := lt_min (lt_of_not_ge h) (by linarith)
  exact (not_lt_of_ge hdc) hlt
have hdb : d = b := le_antisymm (min_le_left _ _) (hbc.trans hdc)
rw [hdb] at hd
exact hd.2

```

□

### Lemma 1.2.14.

For compact sets  $K \subseteq A$  and  $L \subseteq B$  in metric spaces,  $K \times L$  is compact in the product metric  $d((x, y), (x', y')) = \max(d(x, x'), d(y, y'))$ .

```

theorem compact_product {A B : Type*} [MetricSpace A] [MetricSpace B]
  (K : Set A) (L : Set B) (hK : IsCompact K) (hL : IsCompact L) :
  IsCompact (K ×s L)

```

*Proof.*

Fix an open cover and  $x \in K$ . For each  $y \in L$ , choose a cover member and a product ball of radius  $r_y > 0$  about  $(x, y)$  contained in it.

```
classical
apply isCompact_of_finite_subcover
intro ι U hU hc
have htube : ∀ x : K, ∃ δ > 0, ∃ s : Finset ι,
  ∀ x' ∈ ball x.val δ, ∀ y ∈ L, (x', y) ∈ ⋃ i ∈ s, U i := by
  intro x
  have hlocal : ∀ y : L, ∃ i : ι, ∃ r > 0, ball (x.val, y.val) r ⊆ U i
  ↦ := by
    intro y
    obtain (i, hi) := mem_iUnion.mp (hc (show (x.val, y.val) ∈ K ×s L
      ↦ from (x.property, y.property)))
    obtain (r, hr, hs) := Metric.isOpen_iff.mp (hU i) (x.val, y.val) hi
    exact (i, r, hr, hs)
```

The balls about the  $y$  coordinates cover  $L$ . Compactness selects finitely many; take a common positive lower bound  $\delta$  for their radii.

```
choose i r hr hs using hlocal
have hcov : L ⊆ ⋃ y : L, ball y.val (r y) := by
  intro y hy
  exact mem_iUnion.mpr ((y, hy), by simp only [mem_ball, dist_self]
    ↦ using hr (y, hy))
obtain (t, ht) := hL.elim_finite_subcover _ (fun y => ball_open _ _)
  ↦ hcov
obtain (δ, hδ, hle) := finite_positive_lower_bound t r (fun y _ => hr y)
```

If  $d(x', x) < \delta$  and  $y' \in L$ , choose a selected ball containing  $y'$ . Both coordinate distances are below its radius, so  $(x', y')$  lies in its original cover member. Thus finitely many members cover the whole strip  $N_\delta(x) \times L$ .

```
refine (δ, hδ, t.image i, ?_)
intro x' hx' y hy
obtain (z, hz) := mem_iUnion.mp (ht hy)
obtain (hzt, hyz) := mem_iUnion.mp hz
have hpair : (x', y) ∈ ball (x.val, z.val) (r z) := by
  change max (dist x' x.val) (dist y z.val) < r z
  exact max_lt (hx'.trans_le (hle z hzt)) hyz
exact mem_iUnion.mpr (i z, mem_iUnion.mpr (Finset.mem_image.mpr (z, hzt,
  ↦ rfl), hs z hpair))
```

Choose such a strip and finite index set for each  $x \in K$ . Their balls cover  $K$ , so finitely many strips suffice.

```
choose δ hδ s hs using htube
have hcov : K ⊆ ⋃ x : K, ball x.val (δ x) := by
  intro x hx
  exact mem_iUnion.mpr ((x, hx), by simp only [mem_ball, dist_self]
    ↪ using hδ (x, hx))
obtain (t, ht) := hK.elim_finite_subcover _ (fun x => ball_open _ _)
↪ hcov
```

Take the union of the finite index sets for those strips. For any  $(x, y) \in K \times L$ , one selected strip contains it, and its associated members cover it. The resulting finite family covers the product.

```
refine (t.biUnion s, ?_)
rintro (x, y) (hx, hy)
obtain (z, hz) := mem_iUnion.mp (ht hx)
obtain (hzt, hxz) := mem_iUnion.mp hz
obtain (i, hi) := mem_iUnion.mp (hs z x hxz y hy)
obtain (his, hxyi) := mem_iUnion.mp hi
exact mem_iUnion.mpr (i, mem_iUnion.mpr (Finset.mem_biUnion.mpr (z, hzt,
  ↪ his), hxyi))
```

□

### Lemma 1.2.15.

For  $n \in \mathbb{N}$  and real endpoints  $a_j \leq b_j$  for  $0 \leq j < n$ , the coordinate box  $\{x : a_j \leq x_j \leq b_j \text{ for all } j\}$  is compact in the finite-coordinate product metric.

```
theorem pi_cell_compact (n : ℕ) (a b : Fin n → ℝ) (hab : ∀ j, a j ≤ b j)
↪ :
  IsCompact {x : Fin n → ℝ | ∀ j, a j ≤ x j ∧ x j ≤ b j}
```

*Proof.*

Induct on  $n$ . With no coordinates, every vector is the same empty tuple, so the box is a compact singleton.

```

induction n with
| zero =>
  have heq : {x : Fin 0 → ℝ | ∀ j, a j ≤ x j ∧ x j ≤ b j} = {fun j =>
    ↪ Fin.elim0 j} := by
    ext x
    simp only [mem_ofPred_eq, mem_singleton_iff]
    constructor
    · intro _
      funext j
      exact Fin.elim0 j
    · intro _ j
      exact Fin.elim0 j
  rw [heq]
  exact singleton_compact _

```

For  $n + 1$  coordinates, separate coordinate 0 from the remaining  $n$ . The map adjoining that first coordinate is continuous, since each of its coordinates is a coordinate projection. The interval for coordinate 0 and the remaining box are compact, so their product is compact.

```

| succ n ih =>
  let K := {x : Fin n → ℝ | ∀ j, a j.succ ≤ x j ∧ x j ≤ b j.succ}
  let f : ℝ × (Fin n → ℝ) → (Fin (n + 1) → ℝ) := fun p => Fin.cons p.1
    ↪ p.2
  have hf : Continuous f := by
    apply continuous_pi
    intro j
    refine Fin.cases ?_ (fun k => ?_) j
    · exact continuous_fst
    · exact (continuous_apply k).comp continuous_snd
  have hk := compact_product (Icc (a 0) (b 0)) K (interval_compact _ _
    ↪ (hab 0))
    (ih (fun j => a j.succ) (fun j => b j.succ) (fun j => hab j.succ))

```

The image of this product is exactly the desired box: its first coordinate lies in the first interval, and the tail lies in the remaining intervals. Conversely, split any vector in the box into its first coordinate and tail. Compactness of continuous images completes the induction.

```

have heq : f '' (Icc (a 0) (b 0) xs K) =
  {x : Fin (n + 1) → ℝ | ∀ j, a j ≤ x j ∧ x j ≤ b j} := by
  ext x
  constructor
  · rintro ⟨⟨y, z⟩, ⟨hy, hz⟩, rfl⟩ j
    exact Fin.cases hy (fun k => hz k) j
  · intro hx
    refine ⟨(x 0, fun j => x j.succ), ⟨hx 0, fun j => hx j.succ⟩, ?_⟩
    funext j
    exact Fin.cases rfl (fun k => rfl) j
rw [← heq]
exact compact_image _ hk f hf

```

□

### Theorem 1.2.16.

For every  $n \in \mathbb{N}$  and real endpoints  $a_j \leq b_j$ , the cell  $\prod_{j=0}^{n-1} [a_j, b_j]$  is compact in  $\mathbb{R}^n$ .

```

theorem cell_compact (n : ℕ) (a b : Fin n → ℝ) (hab : ∀ j, a j ≤ b j) :
  IsCompact {x : EuclideanSpace ℝ (Fin n) | ∀ j, a j ≤ x j ∧ x j ≤ b
    ↪ j}

```

*Proof.*

The finite-coordinate and Euclidean descriptions have the same coordinates and topology. The identity on coordinates from the product description to the Euclidean description is continuous. Its image is exactly the same coordinate box, so the preceding result and compactness of continuous images apply. This includes  $n = 0$ .

```

have h := compact_image _ (pi_cell_compact n a b hab)
  (fun x : Fin n → ℝ => (WithLp.toLp 2 x : EuclideanSpace ℝ (Fin n)))
  (PiLp.continuous_toLp 2 (fun _ : Fin n => ℝ))
convert h using 1
ext x
constructor
· intro hx
  exact (WithLp.ofLp x, hx, rfl)
· rintro ⟨y, hy, rfl⟩
  exact hy

```

□

### Corollary 1.2.17.

For  $a, b \in \mathbb{R}$  with  $a \leq b$ , the closed interval  $[a, b]$  is compact.

```

theorem closed_interval_compact (a b : ℝ) (hab : a ≤ b) : IsCompact (Icc
  ↪ a b)

```

*Proof.*

This is the real interval result established in the construction of compact cells.

```

exact interval_compact a b hab

```

□

### Theorem 1.2.18.

For  $n \in \mathbb{N}$  and  $E \subseteq \mathbb{R}^n$ ,  $E$  is compact if and only if it is closed and bounded. This is the **Heine–Borel theorem**.

```

theorem heine_borel (n : ℕ) (E : Set (EuclideanSpace ℝ (Fin n))) :
  IsCompact E ↔ IsClosed E ∧ bounded E

```

*Proof.*

A compact set is closed and bounded by the metric-space results above. Conversely, suppose  $E$  is closed and bounded. Euclidean space has a zero vector even in dimension zero, so choose a ball  $N_r(p)$  containing  $E$ .

```

constructor
· intro h
  exact ⟨compact_closed E h, compact_bounded E h⟩
· rintro ⟨hc, hb⟩
  obtain ⟨p, r, hr, hs⟩ := hb ⟨0⟩

```

The box with coordinate intervals  $[p_j - r, p_j + r]$  is compact. Each coordinate difference is bounded in absolute value by the Euclidean norm, so a point in  $N_r(p)$  lies in this box. Hence  $E$  is a closed subset of a compact set and is compact.

```

let K : Set (EuclideanSpace ℝ (Fin n)) := {x | ∀ j, p j - r ≤ x j ∧ x j
  ↪ ≤ p j + r}
have hk : IsCompact K := cell_compact n _ _ (fun j => by linarith)
apply closed_subset_compact K E hk hc
intro x hx j
have hd : ‖x - p‖ < r := by simp only [mem_ball, dist_eq_norm] using hs
  ↪ hx
have hj : ‖(x - p) j‖ ≤ ‖x - p‖ := PiLp.norm_apply_le _ _
change ‖x j - p j‖ ≤ ‖x - p‖ at hj
rw [Real.norm_eq_abs] at hj
have hcoord := abs_le.mp (hj.trans (le_of_lt hd))
constructor <=> linarith

```

□

### 1.3 Connected Sets

**Definition 1.3.1.** Sets  $A, B \subseteq X$  are **separated** if  $\overline{A} \cap B = A \cap \overline{B} = \emptyset$ . A set  $E$  is **connected** if it cannot be expressed as the union of two nonempty separated sets; otherwise it is **disconnected**. The empty set is connected under this convention. The union equality automatically makes both sets subsets of  $E$ .

Two sets are separated when neither meets the closure of the other.

```

def separated {X : Type*} [MetricSpace X] (A B : Set X) : Prop :=
  Disjoint (closure A) B ∧ Disjoint A (closure B)

```

A connected set admits no separation into two nonempty parts.

```
def connected {X : Type*} [MetricSpace X] (E : Set X) : Prop :=
  ∀ A B : Set X, A.Nonempty → B.Nonempty → E = A ∪ B → ¬ separated A B
```

### Lemma 1.3.2.

If every point of  $A \subseteq \mathbb{R}$  is at most  $c$ , the same is true for every point of  $\overline{A}$ .

```
theorem closure_upper_bound (A : Set ℝ) (c : ℝ) (h : ∀ x ∈ A, x ≤ c) :
  ∀ x ∈ closure A, x ≤ c
```

*Proof.*

If  $x \in \overline{A}$  were greater than  $c$ , choose a point  $y \in A$  within  $x - c$  of  $x$ . Then  $|x - y| < x - c$  would imply  $y > c$ , contradicting the upper bound.

```
intro x hx
by_contra hn
have hc : c < x := lt_of_not_ge hn
obtain ⟨y, hy, hd⟩ := Metric.mem_closure_iff.mp hx (x - c) (sub_pos.mpr
  ↪ hc)
rw [Real.dist_eq, abs_lt] at hd
have hyc := h y hy
linarith
```

□

### Lemma 1.3.3.

If every point of  $A \subseteq \mathbb{R}$  is at least  $c$ , the same is true for every point of  $\overline{A}$ .

```
theorem closure_lower_bound (A : Set ℝ) (c : ℝ) (h : ∀ x ∈ A, c ≤ x) :
  ∀ x ∈ closure A, c ≤ x
```

*Proof.*

If  $x \in \overline{A}$  were less than  $c$ , a point  $y \in A$  within  $c - x$  of  $x$  would satisfy  $y < c$ . This contradicts the lower bound.

```
intro x hx
by_contra hn
have hc : x < c := lt_of_not_ge hn
obtain ⟨y, hy, hd⟩ := Metric.mem_closure_iff.mp hx (c - x) (sub_pos.mpr
  ↪ hc)
rw [Real.dist_eq, abs_lt] at hd
have hcy := h y hy
linarith
```

□

#### Lemma 1.3.4.

Suppose  $E \subseteq \mathbb{R}$  contains every point strictly between any two of its elements. It is impossible to have  $E = A \cup B$  with  $A, B$  separated and points  $a \in A, b \in B, a < b$ .

```
theorem no_separation_of_between (E A B : Set ℝ)
  (hbetween : ∀ a ∈ E, ∀ b ∈ E, ∀ x : ℝ, a < x → x < b → x ∈ E)
  (heq : E = A ∪ B) (hsep : separated A B)
  (a b : ℝ) (ha : a ∈ A) (hb : b ∈ B) (hab : a < b) : False
```

*Proof.*

Let  $S = A \cap [a, b]$ . It contains  $a$  and is bounded above by  $b$ , so  $c = \sup S$  exists and satisfies  $a \leq c \leq b$ . The supremum is in  $\overline{S} \subseteq \overline{A}$ .

```
let S := A ∩ Icc a b
have hane : a ∈ S := ⟨ha, le_refl, hab.le⟩
have hne : S.Nonempty := ⟨a, hane⟩
have hbd : BddAbove S := ⟨b, fun x hx => hx.2.2⟩
let c := sSup S
have hac : a ≤ c := le_csSup hbd hane
have hcb : c ≤ b := csSup_le hne (fun x hx => hx.2.2)
have hcS : c ∈ closure S := (supremum_in_closure S hne hbd).1
have hcA : c ∈ closure A := (closure_properties S).2.2 (closure A)
  (closure_closed A) (fun x hx => subset_closure hx.1) hcS
```

Separation implies  $c \notin B$ , and thus  $c < b$ . If  $c = a$ , it is already in  $E$ ; otherwise it is strictly between  $a$  and  $b$ , so the between-points hypothesis puts it in  $E$ . Since  $c \notin B$ , we have  $c \in A$ .

```

have hcnotB : c ∉ B := fun h => Set.disjoint_left.mp hsep.1 hcA h
have hcltb : c < b := lt_of_le_of_ne hcb (fun h => hcnotB (h.symm ▸ hb))
have hcE : c ∈ E := by
  by_cases he : a = c
  · rw [← he, heq]
    exact Or.inl ha
  · exact hbetween a (heq.symm ▸ Or.inl ha) b (heq.symm ▸ Or.inr hb) c
    (lt_of_le_of_ne hac he) hcltb
have hcain : c ∈ A := (show c ∈ A ∪ B from heq ▸ hcE).resolve_right
  ↪ hcnotB

```

Now separation gives  $c \notin \overline{B}$ . Consequently some  $r > 0$  satisfies  $d(c, y) \geq r$  for all  $y \in B$ .

```

have hcnotclB : c ∉ closure B := fun h => Set.disjoint_left.mp hsep.2
  ↪ hcain h
have hex : ∃ r > 0, ∀ y ∈ B, r ≤ dist c y := by
  by_contra h
  push Not at h
  exact hcnotclB (Metric.mem_closure_iff.mpr h)
obtain (r, hr, hfar) := hex

```

Choose  $\delta = \min(r, b - c)/2$ . It is positive, less than both  $r$  and  $b - c$ , and  $x = c + \delta$  lies strictly between  $a$  and  $b$ . Hence  $x \in E$ .

```

let δ := min r (b - c) / 2
have hδ : 0 < δ := half_pos (lt_min hr (sub_pos.mpr hcltb))
have hδr : δ < r := (half_lt_self (lt_min hr (sub_pos.mpr
  ↪ hcltb))).trans_le (min_le_left _)
have hδb : δ < b - c := (half_lt_self (lt_min hr (sub_pos.mpr
  ↪ hcltb))).trans_le (min_le_right _)
let x := c + δ
have hax : a < x := by dsimp [x]; linarith
have hxb : x < b := by dsimp [x]; linarith
have hxE := hbetween a (heq.symm ▸ Or.inl ha) b (heq.symm ▸ Or.inr hb) x
  ↪ hax hxb

```

The distance  $d(c, x) = \delta < r$  excludes  $x$  from  $B$ . Therefore  $x \in A \cap [a, b] = S$ , yet  $x > c = \sup S$ , a contradiction. Choosing half the available radius ensures that the new point is inside the excluded ball, not merely on its boundary.

```

have hxnotB : x ∉ B := by
  intro hx
  have hdist := hfar x hx
  have hd : dist c x = δ := by
    rw [Real.dist_eq, show c - x = -δ by dsimp [x]; ring, abs_neg,
      ↪ abs_of_pos hδ]
  rw [hd] at hdist
  linarith
have hxA := (show x ∈ A ∪ B from heq ▸ hxE).resolve_right hxnotB
have hxc : x ≤ c := le_csSup hbd (show x ∈ S from ⟨hxA, hax.le, hxb.le⟩)
dsimp [x] at hxc
linarith

```

□

### Theorem 1.3.5.

A set  $E \subseteq \mathbb{R}$  is connected if and only if, whenever  $a, b \in E$  and  $a < x < b$ , one has  $x \in E$ .

```

theorem connected_iff_between (E : Set ℝ) :
  connected E ↔ ∀ a ∈ E, ∀ b ∈ E, ∀ x : ℝ, a < x → x < b → x ∈ E

```

*Proof.*

Suppose  $E$  is connected and a point  $x$  between  $a, b \in E$  were missing. Split  $E$  into  $A = E \cap (-\infty, x)$  and  $B = E \cap (x, \infty)$ . They cover  $E$ , and  $a \in A, b \in B$  make both nonempty.

```
classical
constructor
· intro hc a ha b hb x hax hxb
  by_contra hx
  let A := E ∩ Iio x
  let B := E ∩ Ioi x
  have heq : E = A ∪ B := by
    ext y
    constructor
    · intro hy
      rcases lt_trichotomy y x with h | h | h
      · exact Or.inl ⟨hy, h⟩
      · exact False.elim (hx (h ▸ hy))
      · exact Or.inr ⟨hy, h⟩
    · rintro (h | h) <=> exact h.1
```

The closure of  $A$  stays at or below  $x$ , so it misses  $B$ . Similarly the closure of  $B$  stays at or above  $x$ , so it misses  $A$ . This is a forbidden separation.

```
apply hc A B ⟨a, ha, hax⟩ ⟨b, hb, hxb⟩ heq
constructor
· apply Set.disjoint_left.mpr
  intro y hy hB
  have hle := closure_upper_bound A x (fun z hz => hz.2.le) y hy
  exact (not_lt_of_ge hle) hB.2
· apply Set.disjoint_left.mpr
  intro y hA hy
  have hle := closure_lower_bound B x (fun z hz => hz.2.le) y hy
  exact (not_lt_of_ge hle) hA.2
```

Conversely, suppose every intermediate point is in  $E$  and assume a separation  $E = A \cup B$ . Choose  $a \in A$  and  $b \in B$ ; separation makes them distinct. If  $a < b$ , the preceding supremum argument contradicts the separation. If  $b < a$ , interchange the two sets and apply the same argument.

```

• intro h A B hA hB heq hsep
  obtain ⟨a, ha⟩ := hA
  obtain ⟨b, hb⟩ := hB
  have hne : a ≠ b := by
    intro he
    exact Set.disjoint_left.mp hsep.1 (subset_closure ha) (he.symm ▸ hb)
  rcases lt_or_gt_of_ne hne with hab | hba
  • exact no_separation_of_between E A B h heq hsep a b ha hb hab
  • have hsep' : separated B A := ⟨hsep.2.symm, hsep.1.symm⟩
    exact no_separation_of_between E B A h (heq.trans (union_comm _ _))
    ↪ hsep' b a hb ha hba

```

□

# Chapter 2

## Sequences and Series

### 2.1 Convergent Sequences

Sequences in this chapter are indexed by  $\mathbb{N} = \{0, 1, 2, \dots\}$ . The Lean declarations lie in `MathematicalAnalysis1.Chapter02` and use the metric-space context from the preceding chapter, with its definitions available. The terms of a sequence retain their indices even when values repeat:  $\{u_n : n \in \mathbb{N}\}$  is its range, whereas “finitely many exceptional terms” refers to finitely many exceptional indices.

**Definition 2.1.1.** A **sequence** in  $X$  is a function  $u : \mathbb{N} \rightarrow X$ , with terms  $u_n = u(n)$ . It **converges to**  $p \in X$  if for every  $\varepsilon > 0$  there is  $N \in \mathbb{N}$  such that  $d(u_n, p) < \varepsilon$  whenever  $n \geq N$ . We write  $u_n \rightarrow p$  or  $\lim_{n \rightarrow \infty} u_n = p$ . A sequence is **convergent** if it has a limit, and **divergent** otherwise.

Eventually every term is within any prescribed positive distance of  $p$ .

```
def convergesTo (u : ℕ → X) (p : X) : Prop :=  
  ∀ ε : ℝ, 0 < ε → ∃ N : ℕ, ∀ n ≥ N, dist (u n) p < ε
```

A convergent sequence has a limit in its ambient space.

```
abbrev convergent (u : ℕ → X) : Prop := ∃ p : X, convergesTo u p
```

A divergent sequence has no limit in that space.

```
abbrev divergent (u : ℕ → X) : Prop := ¬ convergent u
```

**Lemma 2.1.2.**

The epsilon definition of  $u_n \rightarrow p$  agrees with the standard Lean limit of a sequence.

```
theorem convergesTo_iff_tendsto (u : ℕ → X) (p : X) :
  convergesTo u p ↔ Filter.Tendsto u Filter.atTop (nhds p)
```

*Proof.*

The neighborhoods of a point in a metric space have positive-radius balls as a basis, and the eventual sets of natural indices have tails as a basis. Expressing the limit with these bases gives exactly the stated epsilon condition.

```
exact Metric.tendsto_atTop.symm
```

□

**Definition 2.1.3.** A sequence is **bounded** if its range  $\{u_n : n \in \mathbb{N}\}$  is bounded.

Boundedness concerns the set of values of the sequence.

```
abbrev boundedSequence (u : ℕ → X) : Prop := bounded (range u)
```

**Lemma 2.1.4.**

If  $u_n \rightarrow p$  in a metric space, then the range of  $u$  is bounded.

```
theorem convergent_bounded (u : ℕ → X) (p : X) (h : convergesTo u p) :
  bounded (range u)
```

*Proof.*

Use convergence with  $\varepsilon = 1$  to choose  $N$  beyond which every term is within 1 of  $p$ . Set

$$R = 1 + \sum_{n=0}^{N-1} d(u_n, p) > 0.$$

```
intro _
obtain ⟨N, hN⟩ := h 1 zero_lt_one
let R := 1 + ∑ n ∈ Finset.range N, dist (u n) p
have hsum : 0 ≤ ∑ n ∈ Finset.range N, dist (u n) p :=
  Finset.sum_nonneg (fun n _ => dist_nonneg)
refine ⟨p, R, by dsimp [R]; linarith, ?_⟩
```

For  $n < N$ , the distance  $d(u_n, p)$  is at most the finite sum defining  $R - 1$ . For  $n \geq N$ , it is less than  $1 \leq R$ . Thus every term lies in  $N_R(p)$ . The argument still works if the initial segment is empty.

```
rintro x (n, rfl)
change dist (u n) p < R
by_cases hn : n < N
· have hle := Finset.single_le_sum (f := fun n => dist (u n) p)
  (fun n _ => dist_nonneg) (Finset.mem_range.mpr hn)
  dsimp [R]
  linarith
· have htail := hN n (Nat.le_of_not_gt hn)
  dsimp [R]
  linarith
```

□

### Lemma 2.1.5.

*If a sequence converges to both  $p$  and  $q$  in a metric space, then  $p = q$ .*

```
theorem limit_unique (u : ℕ → X) (p q : X)
  (hp : convergesTo u p) (hq : convergesTo u q) : p = q
```

*Proof.*

If  $p \neq q$ , let  $r = d(p, q) > 0$ . Beyond the larger of the two thresholds for radius  $r/2$ , a single term is within  $r/2$  of both points. The triangle inequality would then give  $d(p, q) < r$ , a contradiction.

```
by_contra hne
have hd : 0 < dist p q := dist_pos.mpr hne
obtain ⟨N, hN⟩ := hp (dist p q / 2) (half_pos hd)
obtain ⟨M, hM⟩ := hq (dist p q / 2) (half_pos hd)
have hn := hN (max N M) (le_max_left _)
have hm := hM (max N M) (le_max_right _)
have ht := dist_triangle p (u (max N M)) q
rw [dist_comm p (u (max N M))] at ht
linarith
```

□

**Proposition 2.1.6.**

*A convergent sequence is bounded and has a unique limit.*

```
theorem convergent_properties (u : ℕ → X) (p : X) (hp : convergesTo u p)
  ↪ :
    bounded (range u) ∧ ∀ q : X, convergesTo u q → p = q
```

*Proof.*

The two preceding arguments give boundedness of the range and equality of any other limit with the given one.

```
exact (convergent_bounded u p hp, fun q hq => limit_unique u p q hp hq)
```

□

**Lemma 2.1.7.**

*For every  $\varepsilon > 0$ , some  $N \in \mathbb{N}$  satisfies  $1/(n+1) < \varepsilon$  for all  $n \geq N$ .*

```
theorem reciprocal_small (ε : ℝ) (hε : 0 < ε) :
  ∃ N : ℕ, ∀ n ≥ N, 1 / ((n : ℝ) + 1) < ε
```

*Proof.*

By the Archimedean property, choose  $N > 1/\varepsilon$ . For  $n \geq N$ , positivity gives  $1 < N\varepsilon < (n+1)\varepsilon$ . Dividing by  $n+1 > 0$  proves the claim.

```
obtain ⟨N, hN⟩ := exists_nat_gt (1 / ε)
refine ⟨N, ?_⟩
intro n hn
have hcast : (N : ℝ) ≤ n := Nat.cast_le.mpr hn
have hprod : 1 < (N : ℝ) * ε := (div_lt_iff₀ hε).mp hN
apply (div_lt_iff₀ (show 0 < (n : ℝ) + 1 by positivity)).mpr
nlinarith
```

□

**Example 2.1.8.** Define  $u_n = 1/(n + 1)$  for  $n \in \mathbb{N}$ . We will verify that  $u_n \rightarrow 0$  in  $\mathbb{R}$ , but that the same sequence diverges in the positive-real subspace  $\mathbb{R}^+$ . The shift makes every denominator positive under our zero-based indexing convention.

**Lemma 2.1.9.**

The real sequence  $u_n = 1/(n + 1)$  converges to 0.

```
theorem reciprocal_converges : convergesTo (fun n :  $\mathbb{N} \Rightarrow 1 / ((n : \mathbb{R}) + 1)$ ) 0
```

*Proof.*

Choose the threshold from the preceding reciprocal estimate. Since each term is positive, its distance from zero is exactly  $1/(n + 1)$ , which is eventually below  $\varepsilon$ .

```
intro  $\varepsilon$  h $\varepsilon$ 
obtain (N, hN) := reciprocal_small  $\varepsilon$  h $\varepsilon$ 
refine (N, ?_)
intro n hn
rw [Real.dist_eq, sub_zero, abs_of_pos (by positivity)]
exact hN n hn
```

□

Each  $1/(n + 1)$  also defines a point of the positive-real subspace.

```
noncomputable def positiveReciprocal (n :  $\mathbb{N}$ ) : {x :  $\mathbb{R}$  // 0 < x} :=
  (1 / ((n :  $\mathbb{R}$ ) + 1), by positivity)
```

**Lemma 2.1.10.**

The sequence of positive-real points  $u_n = 1/(n + 1)$  has no limit in  $\mathbb{R}^+$ .

```
theorem positive_reciprocal_diverges :
  ¬ ∃ p : {x :  $\mathbb{R}$  // 0 < x}, convergesTo positiveReciprocal p
```

*Proof.*

If it converged to a positive point  $p$  in the subspace, it would converge to the same real point because the distances are unchanged. Its real limit is already zero. Uniqueness would give  $p = 0$ , contradicting  $p > 0$ .

```
rintro ⟨p, hp⟩
have hreal : convergesTo (fun n : ℕ => 1 / ((n : ℝ) + 1)) p.val := hp
have heq := limit_unique _ p.val 0 hreal reciprocal_converges
exact (ne_of_gt p.property) heq
```

□

### Lemma 2.1.11.

$u_n \rightarrow p$  if and only if, for every  $\varepsilon > 0$ , the set of indices  $\{n \in \mathbb{N} : d(u_n, p) \geq \varepsilon\}$  is finite.

```
theorem convergesTo_iff_finite_exceptions (u : ℕ → X) (p : X) :
  convergesTo u p ↔ ∀ ε : ℝ, 0 < ε → {n : ℕ | ε ≤ dist (u n) p}.Finite
```

*Proof.*

If  $u_n \rightarrow p$ , choose its threshold  $N$ . Every exceptional index is less than  $N$ , so the exception set is contained in a finite initial segment.

```
constructor
```

Conversely, a finite set of natural indices has an upper bound  $N$ , including when it is empty. No index  $n \geq N + 1$  can be exceptional. Therefore every such term is within  $\varepsilon$  of  $p$ , proving convergence.

```

• intro h ε hε
  obtain ⟨N, hN⟩ := h ε hε
  apply (Finset.range N).finite_toSet.subset
  intro n hn
  apply Finset.mem_range.mpr
  by_contra hnot
  exact (not_lt_of_ge hn) (hN n (Nat.le_of_not_gt hnot))
• intro h ε hε
  obtain ⟨N, hN⟩ := (h ε hε).bddAbove
  refine ⟨N + 1, ?_⟩
  intro n hn
  by_contra hdist
  have hle := hN (show n ∈ {n : ℕ | ε ≤ dist (u n) p} from le_of_not_gt
    ↪ hdist)
  omega

```

□

### Lemma 2.1.12.

If  $p \in E'$ , there is a sequence  $(u_n)$  of points in  $E \setminus \{p\}$  that converges to  $p$ .

```

theorem sequence_at_limit_point (E : Set X) (p : X) (hp : p ∈
  ↪ limitPoints E) :
  ∃ u : ℕ → X, (∀ n, u n ∈ E ∧ u n ≠ p) ∧ convergesTo u p

```

*Proof.*

For each  $n \in \mathbb{N}$ , use the limit-point condition to choose  $u_n \in E$ , distinct from  $p$ , with  $d(u_n, p) < 1/(n + 1)$ .

```

have hex : ∀ n : ℕ, ∃ q ∈ E, q ≠ p ∧ dist q p < 1 / ((n : ℝ) + 1) :=
  fun n => hp _ (by positivity)
choose u hu hne hd using hex
refine ⟨u, fun n => ⟨hu n, hne n⟩, ?_⟩

```

Given  $\varepsilon > 0$ , the reciprocal estimate makes  $1/(n+1) < \varepsilon$  for all sufficiently large  $n$ . The chosen distance is smaller still, so  $u_n \rightarrow p$ .

```
intro ε hε
obtain ⟨N, hN⟩ := reciprocal_small ε hε
exact ⟨N, fun n hn => (hd n).trans (hN n hn)⟩
```

□

### Theorem 2.1.13.

*A sequence converges to  $p$  exactly when every neighborhood  $N_\varepsilon(p)$  contains all but finitely many of its terms, counted by indices. Also, if  $p \in E'$ , a sequence in  $E \setminus \{p\}$  converges to  $p$ .*

```
theorem neighborhood_characterization (u : ℕ → X) (p : X) :
  (convergesTo u p ↔ ∀ ε : ℝ, 0 < ε → {n : ℕ | ε ≤ dist (u n)
    ↪ p}.Finite) ∧
  (∀ E : Set X, p ∈ limitPoints E →
    ∃ v : ℕ → X, (∀ n, v n ∈ E ∧ v n ≠ p) ∧ convergesTo v p)
```

*Proof.*

The finite-exception characterization proves the first assertion, including its converse. The construction from shrinking deleted neighborhoods proves the second assertion, with the terms explicitly required to belong to  $E \setminus \{p\}$ .

```
exact ⟨convergesTo_iff_finite_exceptions u p, fun E hp =>
  ↪ sequence_at_limit_point E p hp⟩
```

□

**2.2 Subsequences****2.3 Cauchy Sequences****2.4 Upper and Lower Limits****2.5 Some Special Sequences****2.6 Series****2.7 Series of Nonnegative Terms****2.8 The Number  $e$** **2.9 The Root and Ratio Tests****2.10 Power Series****2.11 Summation by Parts****2.12 Absolute Convergence****2.13 Addition and Multiplication of Series****2.14 Rearrangements**

## **Chapter 3**

# **Continuity**

### **3.1 Limits of Functions**

### **3.2 Continuous Functions**

### **3.3 Continuity and Compactness**

### **3.4 Continuity and Connectedness**

### **3.5 Discontinuities**

### **3.6 Monotonic Functions**

### **3.7 Infinite Limits and Limits at Infinity**

## **Chapter 4**

# **Differentiation**

**4.1 The Derivative of a Real Function**

**4.2 Mean Value Theorems**

**4.3 The Continuity of Derivatives**

**4.4 L'Hospital's Rule**

**4.5 Derivatives of Higher Order**

**4.6 Taylor's Theorem**

**4.7 Differentiation of Vector-valued Functions**

## **Chapter 5**

# **The Riemann-Stieltjes Integral**

### **5.1 Definition and Existence of the Integral**

### **5.2 Properties of the Integral**

### **5.3 Integration and Differentiation**

### **5.4 Integration of Vector-valued Functions**

### **5.5 Rectifiable Curves**

# Bibliography

- [1] Walter Rudin, *Principles of Mathematical Analysis*, McGraw-Hill International Editions.
- [2] Kenneth Ross, *Elementary Analysis*, Springer Nature.
- [3] Steven Douglass, *Introduction to Mathematical Analysis*, Pearson.
- [4] Terence Tao, *Analysis I*, Springer Nature.