



MATHEMATICS DIVISION
INSTITUTE OF MATHEMATICAL SCIENCES AND PHYSICS
College of Arts and Sciences
University of the Philippines Los Baños

**A MATHEMATICAL MODEL OF AMERICAN FOULBROOD
TRANSMISSION THROUGH DRIFTING BEHAVIOUR OF EUROPEAN
HONEYBEES (APIS MELLIFERA L.)**

Kieffer Patani Santos

A SPECIAL PROBLEM SUBMITTED IN PARTIAL FULFILLMENT
OF THE REQUIREMENTS FOR THE DEGREE OF
BACHELOR OF SCIENCE IN APPLIED MATHEMATICS

June 2024

The special problem attached hereto, entitled

**“A MATHEMATICAL MODEL OF AMERICAN FOULBROOD TRANSMISSION THROUGH
DRIFTING BEHAVIOUR OF EUROPEAN HONEYBEES (APIS MELLIFERA L.)”,**

prepared and submitted by **KIEFFER PATANI SANTOS** in partial fulfillment of the requirements for the degree of BACHELOR OF SCIENCE IN APPLIED MATHEMATICS is hereby accepted.

DR. EDUARDO O. JATULAN

Special Problem Adviser

(Date Signed)

DR. MARK LEXTER D. DE LARA

Head

Mathematics Division,

Institute of Mathematical Sciences and Physics

(Date Signed)

Permission is given for the following people to have access to this special problem:

Available to the general public	Yes
Available only after consultation with author/SP adviser	No
Available only to those bound by confidentiality agreement	No

BIOGRAPHICAL SKETCH

Born on August 30, 2001, in Makati City, Kieffer P. Santos is the eldest child of Antonio P. Santos and Ruth P. Santos, and sibling to Seymour P. Santos, Calahan P. Santos, and Austein P. Santos. Growing up, he received his primary education at Golden Lampstand Grade School in San Pedro, Laguna, followed by his secondary education at Immaculate Heart of Mary School, also in San Pedro. In September 2020, Kieffer embarked on his academic journey at the University of the Philippines Los Baños (UPLB), where he pursued a degree in BS Applied Mathematics. With a keen interest in mathematical applications, Kieffer is committed to leveraging his education to contribute meaningfully to his field and beyond.

KIEFFER P. SANTOS

ACKNOWLEDGEMENT

I would like to express my heartfelt gratitude to several individuals who have contributed to my academic journey and the completion of this degree.

First and foremost, I extend my deepest appreciation to my parents, Antonio P. Santos and Ruth P. Santos, for their support and efforts throughout my education.

I am also grateful to my siblings, Seymour P. Santos, Calahan P. Santos, and Austein P. Santos, for their constant encouragement and understanding.

Additionally, I would like to thank the faculty and staff of Golden Lampstand Grade School and Immaculate Heart of Mary School for their dedication to nurturing my academic growth during my formative years.

Also, I am indebted to the University of the Philippines Los Baños (UPLB) for providing me with the opportunity to pursue my passion in Applied Mathematics and for fostering an enriching learning environment. I am particularly grateful to my special problem adviser, Sir Eduardo O. Jatulan, whose guidance, expertise, and mentorship have been invaluable in shaping my academic and personal development.

Furthermore, I would like to express my appreciation to my dear applied math friends Rhencis, Axl, JB, Chen, and others, whose camaraderie and encouragement have provided invaluable assistance throughout my academic journey. Their presence has made the challenges more manageable and the successes more meaningful.

I am deeply grateful to my dormmates Joshua and Kurt, whose friendship and companionship have significantly enriched my university experience. In particular, I want to express special appreciation to my best friend Joshua, whose dependable support and brotherhood have been a constant source of joy and strength throughout my academic journey. Their companionship has truly enriched my university experience, creating lasting memories and a sense of belonging that I will always cherish.

Finally, I extend my deepest gratitude to Denise, my girlfriend, whose unwavering love, understanding, and encouragement have been the cornerstone of my academic journey. Her steadfast support has not only been a source of inspiration but also a driving force behind my achievements.

Each of these individuals and institutions has played a significant role in shaping my academic and personal development, and for that, I am truly grateful.

To the ALMIGHTY FATHER, thank you for the guidance you gave me and to my prayers that you answered. I love you all.

ABSTRACT

KIEFFER P. SANTOS, University of the Philippines Los Baños, June 2024

A MATHEMATICAL MODEL OF INTER-COLONY SPREAD OF AMERICAN FOULBROOD IN EUROPEAN HONEYBEES (*APIS MELLIFERA* L.)

Major Professor: Eduardo O. Jatulan

American Foulbrood (AFB) poses a significant threat to European honeybee colonies worldwide, impacting bee health and apiary productivity. This study aims to deepen our comprehension of the inter-colony transmission dynamics of AFB among European honeybee colonies by developing a comprehensive compartmental model. Incorporating influential factors such as bee drifting between colonies, the model offers a nuanced exploration of disease ecology in honeybee communities. The research highlights that while a source colony initiates infection, the receiving colony exhibits a higher peak of infected bees, elevating the risk of disease transmission to neighboring colonies. Findings also suggest that bee drifting may slow infection spread across colonies. Furthermore, simulations indicate that a combination of lower drift rates and higher rejection rates delays the onset of brood infection in the receiving colony while accelerating it in the source colony. Additionally, infected broods are found to have a more significant impact on colony health than infected adult bees, warranting further study. The simulations show that lower contamination caused by these two factors—hygienic behavior of adult bees and beekeeper cleaning habits—significantly delays the onset of brood infection, emphasizing their critical role in managing AFB outbreaks. This research significantly advances our understanding of disease ecology in honeybee populations, crucial for sustainable beekeeping practices and honeybee population preservation.

Mathematics Subject Classification (2020): 92-10, 92B05, 92D25, 92D30, 92D50

Keywords: American Foulbrood (AFB), European honeybees (*Apis mellifera* L.), epidemiology, population dynamics

INTRODUCTION

American Foulbrood (AFB), caused by *Paenibacillus larvae*, is a devastating disease primarily targeting the brood of honeybee colonies, especially during the pre-pupal and pupal stages. Characteristic symptoms include sunken cappings and a foul smell, with the disease spreading through robbing, drifting bees, and human activities [1, 5, 16]. The resilience of AFB spores, which can remain viable for over 50 years, underscores the necessity for preventive measures and prompt action [5]. The transmission of AFB among bee colonies also occurs through extracted honey supers, transfer of brood and honey frames, and other contaminated hive parts [16]. Additionally, factors such as robbing, drift, and the introduction of infected queens and package bees can facilitate the spread of AFB within and between apiaries [16]. It is important to note that AFB infections do not stem from just a single bacterial spore; instead, it typically takes millions of spores to successfully infect larvae within a colony [16]. Thus, understanding the diverse transmission pathways and implementing rigorous preventive measures are essential for effective AFB management and the preservation of honeybee populations. Control and treatment methods for AFB, such as sealing and burning infected colonies, require rigorous colony management practices for prevention and effective disease extinction, with an understanding of transmission routes and proper hive management crucial for controlling AFB spread and maintaining healthy bee colonies [1, 4, 5, 16].

The lifecycle of European honeybees (*Apis mellifera* L.) progresses through distinct stages from egg to adult, vital for colony survival dependent on factors like foraging behaviour and resource availability, with peak seasons comprising substantial foraging activity requiring approximately 50 pounds of pollen and 100 pounds of honey annually for colony sustenance [7, 15, 18]. Influential factors impacting colony survival and disease dynamics include transmission rates and disease-induced death rates, with disease spread through robbing emphasizing the importance of reducing drift for disease mitigation and understanding disease dynamics elucidating mechanisms underlying colony survival or collapse [2, 3, 11]. Inter-colony behaviour, including drifting and robbing, plays a pivotal role in disease transmission within apiaries, with mite immigration contributing to varroa mite infestations, highlighting the role of worker bee behaviour and guard bees in disease transmission [14, 19, 21]. Comparisons between environmental hazards and infectious diseases underscore the complexity of honeybee colony dynamics, emphasizing the severity of colony-level infections and the interplay between environmental factors and evolutionary adaptations [2, 3].

Recent studies have delved into understanding disease dynamics in honeybee populations. Betti, Wahl, and Zamir [2] (2014) developed a mathematical model to predict

colony fate by linking disease spread dynamics with colony demographics, highlighting the critical roles of transmission and disease-induced death rates. Interestingly, increased disease severity might enhance colony survival, whereas early infection, especially within 20 days before winter, significantly harms colony health. Their model identifies hive bee recruitment age as an early survival indicator and notes that infectious diseases pose greater threats to colonies than environmental hazards like pesticides. Betti and Shaw [3] (2021) created a multi-scale mathematical model to address inter-colony disease spread in honeybee populations. They found that increased forager and drone drift accelerates disease transmission, highlighting the importance of effective guarding behavior. Their study also identifies specific behaviors in certain conditions, suggesting further research into extending the model to entire apiaries and exploring the evolutionary aspects of honeybee behavior. Datta, Bull, Budge, and Keeling [4] (2013) studied the 2010 AFB epidemic on Jersey Island using a spatial SIR model with an owner network, applying a Markov chain Monte Carlo (MCMC) scheme to determine parameters and infection times. They found that both distance- and owner-based transmissions contributed to AFB spread. Validation with a stochastic SIR model showed consistent infection levels. Their analysis indicated that earlier inspections reduced epidemic size and increased the likelihood of AFB eradication. Gavina, Rabajante, and Cervancia [8] (2014) examined the strategic placement of eusocial bee colonies to address forager competition and suboptimal crop pollination due to limited nectar and pollen sources. They suggested that optimal beehive distribution could mitigate these issues. Using linear programming, they developed models that considered factors like beekeeper preferences, colony numbers, colony strength, plant cluster carrying capacity, and apiary spatial orientation. Meanwhile, Jatulan, Rabajante, Banaay, Fajardo, and Jose [9] (2015) developed a mathematical model to analyze the intra-colony spread of AFB in European honeybees. Their compartmental (SI framework) model predicted that untreated AFB infection leads to colony collapse. They identified infection thresholds based on equilibrium states' stability and emphasized the number of infected cell combs as a key factor driving disease spread. Their findings provide valuable insights for forecasting AFB transmission timelines and evaluating control strategies to prevent potential epidemics. Khoury, Myerscough, and Barron [11] (2011) focused on forager death rates' impact on colony failure, and Prado, Requier, Crauser, Le Conte, Bretagnolle, and Alaux [13] (2020) explored honeybee lifespan dynamics. Additionally, Šekulja, Pechhacker, and Licek [21] (2014) studied drifting's role in AFB transmission, especially during young bee orientation flights. These studies collectively enhance our understanding of disease management in honeybee populations.

The primary objective of this study is to develop a compartmental model (SI model) to investigate the dynamics of inter-colony spread of AFB in European honeybee populations,

aiming to incorporate key factors such as drifting rate, effectiveness of drifting bees in integrating into other colonies, and return rate of unsuccessful drifting bees into the model to comprehensively capture AFB transmission dynamics between honeybee colonies. To the best of our knowledge, this represents the first mathematical model addressing inter-colony AFB transmission. Through analysis and numerical simulations, the study seeks to identify factors influencing the spread of AFB and assess the accuracy and effectiveness of the model. The significance of the study lies in its comprehensive analysis of AFB transmission dynamics, providing insights crucial for devising effective disease management strategies and safeguarding the economic viability of beekeeping operations. Additionally, the study's implications extend to biodiversity conservation, public health, and scientific research, highlighting its importance in addressing a pressing threat to honeybee populations and the ecosystems they support. While the study focuses on understanding the influence of factors like colony density, bee movement, behaviour, and health on AFB transmission dynamics and offers insights into long-term AFB exposure effects and critical control points for mitigating epidemics, it acknowledges limitations inherent in mathematical modeling, including simplifications and data availability constraints, which may affect the accuracy and generalizability of findings to real-world scenarios. Therefore, it's important to exercise prudence when applying the findings of this study to practical beekeeping and disease management practices.

METHODOLOGY

1.1 Main Assumptions

In this model, we assume that the two colonies are maintained within a controlled environment where food resources are abundant and consistently available, thereby eliminating the possibility of robbing behavior among the bees. This controlled setting ensures that competition for resources does not influence bee behavior, focusing the study on other dynamics. Additionally, it is presumed that there are sufficient spores present for an infected adult bee to effectively transmit American Foulbrood (AFB) within and between colonies. Due to the continuous interaction within the hive, we do not differentiate between house bees and forager bees, treating them collectively as adult bees. This simplification allows for a more streamlined model, focusing on the overall dynamics of the adult bee population without needing to account for the specific roles of individual bees. Brood can become infected through two primary pathways: either through feeding by spore-carrying adult bees or when eggs are laid in an infected cell comb. The queen's egg-laying rate is assumed to be dependent on the adult honeybee population, reflecting the natural regulation of reproduction based on colony size and health. Furthermore, we assume that 30% of the adult bee population comprises potential foragers capable of drifting. In this model, guard bees do not kill the drifting foragers but instead reject them. Rejected drifters face two possible outcomes: they may either return to their original colony or become lost within the model's dynamics, failing to reintegrate into any colony.

1.2 Mathematical Model

This mathematical model builds upon the model presented in Jatulan et al.'s paper [9] by incorporating an inter-colony interaction. Specifically, it focuses on modeling two colonies, designated as Colony 1 and Colony 2, to illustrate how different behaviours impact disease transmission between colonies. The mathematical model for Colony 1 is given by the following system:

$$\frac{dB_{1s}}{dt} = \alpha_1 P_1 B_1 + \frac{C_1}{v} M_1 - \mu_{B_s} \varphi_1 B_{1s} \quad (1)$$

$$\frac{dB_1}{dt} = M_1 - \alpha_1 P_1 B_1 - \frac{C_1}{v} M_1 - \beta_1 B_1 \quad (2)$$

$$\frac{dB_{1a}}{dt} = \beta_1 B_1 - \beta_2 B_{1a} \quad (3)$$

$$\frac{dC_1}{dt} = \mu_{B_s} \varphi_1 B_{1s} (\sigma(v - C_1)) \quad (4)$$

$$\frac{dA_{1s}}{dt} = \beta_2 \alpha_1 P_1 B_{1a} + \alpha_2 \frac{C_1}{v} A_1 + \alpha_3 P_1 A_1 - \mu A_{1s} - d_1 A_{1s} + r_1 \gamma_1 d_1 A_{1s} \quad (5)$$

$$\frac{dA_1}{dt} = \beta_2 (1 - \alpha_1 P_1) B_{1a} - \alpha_2 \frac{C_1}{v} A_1 - \alpha_3 P_1 A_1 - \mu A_1 - d_1 A_1 + r_1 \gamma_1 d_1 A_1 \quad (6)$$

The dynamics of Colony 2 exhibit similarities to Colony 1, albeit with distinct subscripts of the state variables and parameters C_n , P_n , M_n , d_n , γ_n , and r_n . The dynamics of Colony 2 are governed by the following system:

$$\frac{dB_{2s}}{dt} = \alpha_1 P_2 B_2 + \frac{C_2}{v} M_2 - \mu_{B_s} \varphi_1 B_{2s} \quad (7)$$

$$\frac{dB_2}{dt} = M_2 - \alpha_1 P_2 B_2 - \frac{C_2}{v} M_2 - \beta_1 B_2 \quad (8)$$

$$\frac{dB_{2a}}{dt} = \beta_1 B_2 - \beta_2 B_{2a} \quad (9)$$

$$\frac{dC_2}{dt} = \mu_{B_s} \varphi_1 B_{2s} (\sigma(v - C_2)) \quad (10)$$

$$\frac{dA_{2s}}{dt} = \beta_2 \alpha_1 P_2 B_{2a} + \alpha_2 \frac{C_2}{v} A_2 + \alpha_3 P_2 A_2 - \mu A_{2s} - d_2 A_{2s} + r_2 \gamma_2 d_2 A_{2s} \quad (11)$$

$$\frac{dA_2}{dt} = \beta_2 (1 - \alpha_1 P_2) B_{2a} - \alpha_2 \frac{C_2}{v} A_2 - \alpha_3 P_2 A_2 - \mu A_2 - d_2 A_2 + r_2 \gamma_2 d_2 A_2 \quad (12)$$

The drifting component's dynamics are governed through the following system:

$$\frac{dA_{21s}}{dt} = d_2 A_{2s} - \gamma_2 d_2 A_{2s} - \mu A_{21s} + \alpha_2 \frac{C_1}{v} A_{21} + \alpha_3 P_1 A_{21} \quad (13)$$

$$\frac{dA_{21}}{dt} = d_2 A_2 - \gamma_2 d_2 A_2 - \mu A_{21} - \alpha_2 \frac{C_1}{v} A_{21} - \alpha_3 P_1 A_{21} \quad (14)$$

$$\frac{dA_{12s}}{dt} = d_1 A_{1s} - \gamma_1 d_1 A_{1s} - \mu A_{12s} + \alpha_2 \frac{C_2}{v} A_{12} + \alpha_3 P_2 A_{12} \quad (15)$$

$$\frac{dA_{12}}{dt} = d_1 A_1 - \gamma_1 d_1 A_1 - \mu A_{12} - \alpha_2 \frac{C_2}{v} A_{12} - \alpha_3 P_2 A_{12} \quad (16)$$

From the equations, the following terms $M_1 = L_1 \frac{A_1 + A_{1s} + A_{21} + A_{21s}}{w_1 + A_1 + A_{1s} + A_{21} + A_{21s}}$, $M_2 = L_2 \frac{A_2 + A_{2s} + A_{12} + A_{12s}}{w_2 + A_2 + A_{2s} + A_{12} + A_{12s}}$ are the number of eggs that a queen can produce per day, $P_1 = \frac{A_{1s} + A_{21s}}{1 + A_1 + A_{1s} + A_{21} + A_{21s}}$, and $P_2 = \frac{A_{2s} + A_{12s}}{1 + A_2 + A_{2s} + A_{12} + A_{12s}}$ are the probability that a brood will come into contact with spore-carrier adult bees. Additionally, $P_n B_n$ represents the number of broods that can be infected by spore-carrier adult bees per day. The parameters introduced, not originally in Jatulan et al.'s paper [9], include d_1 and d_2 , representing the drifting rate of honey bees times the percentage of their foragers; γ_1 and γ_2 , indicating the rejection rate of drifting honey bees not accepted by the other colony; and finally r_1 and r_2 , signifying the return rate of drifting adult bees to their original colony upon failure to integrate into the other colony.

The other key parameters that are in Jatulan et al.'s paper [9], together with the new added ones are presented in the table below:

Table 1. Significant parameters utilized in the model.

Parameters	Description	[Range], Default Value	Reference
L_n	The queen's maximum laying rate per day	[1000, 3000], 1500	[9, 10, 15]
w_n	A coefficient that influences the rate at which the term M_n approaches L_n as the number of adult bees in a colony gets large [11].	21000	[9, 11]
α_1	The rate at which broods are infected by colony spore-carrying adult bees	[0,1], 0.001	[9]
α_2	The rate at which adult bees become infected by the infected cell	[0,1], 1	[9]
α_3	The rate at which adult bees become infected upon contact with spore-carrier adult bees	[0,1], 0.0005	[9]
β_1	The rate at which broods become immune with the spores	$\frac{1}{3}$	[9], calculated from [12]
β_2	The rate at which immune broods become adult bees	$\frac{1}{17}$	calculated from [12]

μ_{B_s}	The death rate of infected broods	variable, $\frac{1}{3}$	[9, 12]
φ_1	The rate at which a diseased dead brood is cleaned by adult bees	variable, 1	[9, 12]
v	The maximum number of cells in the colony	10000	[9]
σ	The percentage of clean cell combs that will be contaminated by AFB spores	0.001	[9]
μ	The death rate of adult bees (both spore-carrier and spore-free)	$\frac{1}{30}$	[9]
d_n	The rate at which honeybees drift multiplied by the percentage of foraging honeybees	[0,1], [0.003,0.3], 0.003*	calculated from [18] and [21]
γ_n	The rejection rate of drifting honeybees that are not accepted by the receiving colony.	0.7	[17]
r_n	The rate at which drifting adult bees return to their original colony after failing to integrate into the other colony.	[0, 1], 0.64	[12, 13]

*the subscript $n = 1, 2$ representing the two colonies.

*~1% drift rate [21] x ~30% are foragers [18]


In the model, we examined 16 state variables, with parameter details provided in the previous table, **Table 1**. We applied the Susceptible-Infectious (SI) framework to depict the transmission of American foulbrood (AFB) among broods and adult bees between colonies.

Table 2. Description of the state variables.


Variables	Description
B_{1s}	The number of infected broods in colony 1
B_1	The number of healthy broods in colony 1 before k hours old (k depends on the susceptibility of bees)
B_{1a}	The number of healthy broods in colony 1 beyond k hours old

C_1	The number of infected cell combs in colony 1 (due to diseased dead brood)
A_{1s}	The number of adult bees with AFB spores in colony 1
A_1	The number of adult bees without AFB spores in colony 1
B_{2s}	The number of infected broods in colony 1
B_2	The number of healthy broods in colony 2 before k hours old (k depends on the susceptibility of bees)
B_{2a}	The number of healthy broods in colony 2 beyond k hours old
C_2	The number of infected cell combs in colony 2 (due to diseased dead brood)
A_{2s}	The number of adult bees with AFB spores in colony 2
A_2	The number of adult bees without AFB spores in colony 2
A_{21s}	The number of infected adult bees from colony 2 that drift into colony 1.
A_{21}	The number of adult bees from colony 2 that drift into colony 1.
A_{12s}	The number of infected adult bees from colony 1 that drift into colony 2.
A_{12}	The number of adult bees from colony 1 that drift into colony 2.

To be able to come up with equations, it is important to identify the relationships among the established variables and parameters. **Figure 1** depicts the intra-colony transmission flow diagram of AFB. The intra-colony behaviour mirrors that described in Jatulan et al.'s paper [9], with supplementary parameters outlining inter-colony behaviour that **Figure 2** illustrates.




(a)




(b)

Figure 1. Compartmental diagram illustrating the interaction of honeybees and the spread of AFB within Colony 1 (a) and Colony 2 (b). Red arrows represent infection, while blue arrows denote transitions between states (refer to Table 2 for the definitions of state variables).



(a)



(b)

Figure 2. Diagram illustrating inter-colony honeybee behaviour between the two colonies. Subfigure (a) depicts the drifting behaviour dynamics, while (b) illustrates infection dynamics. In (a), dark blue arrows represent honeybees leaving their colony to drift to the other colony, while orange arrows indicate the return movement of foreign honeybees either back to their original colony or outside the two colonies.

Upon merging all the diagrams, the resulting comprehensive diagram is displayed in **Figure 3**. **Figure 3** provides an overview of the complete mathematical model depicting the spread of AFB through drifting behaviour, constituting the primary focus of this study.




Figure 3. The resulting comprehensive model diagram depicting the AFB transmission dynamics.

Lastly, Table 3 provides explanations for the terms utilized in each equation, enhancing clarity and facilitating a better understanding of the dynamics involved.

Table 3. Explanation of terms used in equations.

Terms	Relevant Equation	Description
$\alpha_1 P_1 B_1 ; \alpha_1 P_2 B_2$	Eqn (1), (2), (7), (8)	The number of broods infected daily through contact with spore-carrying adult bees
$\frac{C_1}{v} M_1 ; \frac{C_2}{v} M_2$	Eqn (1), (2), (7), (8)	The number of broods infected daily when laid in infected cells. The expression $\frac{C_n}{v}$ represents the probability that a brood will be laid in an infected cell.
$\mu_{B_s} \varphi_1 B_{1s} ; \mu_{B_s} \varphi_1 B_{2s}$	Eqn (1), (7)	The number of cells infected after an infected brood die.
$\beta_1 B_1 ; \beta_1 B_2$	Eqn (2), (3), (8), (9)	The number of young broods that develop resistance to AFB spores. This means the broods will survive

		even if they carry spores. However, immunity to spores does not make the broods spore-free; they can still be spore carriers.
$\beta_2 B_{1a} ; \beta_2 B_{2a}$	Eqn (3), (9)	The number of broods that mature into adult bees each day. Broods that survive AFB may still carry spores when they become adult bees.
$\mu_{B_s} \varphi_1 B_{1s} (\sigma(v - C_1) ;$ $\mu_{B_s} \varphi_1 B_{2s} (\sigma(v - C_2)$	Eqn (4), (10)	The number of cell combs that can be contaminated by adult bees during the cleaning or removal of diseased dead brood.
$\beta_2 \alpha_1 P_1 B_{1a} ; \beta_2 \alpha_1 P_2 B_{2a}$	Eqn (5), (11)	The number of broods that develop into spore-carrier adult bees after acquiring resistance.
$\beta_2 (1 - \alpha_1 P_1) B_{1a} ;$ $\beta_2 (1 - \alpha_1 P_2) B_{2a}$	Eqn (6), (12)	The daily number of broods that develop into spore-free adult bees.
$\alpha_2 \frac{C_1}{v} A_1 ; \alpha_2 \frac{C_2}{v} A_2 ;$ $\alpha_2 \frac{C_1}{v} A_{21} ; \alpha_2 \frac{C_2}{v} A_{12}$	Eqn (5), (6), (11), (12), (13), (14), (15), (16)	The number of adult bees that become spore carriers upon contacting infected cells. Here, $\frac{C_n}{v}$ represents the probability that a spore-free adult bee will come into contact with an infected cell.
$\alpha_3 P_1 A_1 ; \alpha_3 P_2 A_2 ; \alpha_3 P_1 A_{21}$ $; \alpha_3 P_2 A_{12}$	Eqn (5), (11), (13), (14), (15), (16)	The number of spore-free adult bees that become spore carriers upon contact with spore-carrier adult bees. The number of spore-carrier bees indirectly affects the spread of spores in the storage area, as per the main assumptions.
$\mu A_{1s} ; \mu A_1 ; \mu A_{2s} ; \mu A_2 ;$ $\mu A_{21s} ; \mu A_{21} ; \mu A_{12s} ; \mu A_{12}$	Eqn (5), (6), (11), (12), (13), (14), (15), (16)	The mortality rate of adult honeybees due to natural causes.

$d_1A_{1s} ; d_1A_1 ; d_2A_{2s} ; d_2A_2$	Eqn (5), (6), (11), (12), (13), (14), (15), (16)	The number of adult bees of appropriate age (forager bees) for leaving the colony that exhibit drifting behavior.
$r_1\gamma_1d_1A_{1s} ; r_1\gamma_1d_1A_1 ;$ $r_2\gamma_2d_2A_{2s} ; r_2\gamma_2d_2A_{2s}$	Eqn (5), (6), (11), (12)	The number of adult bees that were initially rejected during drifting but return to their original colony.
$\gamma_2d_2A_{2s} ; \gamma_2d_2A_2 ; \gamma_1d_1A_{1s}$ $; \gamma_1d_1A_1$	Eqn (13), (14), (15), (16)	This terms for example, $\gamma_2d_2A_{2s}$ is the simplified form of $-r_2\gamma_2d_2A_{2s} - (1-r_2)\gamma_2d_2A_{2s}$. It represents the number of adult bees that initially drifted away but got rejected, accounting for those that successfully returned to their original colony and those that didn't or got lost within the model's dynamics.

RESULTS

This section presents the outcomes of the simulation and numerical experiments conducted with the proposed model. The results are depicted graphically to enhance clarity and aid visualization.

For the first result of inter-colony dynamics, we incorporated the drifting behaviour of honeybees into the model of the two healthy colonies. The parameters $d_n = 0.003$, $\gamma_n = 0.7$, and $r_n = 0.64$ for all $n = 1, 2$ is incorporated and the results are shown in the figure below.




Figure 4. Dynamics of Two Healthy Colonies with Drifting Behavior. The parameters used are in **Table 1** except that $d_n = 0.003$, $\gamma_n = 0.7$, $r_n = 0.64$ for all $n = 1, 2$. The initial conditions are also $(0, 0, 0, 0, 10000, 0, 0, 0, 0, 0, 10000, 0, 0, 0, 0, 0)$ indicating that both are healthy.

The graph illustrated that when drifting occurs, the behavior of the colonies remains largely similar to when there is no drifting, albeit with the addition of foreign bees to the colony dynamics. Notably, there is only a marginal difference in the population of the original bees within the colony, but this is accompanied by the addition of the population of foreign bees. Based on the findings from **Figure 4**, we can also determine the percentage of honeybees that have drifted from the other colony, as further illustrated in **Figure 5**.




Figure 5. Percentage of drifted honeybees over time. This example from Colony 1 demonstrates behavior similar to that of Colony 2, as evidenced by the results in **Figure 4**. Therefore, we presented the results for Colony 1, recognizing their equivalence to those of Colony 2.

To further analyze the impact of drifting on the colony, we varied the drifting rate by adjusting the parameter d_n as detailed in **Table 1**. Specifically, we conducted 10 simulations with drifting rates ranging from 1% ($d_n = 0.003$) to 100% ($d_n = 0.3$) for a healthy Colony 1. The resulting effects are depicted in **Figure 6**.




Figure 6. Effects of different values of drift rate to the healthy Colony 1.

From the graph above, we observed that as the drifting rate increases for a healthy Colony 1, it may result in a weakened colony, potentially leading to collapse, particularly at higher drift rates.

Now, we delved deeper into studying the parameters influencing the dynamics of a healthy colony in drifting. In the following heatmap, we varied both the drifting rates, d_1 and d_2 , of the two colonies.




Figure 7. Dynamics of adult honeybees in two healthy colonies with varied drifting rates

The heatmap above illustrated the outcomes for the adult honey bee populations of Colony 1 and Colony 2, indicating whether they will survive or collapse as both their drifting rates are varied. From this analysis, we can conclude that both colonies will survive regardless of whether drifting is at 0% or 100%, as long as the percentage of forager bees (honeybees of an appropriate age capable of leaving the colony) in both colonies remains at 30%. If we include varying the percentage of forager bees and expand the range of d_n from 0% to 100%, the resulting heatmap is displayed below.




Figure 8. Dynamics of two colonies with varied drifting rates including the percentage of forager honeybees.

From the heatmap, we can identify points where colony collapse occurs. While there isn't a specific percentage of the total population that guarantees collapse, as colony losses are less frequently associated with collapse [6], we can set collapse as occurring when the

percentage of the current population to the original initial population is less than 1% (1000). In the heatmap, areas with a deeper blue color indicate collapse.

Additionally, we analyzed the combined effects of drift rate d_n and return rate r_n on Colony 2 to determine whether it will survive or collapse under varied parameters. By varying d_n and r_n and plotting this in a heatmap, it can visualize how these two parameters interact and influence each other within the context of honeybee behavior and colony dynamics. Their potential implications are:

- High d_n , Low r_n : This scenario suggests high levels of drift, where bees frequently leave their original colony but successfully integrate into new colonies. This could be due to weak guarding behavior in the receiving colony or aggression from resident bees, leading to the drifting bees either dying or becoming lost outside the two-colony system.
- Low d_n , High r_n : Here, while bees may initially show less inclination to drift, those that do are unsuccessful at integrating into other colonies. This may imply a less hospitable environment outside or strong guarding behavior by the receiving colony.
- High d_n , High r_n : This combination could indicate a dynamic in which bees frequently drift but struggle to integrate into new colonies, possibly suggesting an unhealthy colony or an inhospitable network of colonies.
- Low d_n , Low r_n : This scenario may suggest a stable colony environment with minimal drifting and few bees returning to their original colony, indicative of a well-connected network of colonies.

In the following figure, we analyzed the effect of varying both the drift rate d_n and return rate r_n in Colony 2.




Figure 9. Effect of varying drift rate and return rate to the survival of the adult honeybees.

From the heatmap shown in **Figure 9**, we can see that adult honeybees in Colony 2 still survived, however it became weak, even when there is no return rate ($r_n = 0$) and has high drift rate ($d_n = 0.03$) in Colony 2. The colony survival under these parameter values is further illustrated in the figure below.




Figure 10. Inter-colony dynamics of the two healthy colony when $r_n = 0$ and $d_n = 0.03$ on Colony 2.

Finally, in this scenario, if we also vary the percentage of foragers and expand the range of d_n from 0% to 100%, the survival of Colony 2 is depicted in the figure below.




Figure 11. Colony 2 survival heatmap.

We can further explore various combinations of parameters to examine the survival or collapse of a healthy colony. However, we will now shift our focus to the primary objective of this study, which involves integrating AFB infection into the inter-colony dynamics. For the first simulation involving the AFB infection, we started by adding an infected adult to Colony 2. The results are shown in **Figure 12**.




Figure 12. Dynamics of the two colonies with drifting infected adult honeybee from Colony 2. The parameters used here is also in **Table 1**. The initial conditions are $(0, 0, 0, 0, 10000, 0, 0, 0, 0, 0, 10000, 1, 0, 0, 0, 0)$, indicating the presence of an infected adult in Colony 2.

From the graphs above, it is evident that both colonies experience mortality between 250 to 300 days after the infection begins in Colony 2. Interestingly, despite the infection originating in Colony 2, the number of infected adult honeybees in Colony 1 peaks higher than that of Colony 2 over time, potentially increasing the risk of spreading the infection to other colonies when a portion of infected bees in Colony 1 drift to neighboring colonies beyond the two-colony system in the model. This result aligns with findings from Betti et al.'s paper [3], where the second colony in their study peaks slightly higher than the source colony. However, the peak we observed is more pronounced. Lastly, our results from Berkeley Madonna suggested that drifting may actually help decrease the spread of infection in both colonies compared to when there is no drifting. This finding exhibits a degree of consistency with the outcomes documented by Betti et al. [3]. The results we got from Berkeley Madonna are summarized in the table below.

Table 4. Day of First Brood Infection Onset.

Scenario	Days of First Brood Infection	
	Colony 1	Colony 2
Infection without drifting (Intra-colony)	38.9 days	38.9 days
Infection started in one colony (Colony 2) with drifting at 1% on both colonies ($d_n = 0.003$)	59.44 days	39.16 days
Infection started in one colony (Colony 2) with drifting at 10% on both colonies ($d_n = 0.03$)	53.98 days	41.54 days
Infection started in one colony (Colony 2) with drifting at 50% on both colonies ($d_n = 0.15$)	56.34 days	50.2 days
Infection started in one colony (Colony 2) with drifting at 10% on Colony 2 ($d_2 = 0.03$)	48.68 days	44.94 days
Infection started in one colony (Colony 2) with drifting at 50% on Colony 2 ($d_2 = 0.15$)	40.88 days	105.72 days
Infection started on both colonies with drifting at 1% on both colonies ($d_n = 0.003$)	39.14 days	39.14 days
Infection started on both colonies with drifting at 10% on both colonies ($d_n = 0.03$)	41.32 days	41.32 days
Infection started on both colonies with drifting at 50% on both colonies ($d_n = 0.15$)	48.78 days	48.78 days

We further explored intriguing scenarios, such as when Colony 2 is weaker than Colony 1, resulting in notable outcomes depicted in **Figure 13**.




Figure 13. Dynamics of the two colonies with drifting infected adult honeybee from a weak Colony 2. Here, we set $L_2 = 150$ and $w_2 = 2100$. We also set the initial value of $A_2 = 1000$ so that we weaken the Colony 2.

From the results, we observed that despite Colony 2 having the initial infected adult, Colony 1 is the first to collapse. This suggests that higher population density may accelerate the spread of AFB transmission within a colony. Further studies could explore whether population density serves as a critical factor in transmission dynamics.

We further investigated the dynamics using the model by varying important parameters that are likely to impact the trajectory of the system. Our initial exploration involves varying the parameter α_2 , which is detailed in **Table 1**, to observe its effects on the colony dynamics of both colonies, particularly with an infected adult honeybee present in Colony 2. The results are shown below.




Figure 14. Effect of varying α_2 on adult honeybee dynamics with infected adult in Colony 2.

From the graphs depicted in **Figure 14**, we can see that for all values of α_2 from $[0,1]$, the effect of α_2 does not help much on the colony survival, specifically the adult honeybee population.

Now, let's examine its impact on the brood infection within the system. We can see the results in the following figure below.




Figure 15. Effect of varying α_2 on brood dynamics with infected adult in Colony 2.

From the results above, it's evident that varying α_2 does not significantly influence the brood dynamics.

Next, we explored other infection rates for adult honeybees, like α_3 . Similar to α_2 , α_3 represents the rate at which adult bees become infected. The following figure showed how varying α_3 affects the adult honeybee population.




Figure 16. Effects of varying α_3 on adult honeybee dynamics with infected adult in Colony 2.

From **Figure 16**, any values from the batch runs ranging $[0,1]$ does not affect much on colony survival. It is observed that increasing values up to 1 only prolong the lifespan of adult honeybees before eventual mortality.

We further investigated the effect of varying α_3 into the brood infection now. The results can be seen from the figure below.




Figure 17. Effect of varying α_3 on brood dynamics with infected adult in Colony 2.

Based on the data presented in the figure above, it is apparent that infected broods will succumb to the infection across all values of α_3 within the range of $[0,1]$. Lower values of the parameter merely delay the eventual mortality of the infected broods.

Next, we explore another parameter related to infection rates: α_1 . Like the previous parameters, α_1 represents an infection rate, specifically concerning broods. We conducted experiments by varying α_1 , and the findings are presented below.




Figure 18. Effects of varying α_1 on adult honeybee dynamics with infected adult in Colony 2.

The findings from the graphs show a significant shift from the previous two parameters. We now observe a potential for adult honeybee survival when parameter α_1 is zero. Because of this, it is intriguing to also see what will happen to the brood dynamics. We can see the results from the following figure.




Figure 19. Effect of varying α_1 on brood dynamics with infected adult in Colony 2.

Correlating with the survival of adult honeybees, a notable observation arises when parameter $\alpha_1 = 0$, despite not being immediately apparent due to the graph displaying a flat line at 0. Both colonies in this scenario (varying α_1) exhibit survival without any infections. This finding holds significant implications, suggesting the critical importance of the infection rate among broods for colony survival and health. Consequently, it can be inferred that infected broods may constitute a more decisive factor in colony health and survival than infected adult honeybees. Further studies into these results is warranted.

Moving on from the infection rates parameters, we also studied the effects of varying γ_2 on the adult honeybee population, which is the rejection rate of drifting honeybees from Colony 2 that are not accepted by the receiving colony, as discussed in **Table 1**. The results are shown below.




Figure 20. Effects of varying γ_2 on adult honeybee dynamics with infected adult in Colony 2.

Based on the findings depicted in **Figure 20**, when Colony 1 consistently rejects all drifting adult honeybees from Colony 2, Colony 1 can survive without experiencing infection within its colony. Conversely Colony 2, where AFB infection initially originated, will still face colony collapse due to the presence of the disease.

Additionally, we investigated another parameter, φ_1 , by conducting 10 batch runs to observe the effects of varying the rate at which diseased dead brood is cleaned by adult bees on the onset of the first brood infection. The results of these experiments are presented below.




Figure 21. Effect of varying φ_1 on brood dynamics with infected adult in Colony 2.

The observed graphs indicate that as the parameter φ_1 decreases, the onset of the first brood infection decelerates, albeit with a higher peak population. Consequently, we conducted

experiments with smaller values of this parameter and investigated their impact on the dynamics of adult honeybees. The findings are presented below.




Figure 22. Effect of varying ϕ_1 [0, 0.1] on brood dynamics with infected adult in Colony 2.




Figure 23. Effects of varying ϕ_1 [0, 0.1] on adult honeybee dynamics with infected adult in Colony 2.

Further observations reveal that smaller values of ϕ_1 lead to delayed onset of AFB infection; however, if left unaddressed, they result in more severe consequences in colony health.

Now, we proceeded to another important parameter, σ , which is the percentage of clean cell combs that will be contaminated by AFB spores (**Table 1**). The following results can be seen in the figures below.




Figure 24. Effect of varying σ on brood dynamics with infected adult in Colony 2.




Figure 25. Effects of varying σ on adult honeybee dynamics with infected adult in Colony 2.

These results indicate that smaller values of σ also slow down the spread of infection within the colony. This suggests that apiary farmers may have more time to address the spread of AFB within their apiaries.

Based on these findings, we further explored the effects of different combinations of two parameters on brood infection onset using heat maps. Specifically, we varied both d_2 and γ_2 , and the resulting heat map is depicted below.




Figure 26. Effects of varying d_2 and γ_2 on brood dynamics with infected adult in Colony 2.

Analyzing the heatmaps, we determined the relationship between different parameter combinations and the onset of the first brood infection in days. Notably, when d_2 is low and γ_2 is high, the onset of brood infection occurs slower in Colony 1 but faster in Colony 2. This suggests that limited drifting in Colony 2 facilitates faster infection spread within their colony. Consequently, rejected adult drifters from Colony 2 have a higher likelihood of getting lost to another colony or returning to their own, accelerating the onset of infection.

Furthermore, we investigated the combined effects of two important parameters, σ and ϕ_1 , on the onset of brood infection. The results are displayed in the heatmaps below.




Figure 27. Effects of varying σ and ϕ_1 on brood dynamics with infected adult in Colony 2.

The heatmaps above illustrate that lower values of both parameters lead to a slower onset of the first brood infection. This observation underscores the significant influence of these

parameters on the timing and severity of AFB outbreaks, highlighting their importance in formulating effective disease control strategies for beekeeping practices.

Finally, the last simulation was the effect of varying the parameters, σ and α_2 . The importance of this relationship lies in how environmental contamination influences the likelihood and rate of infection transmission within the colony. A higher σ value means that more clean cell combs are contaminated, which in turn increases the exposure of adult bees to AFB spores and thus increases the likelihood of infection transmission, as captured by α_2 . The heatmap is shown below.




Figure 28. Effects of varying σ and α_2 on brood dynamics with infected adult in Colony 2.

From the heatmap above, we can see that α_2 does not really affect the first infection on broods. It is more important to look at σ , which denotes the percentage of clean cell combs contaminated by AFB spores, is particularly significant for assessing the vulnerability of honeybee colonies to AFB infection. A higher value of sigma indicates a greater likelihood of AFB contamination in the colony's brood cells, leading to more rapid and extensive spread of the disease.

The obtained result is further elaborated upon in the Discussion section of the paper.

DISCUSSION

dfkopgdsfkopg kdsopfgksdopfgkds opkop kfdsgopsdfk gopdsfk gsdfk opdf gkfdopgk
dsfopg ksdfopg kdsfpog ksdfopg ksdfopfg ksdfopg ksdfopg skdfopg skdfgpo sfkdg sdfgoi dfjg
iodsfj giodsfj goisdjf goisdjf goisdjfgo ijsdfo gijsdf giosjd goisdjf giosdj giosdjg iodsfi giosdjf
goisdjf gosdijf giosdjf giodsjf giojdsfgio jsdgfio jsdfio gjdsf iogjiodsf jgsiodf jgsiod fjiodsf
jgiodsfj giosdf jiosdfj giosdjf giosdjf giosdjg iosjdfgiojdsfgioijsdfogkmdsflkgmerlkm lkm
lskegmslkm lkerm lkem lkemglkerm ioefio erkfo wkop wekopfwk efpkdsfo ok opewk
fpweokfopewkf powekf pwoeokf pwoefk ewpofk weopf kweopf kweopf k we dfgl;k dsf;lkg
sdl;fgk sdl; ks;dll kl;sd ksl;dfgk ;sldf kgsdpfogksodpfkgs dfgp sdfkg opsdkg opk gpokf pok
op kfgop kodf opko kopopgodp opdop o opg op op o opsogp gsfd opgksdff opdf gpdsfop gkpdkf
posdfp kopsdfo gsodfp gopsgdopk opdkfg

REFERENCES

1. American Foulbrood. Texas Apiary Inspection Service TAIS. (2017, May). <https://txbeeinspection.tamu.edu/american-foulbrood/>
2. Betti, M. I., Wahl, L. M., & Zamir, M. (2014). Effects of infection on Honey Bee Population Dynamics: A model. PLoS ONE, 9(10). <https://doi.org/10.1371/journal.pone.0110237>
3. Betti, M., & Shaw, K. (2021). A multi-scale model of disease transfer in honey bee colonies. *Insects*, 12(8), 700. <https://doi.org/10.3390/insects12080700>
4. Datta, S., Bull, J. C., Budge, G. E., & Keeling, M. J. (2013). Modelling the spread of American Foulbrood in Honeybees. *Journal of The Royal Society Interface*, 10(88), 20130650. <https://doi.org/10.1098/rsif.2013.0650>
5. Established pests. Bee Aware. (2014, July). <https://beeaware.org.au/archive-pest/american-foulbrood/#ad-image-0>
6. Ferrier, P. (2018, October 1). Despite elevated loss rate since 2006, U.S. honey bee colony numbers are stable. USDA ERS - Despite Elevated Loss Rate Since 2006, U.S. Honey Bee Colony Numbers Are Stable. <https://www.ers.usda.gov/amber-waves/2018/october/despite-elevated-loss-rate-since-2006-u-s-honey-bee-colony-numbers-are-stable/>
7. Garvey, K. K. (2013, September 24). The foraging force of a honey bee colony. ANR Blogs. <https://ucanr.edu/blogs/blogcore/postdetail.cfm?postnum=11592>
8. Gavina, M. K., Rabajante, J. F., & Cervancia, C. R. (2014). Mathematical programming models for determining the optimal location of beehives. *Bulletin of Mathematical Biology*, 76(5), 997–1016. <https://doi.org/10.1007/s11538-014-9943-9>
9. Jatulan, E. O., Rabajante, J. F., Banaay, C. G., Fajardo, A. C., & Jose, E. C. (2015). A mathematical model of intra-colony spread of American Foulbrood in European

- honeybees (*apis mellifera* L.). PLOS ONE, 10(12).
<https://doi.org/10.1371/journal.pone.0143805>
10. Katzav-Gozansky, T., Ibarra, F., Francke, W., Hefetz, A., & Soroker, V. (2001). Dufour's gland secretion of the queen honeybee (*apis mellifera*): An egg discriminator pheromone or a queen signal? *Behavioural Ecology and Sociobiology*, 51(1), 76–86.
<https://doi.org/10.1007/s002650100406>
 11. Khoury, D. S., Myerscough, M. R., & Barron, A. B. (2011). A quantitative model of Honey Bee Colony Population Dynamics. *PLoS ONE*, 6(4).
<https://doi.org/10.1371/journal.pone.0018491>
 12. Laomettachit, T., Liangruksa, M., Termsaithong, T., Tangthanawatsakul, A., & Duangphakdee, O. (2021). A model of infection in honeybee colonies with social immunity. *PLOS ONE*, 16(2). <https://doi.org/10.1371/journal.pone.0247294>
 13. Prado, A., Requier, F., Crauser, D., Le Conte, Y., Bretagnolle, V., & Alaux, C. (2020). Honeybee lifespan: The critical role of pre-foraging stage. *Royal Society Open Science*, 7(11), 200998. <https://doi.org/10.1098/rsos.200998>
 14. Randy. (2023, May 8). A study on Bee Drift and mite immigration: Part 1. Scientific Beekeeping. <https://scientificbeekeeping.com/a-study-on-bee-drift-and-mite-immigration-part-1/#:~:text=the%20drone%20factor,-,HOW%20MUCH%20DRIFT%20IS%20THERE%20OF%20WORKER%20BEES%3F,hives%20%5B%5B13%5D%5D>.
 15. Schmus, B. (2022, October 27). The honey bee lifecycle. The Best Bees Company. <https://bestbees.com/2022/08/08/bee-lifecycle/>
 16. Spread of AFB: The Management Agency, National American Foulbrood Pest Management Plan New Zealand. The Management Agency, National American Foulbrood Pest Management Plan New Zealand | Elimination of American Foulbrood (AFB) Disease in New Zealand. (2019, April 11). <https://afb.org.nz/spread-of-afb/>
 17. Staughton, J. (2024, March 19). What happens to bees when they get lost?. Science ABC. <https://www.scienceabc.com/nature/animals/what-happens-to-bees-when-they-get-lost.html>
 18. Van Der Steen, J. (2015, February). The Foraging Honey Bee. Wageningen; Plant Research International.
 19. Wildflowermeadows. (2019, October 22). Drifting. Wildflower Meadows. https://wildflowermeadows.com/2019/10/drifting/?fbclid=IwAR1bdt9f_t3ocsow-a6RSkcIFk_YGo_zio-1A6s8mpt3OO-

b98wQymae2yE#:~:text=Looking%20at%20a%20large%20apiary,the%20correct%20c
olony%20each%20time

20. Willingham, R., Klopchin, J., & Ellis, J. (2021, July 20). Eny-163/IN1064: Robbing behaviour in Honey Bees. Ask IFAS - Powered by EDIS. <https://edis.ifas.ufl.edu/publication/IN1064>
21. Šekulja, D., Pechhacker, H., & Licek, E. (2014). Drifting Behaviour of Honey Bees (*Apis Mellifera Carnica* Pollman, 1879) in the Epidemiology of American Foulbrood. *Zbornik Veleučilišta u Rijeci*, 2(1), 345–358. <https://hrcak.srce.hr/en/file/443139>