

## Convolution

$(f * g)(t) = \int_{-\infty}^{\infty} f(\tau)g(t - \tau)d\tau$ , additionally for  $f, g : [0, \infty) \rightarrow \mathbb{R}$  borders are from 0 to  $t$ .

## Laplace Transform

$F(s) = \mathcal{L}(f(t)) = \int_0^{\infty} f(t)e^{-st}dt$  is the Laplace transform of  $f(t)$ .

### 0.1 Rules

- $\mathcal{L}(y(t)) = Y(s)$
- $\mathcal{L}(y'(t)) = s \cdot Y(s)$
- $\mathcal{L}(y''(t)) = s^2 \cdot Y(s)$

## Misc

$y(t)$  is output,  $u(t)$  is input. Transfer function:  $G(s) = \frac{Y(s)}{U(s)}$

TODO: Table 1.1

Response to an input:

1. Given:  $ay'(t) + by(t) = cu(t)$
2. Laplace:  $a \cdot s \cdot Y(s) + bY(s) = cU(s)$
3. Transfer function:  $G(s) = \frac{Y(s)}{U(s)} = \frac{c}{a \cdot s + b}$
4. Transfer given input:  $\mathcal{L}(u(t)) = U(s)$
5. Get output:  $Y(s) = G(s) \cdot U(s)$
6. Transfer back output, hint partial fractions!