

Questions

1. Given Euler angles $\theta_x, \theta_y, \theta_z$, the corresponding rotation matrix is given by

$$R = R_x R_y R_z = \begin{pmatrix} C_y C_z & -C_y S_z & S_y \\ S_x S_y C_z + C_x S_z & -S_x S_y S_z + C_x C_z & -S_x C_y \\ -C_x S_y C_z + S_x S_z & C_x S_y S_z + S_x C_z & C_x C_y \end{pmatrix}$$

where

$$\begin{aligned} C_x &= \cos \theta_x & S_x &= \sin \theta_x \\ C_y &= \cos \theta_y & S_y &= \sin \theta_y \\ C_z &= \cos \theta_z & S_z &= \sin \theta_z \end{aligned}$$

Consider the matrix

$$R = \begin{pmatrix} 0 & 0 & 1 \\ \frac{\sqrt{2}}{2} & \frac{\sqrt{2}}{2} & 0 \\ -\frac{\sqrt{2}}{2} & \frac{\sqrt{2}}{2} & 0 \end{pmatrix}$$

Find angles $\theta_x, \theta_y, \theta_z$ such that $R = R_x R_y R_z$. Are there more choices of $\theta_x, \theta_y, \theta_z$?

Solution: One possible solution is $(45^\circ, 90^\circ, 0^\circ)$. This is not the only possibility.

2. Given an axis \mathbf{r} (unit vector) and an angle θ , the matrix representing the rotation of θ around \mathbf{r} is given by

$$R_{\mathbf{r}\theta} = \begin{pmatrix} tx^2 + c & txy - sz & txz + sy \\ txy + sz & ty^2 + c & tyz - sx \\ txz - sy & tyz + sx & tz^2 + c \end{pmatrix}$$

where

$$\begin{aligned} \mathbf{r} &= (x, y, z) \\ c &= \cos \theta \\ s &= \sin \theta \\ t &= 1 - \cos \theta \end{aligned}$$

Consider the rotation matrix

$$R = \begin{pmatrix} 0 & 0 & 1 \\ 0 & -1 & 0 \\ 1 & 0 & 0 \end{pmatrix}$$

Determine the rotation angle and rotation axis.

Solution: $180^\circ, (1, 0, 1)^T$ (your answer may differ by a positive scaling factor)

3. Determine the quaternion that corresponds to rotation around the axis $(-1, -2, -2)^T$ with $\theta = 270^\circ$.

Solution: $(-\frac{\sqrt{2}}{2}, -\frac{\sqrt{2}}{6}, -\frac{\sqrt{2}}{3}, -\frac{\sqrt{2}}{3})$

4. Find the quaternion q that corresponds to rotation around the axis $(1, 1, 0)$ with 90° . Consider the point $(1, 1, 1)$. Let $p_1 = (0, 1, 1, 1)$ be the corresponding quaternion. Calculate qp_1q^{-1} . Let $p_2 = (0, 1, 1, 0)$. Argue that qp_2q^{-1} equals p_2 . Show this by performing the exact calculations.

Solution: $(\frac{\sqrt{2}}{2}, \frac{1}{2}, \frac{1}{2}, 0)$