

# The Determinant and Arc Indices of $\theta$ -Curves and Handcuff-Graphs

---

Eunchan Cho<sup>1</sup>   Jeongwon Shin<sup>1</sup>   Boyeon Seo<sup>1</sup>   Minho Choi<sup>1</sup>

NOV 22, 2025  
R&E 2025


<sup>1</sup>Korea Science Academy of KAIST

# Introduction

---


# $\theta$ -Curves

- A  **$\theta$ -curve**  $T$  is a graph embedded in  $S^3$ , which consists of two vertices  $v_1, v_2$  and three edges  $e_1, e_2, e_3$ , such that each edge joins the vertices.
- A **constituent knot**  $T_{ij}$ ,  $1 \leq i < j \leq 3$ , is a subgraph of  $T$  that consists of two vertices  $v_1, v_2$  and two edges  $e_i, e_j$ .
- $\theta$ -curves are roughly classified by comparing the triples of constituent knots.
- A  $\theta$ -curve is said to be **trivial** if it can be embedded in a 2-sphere in  $S^3$ .



# Handcuff Graphs

- A **handcuff graph**  $H$  is a graph embedded in  $S^3$  consisting of two vertices ( $v_1, v_2$ ) and three edges ( $e_1, e_2, e_3$ ), where  $e_3$  has distinct endpoints  $v_1$  and  $v_2$ , and  $e_1$  and  $e_2$  are loops based at  $v_1$  and  $v_2$ .
- A **constituent link**  $H_{12}$ , is a subgraph of  $H$  that consists of two vertices  $v_1, v_2$  and two edges  $e_1, e_2$ .



- **Arc presentation** of a  $\theta$ -curve or handcuff graph is an embedding of them.
- It is contained in the union of finitely many half planes (called **pages**).
- The embedding is with the common boundary line (called **axis**).
- Each vertex lies in the axis.
- Each page contains a properly embedded single arc.
- **Arc index**, is the minimal number of pages among all possible arc presentations of graph.
- This arc presentation with the minimal number of pages is **minimal arc presentation**.

# Arc Presentation



Trefoil




Open Book



Grid Diagram



$\theta_{5,2}$



Open Book




Grid Diagram



$2_1$

$\Phi_{2,1}$



Open Book





Grid Diagram

## Grid Diagram

- The **grid diagram** of  $\theta$ -curve or handcuff graph is a diagram with only vertical and horizontal strands.
- $(\text{number of vertical strands}) + 1 = (\text{number of horizontal strands})$
- At every crossing, the vertical strand crosses over the horizontal strand.
- No two horizontal strands are in the same row.
- No two vertical strands are in same column.

# Grid Diagram

- A grid diagram gives rise to an arc presentation and vice versa.







# Arc Presentation of the $\theta$ -Curve and Handcuff Graph

## Theorem

*Every  $\theta$ -curve and handcuff graph admit a grid diagram.*

## PROOF



## Corollary

*Every  $\theta$ -curve and handcuff graph admit a arc presentation.*


## Determinant of $\theta$ -curve and Handcuff graph

---

# THC-cromwell matrix

- The **Cromwell Matrix** of a knot is an  $n \times n$  binary matrix each of whose rows and columns has exactly two 1s.
- The **THC-cromwell matrix** is an expansion of cromwell matrix into  $\theta$ -curves and handcuff graphs.

$$\begin{bmatrix} 1 & 0 & 1 & 0 & 0 \\ 0 & 1 & 0 & 1 & 0 \\ 0 & 0 & 1 & 0 & 1 \\ 1 & 0 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 & 1 \end{bmatrix}$$




$$\begin{bmatrix} 1 & 1 & 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 & 0 & 1 \\ 1 & 0 & 0 & 0 & 1 & 0 \\ 0 & 1 & 1 & 0 & 0 & 1 \end{bmatrix}$$

# Determinants of the cromwell matrices of Knot

## Theorem


Let  $K$  be a knot. Then the determinant of a cromwell matrix of  $K$  is 0 or  $\pm 2$ .

## PROOF



## H-deletion of THC-cromwell matrices

- The **vertex-row** of THC-cromwell matrix  $M$  is a row which contains three 1s,  $M_{ia}, M_{ib}, M_{ic}$ , where  $a < b < c$ , as its elements.
- The **side-column** of THC-cromwell matrix  $M$  is a column which contains the leftmost 1 of vertex-row ( $M_{ia}$ ) or the rightmost 1 of vertex row ( $M_{ic}$ ).
- The **H-deletion** Matrix of the THC-cromwell matrix  $G$  is  $(n - 1) \times (n - 1)$  matrix which deleted vertex-row and its two side-columns from the matrix  $G$ .



# Determinants of the THC-cromwell matrices

## Theorem


Let  $M$  be a THC-cromwell matrix of  $\theta$ -curve or handcuff graph.

- $\det^*(M) = \pm 1 \iff M$  represents  $\theta$ -curve
- $\det^*(M) = 0$  or  $\pm 2 \iff M$  represents handcuff graph


where  $\det^*(M)$  = determinant of  $H$ -deletion matrix of  $M$

## PROOF

$$\begin{bmatrix} 1 & 0 & 1 & 0 & 1 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 & 1 \\ 1 & 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 & 1 \\ 0 & 1 & 0 & 1 & 0 & 1 & 0 \end{bmatrix}$$



$H$ -deletion  $\rightarrow$




# Proof of Theorem

## CASE 1. When $M$ represents $\theta$ -curve

### i Line-shape

$$\begin{bmatrix} 0 & 1 & 1 & 0 & 0 & 0 & 1 \\ 1 & 0 & 1 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 & 1 \\ 0 & 1 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 1 & 0 \\ 1 & 0 & 0 & 0 & 1 & 0 & 0 \end{bmatrix}$$



$\xrightarrow{H\text{-deletion}}$

$$\begin{bmatrix} 1 & 1 & 0 & 0 & 1 \\ 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 1 & 0 & 1 \\ 1 & 0 & 0 & 1 & 0 \end{bmatrix}$$

$\xrightarrow[\text{operations}]{\text{row/column}}$

$$\begin{bmatrix} 1 & 0 & 0 & 0 & 0 \\ 1 & 1 & 0 & 0 & 0 \\ 0 & 1 & 1 & 0 & 1 \\ 0 & 0 & 1 & 1 & 0 \\ 0 & 0 & 0 & 1 & 0 \end{bmatrix}$$

$\xrightarrow{\text{subtracting}}$

$$\begin{bmatrix} 1 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 1 \\ 0 & 0 & 0 & 1 & -1 \\ 0 & 0 & 0 & 0 & 1 \end{bmatrix}$$

So  $\det^*(M) = \pm 1$