

The Arc Index of Theta-Curve and Handcuff Graph


Eunchan Cho¹ Jeongwon Shin¹ Boyeon Seo¹ Minho Choi¹

SEP 6, 2025

Introduction


Theta-Curves

- A **theta-curve** T is a graph embedded in S^3 , which consists of two vertices v_1, v_2 and three edges e_1, e_2, e_3 , such that each edge joins the vertices.
- A **constituent knot** T_{ij} , $1 \leq i < j \leq 3$, is a subgraph of T that consists of two vertices v_1, v_2 and two edges e_i, e_j .
- Theta-curves are roughly classified by comparing the triples of constituent knots.
- A theta-curve is said to be **trivial** if it can be embedded in a 2-sphere in S^3 .




Handcuff Graphs

- A **handcuff graph** H is a graph embedded in S^3 consisting of two vertices (v_1, v_2) and three edges (e_1, e_2, e_3), where e_3 has distinct endpoints v_1 and v_2 , and e_1 and e_2 are loops based at v_1 and v_2 .
- A **constituent link** H_{12} , is a subgraph of H that consists of two vertices v_1, v_2 and two edges e_1, e_2 .



Reidemeister Moves for Theta-Curves and Handcuff Graphs



- **Arc presentation** of a theta-curve or handcuff graph is an embedding of them.
- It is contained in the union of finitely many half planes (called **pages**).
- The embedding is with the common boundary line (called **axis**).
- Each vertex lies in the axis.
- Each page contains a properly embedded single arc.
- **Arc index**, is the minimal number of pages among all possible arc presentations of graph.
- This arc presentation with the minimal number of pages is **minimal arc presentation**.

Arc Presentation



Trefoil




Open Book



Grid Diagram



$\theta_{5,2}$



Open Book



Grid Diagram



2_1

$\Phi_{2,1}$



Open Book





Grid Diagram

Grid Diagram

- The **grid diagram** of theta-curve or handcuff graph is a diagram with only vertical strand and horizontal strands.
- $(\text{number of vertical strands}) + 1 = (\text{number of horizontal strands})$
- At every crossing, the vertical strand crosses over the horizontal strand.
- No two horizontal strands are in the same row.
- No two vertical strands are in same column.

Grid Diagram

- A grid diagram gives rise to an arc presentation and vice versa.





Arc Presentation of the Theta-Curve and Handcuff Graph

Theorem

Every theta-curve and handcuff graph admit a grid diagram.

PROOF




Corollary

Every theta-curve and handcuff graph admit a arc presentation.

Classifying by Determinant

THC-cromwell matrix

- The **THC-cromwell matrix** is an expansion of cromwell matrix into θ -curves and handcuff graphs



$$\begin{bmatrix} 1 & 1 & 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 & 0 & 1 \\ 1 & 0 & 0 & 0 & 1 & 0 \\ 0 & 1 & 1 & 0 & 0 & 1 \end{bmatrix}$$

Determinant of the cromwell matrices of Knot


Theorem

Let K be any knot then its determinant of the cromwell matrix is 0 or ± 2 .

PROOF



grid diagram



THC-cromwell

THC-cromwell

$$\begin{bmatrix} 1 & 0 & 1 & 0 & 0 \\ 0 & 1 & 0 & 1 & 0 \\ 0 & 0 & 1 & 0 & 1 \\ 1 & 0 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 & 1 \end{bmatrix}$$

row/column
operations

$$\begin{bmatrix} 1 & 1 & 0 & 0 & 0 \\ 0 & 1 & 1 & 0 & 0 \\ 0 & 0 & 1 & 1 & 0 \\ 0 & 0 & 0 & 1 & 1 \\ 1 & 0 & 0 & 0 & 1 \end{bmatrix}$$

CASE 1. When n is an even number.

$$\begin{bmatrix} 1 & 1 & 0 & 0 \\ 0 & 1 & 1 & 0 \\ 0 & 0 & 1 & 1 \\ 1 & 0 & 0 & 1 \end{bmatrix} \mapsto \begin{bmatrix} 1 & 1 & 0 & 0 \\ 0 & 1 & 1 & 0 \\ 0 & 0 & 1 & 1 \\ 2 & 2 & 2 & 2 \end{bmatrix} \mapsto \begin{bmatrix} 1 & 1 & 0 & 0 \\ 0 & 1 & 1 & 0 \\ 0 & 0 & 1 & 1 \\ 0 & 0 & 0 & 0 \end{bmatrix}$$

So the determinant of K is 0.

CASE 2. When n is an odd number.

$$\begin{bmatrix} 1 & 1 & 0 & 0 & 0 \\ 0 & 1 & 1 & 0 & 0 \\ 0 & 0 & 1 & 1 & 0 \\ 0 & 0 & 0 & 1 & 1 \\ 1 & 0 & 0 & 0 & 1 \end{bmatrix} \mapsto \begin{bmatrix} 1 & 1 & 0 & 0 & 0 \\ 0 & 1 & 1 & 0 & 0 \\ 0 & 0 & 1 & 1 & 0 \\ 0 & 0 & 0 & 1 & 1 \\ 2 & 2 & 2 & 2 & 2 \end{bmatrix} \mapsto \begin{bmatrix} 1 & 1 & 0 & 0 & 0 \\ 0 & 1 & 1 & 0 & 0 \\ 0 & 0 & 1 & 1 & 0 \\ 0 & 0 & 0 & 1 & 1 \\ 0 & 0 & 0 & 0 & 2 \end{bmatrix}$$

So the determinant of K is ± 2 .

□

H-deletion of THC-cromwell matrices

- The **H-deletion** Matrix of the THC-cromwell matrix G is $(n - 1) \times (n - 1)$ matrix which deleted vertex-row and its two side-rows from the matrix G .

$$\begin{bmatrix} 1 & 0 & 1 & 0 & 0 & 1 \\ 0 & 1 & 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 & 1 & 0 \\ 1 & 0 & 0 & 1 & 0 & 0 \\ 0 & 1 & 0 & 0 & 1 & 1 \end{bmatrix} \mapsto \begin{bmatrix} 1 & 0 & 1 & 0 & 0 & 1 \\ 0 & 1 & 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 & 1 & 0 \\ 1 & 0 & 0 & 1 & 0 & 0 \\ 0 & 1 & 0 & 0 & 1 & 1 \end{bmatrix} \mapsto \begin{bmatrix} 1 & 0 & 1 & 0 \\ 0 & 1 & 0 & 1 \\ 0 & 0 & 1 & 0 \\ 1 & 0 & 0 & 1 \end{bmatrix}$$

Determinant of the THC-cromwell matrices

Theorem



Let M be any THC-cromwell matrix of θ -curve or handcuff graph.

- $\det(M) = \pm 1 \iff M$ represents θ -curve
- $\det(M) = 0$ or $\pm 2 \iff M$ represents handcuff graph

* $\det(M)$ = determinant of H -deletion matrix of M

PROOF


$$\begin{bmatrix} 1 & 0 & 1 & 0 & 1 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 & 1 \\ 1 & 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 & 1 \\ 0 & 1 & 0 & 1 & 0 & 1 & 0 \end{bmatrix}$$



CASE 1. When M represents θ -curve

① Line-shape

$$\begin{pmatrix} 0 & 1 & 1 & 0 & 0 & 0 & 1 \\ 1 & 0 & 1 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 & 1 \\ 0 & 1 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 1 & 0 \\ 1 & 0 & 0 & 0 & 1 & 0 & 0 \end{pmatrix}$$



$\xrightarrow{H\text{-deletion}}$

$$\begin{bmatrix} 1 & 1 & 0 & 0 & 1 \\ 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 1 & 0 & 1 \\ 1 & 0 & 0 & 1 & 0 \end{bmatrix}$$

and simplify with

$\xrightarrow[\text{operations}]{\text{row/column}}$

$$\begin{bmatrix} 1 & 0 & 0 & 0 & 0 \\ 1 & 1 & 0 & 0 & 0 \\ 0 & 1 & 1 & 0 & 1 \\ 0 & 0 & 1 & 1 & 0 \\ 0 & 0 & 0 & 1 & 0 \end{bmatrix}$$

$\xrightarrow{\text{subtracting}}$


$$\begin{bmatrix} 1 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 1 \\ 0 & 0 & 0 & 1 & -1 \\ 0 & 0 & 0 & 0 & 1 \end{bmatrix}$$

So $\det(M) = \pm 1$

CASE 1. When M represents θ -curve

ii) T-shape

$$\begin{bmatrix} 1 & 0 & 1 & 0 & 1 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 & 1 \\ 1 & 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 & 1 \\ 0 & 1 & 0 & 1 & 0 & 1 & 0 \end{bmatrix}$$



H -deletion

$$\begin{bmatrix} 1 & 0 & 0 & 0 & 1 \\ 0 & 0 & 1 & 0 & 0 \\ 0 & 1 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 1 \\ 1 & 0 & 1 & 1 & 0 \end{bmatrix}$$

and simplify with

row/column
operations

$$\begin{bmatrix} 1 & 0 & 0 & 0 & 0 \\ 1 & 1 & 0 & 1 & 0 \\ 0 & 1 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 & 1 \\ 0 & 0 & 0 & 0 & 1 \end{bmatrix}$$



regioning

$$\begin{bmatrix} 1 & 0 & 0 & 0 & 0 \\ 1 & 1 & 0 & 1 & 0 \\ 0 & 1 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 & 1 \\ 0 & 0 & 0 & 0 & 1 \end{bmatrix}$$

So $\det(M) = \pm 1$

CASE 2. When M represents handcuff graph

$$\begin{bmatrix} 0 & 0 & 1 & 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 & 1 \\ 1 & 0 & 1 & 0 & 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 & 1 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 & 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 1 & 0 & 1 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 1 & 0 & 0 \end{bmatrix}$$



CASE 2. When M represents handcuff graph


T-loop

$$\begin{bmatrix} 0 & 0 & 1 & 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 & 1 \\ 1 & 0 & 1 & 0 & 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 & 1 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 & 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 1 & 0 & 1 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 1 & 0 & 0 \end{bmatrix}$$

and simplify with

$$\begin{array}{c} \text{row/column} \\ \hline \text{operations} \end{array} \rightarrow \begin{bmatrix} 1 & 1 & 0 \\ 1 & 1 & 1 \\ 0 & 0 & 1 \end{bmatrix}$$

So $\det(M) = 0$ or ± 2



CASE 2. When M represents handcuff graph

① T-loop & Line-shape

$$\begin{bmatrix} 0 & 0 & 1 & 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 & 1 \\ 1 & 0 & 1 & 0 & 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 & 1 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 & 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 1 & 0 & 1 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 1 & 0 & 0 \end{bmatrix}$$

and simplify with


operations \rightarrow

$$\begin{bmatrix} T-loop & * \\ 0 & Line-shape \end{bmatrix}$$

seperating \rightarrow

$$\begin{bmatrix} Knot & * & * \\ 0 & Line-shape & * \\ 0 & 0 & Line-shape \end{bmatrix}$$

So $\det(M) = 0$ or ± 2



H -deletion \rightarrow

$$\begin{bmatrix} 0 & 1 & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 & 0 & 1 \\ 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 1 & 0 & 0 \\ 1 & 0 & 1 & 0 & 1 & 0 \\ 1 & 0 & 0 & 0 & 1 & 0 \end{bmatrix}$$

CASE 2. When M represents handcuff graph

② Knot & Line-shape

cromwell matrix $\xrightarrow{H\text{-deletion}}$

H-deletion
matrix

$\xrightarrow[\text{operations}]{\text{seperating}}$

$$\begin{bmatrix} \text{Knot} & * \\ 0 & \text{Line - shape} \end{bmatrix}$$

So $\det(M) = 0$ or ± 2

