

The Determinant and Arc Indices of θ -Curves and Handcuff-Graphs


Eunchan Cho¹ Jeongwon Shin¹ Boyeon Seo¹ Minho Choi¹

NOV 22, 2025
R&E 2025

Introduction


θ -Curves

- A **θ -curve** T is a graph embedded in S^3 , which consists of two vertices v_1, v_2 and three edges e_1, e_2, e_3 , such that each edge joins the vertices.
- A **constituent knot** T_{ij} , $1 \leq i < j \leq 3$, is a subgraph of T that consists of two vertices v_1, v_2 and two edges e_i, e_j .
- θ -curves are roughly classified by comparing the triples of constituent knots.
- A θ -curve is said to be **trivial** if it can be embedded in a 2-sphere in S^3 .




Handcuff Graphs

- A **handcuff graph** H is a graph embedded in S^3 consisting of two vertices (v_1, v_2) and three edges (e_1, e_2, e_3), where e_3 has distinct endpoints v_1 and v_2 , and e_1 and e_2 are loops based at v_1 and v_2 .
- A **constituent link** H_{12} , is a subgraph of H that consists of two vertices v_1, v_2 and two edges e_1, e_2 .



Reidemeister Moves for θ -Curves and Handcuff Graphs



- **Arc presentation** of a θ -curve or handcuff graph is an embedding of them.
- It is contained in the union of finitely many half planes (called **pages**).
- The embedding is with the common boundary line (called **axis**).
- Each vertex lies in the axis.
- Each page contains a properly embedded single arc.
- **Arc index**, is the minimal number of pages among all possible arc presentations of graph.
- This arc presentation with the minimal number of pages is **minimal arc presentation**.

Arc Presentation



Trefoil




Open Book



Grid Diagram



$\theta_{5,2}$



Open Book




Grid Diagram



2_1

$\Phi_{2,1}$



Open Book





Grid Diagram

Grid Diagram

- The **grid diagram** of θ -curve or handcuff graph is a diagram with only vertical and horizontal strands.
- $(\text{number of vertical strands}) + 1 = (\text{number of horizontal strands})$
- At every crossing, the vertical strand crosses over the horizontal strand.
- No two horizontal strands are in the same row.
- No two vertical strands are in same column.

Grid Diagram

- A grid diagram gives rise to an arc presentation and vice versa.





Arc Presentation of the θ -Curve and Handcuff Graph

Theorem

Every θ -curve and handcuff graph admit a grid diagram.

PROOF



Corollary

Every θ -curve and handcuff graph admit a arc presentation.

Lower and Upper Bounds of Arc Index





Lower Bounds from Constituent Knots

Theorem

Let T be any θ -curve and K_1, K_2, K_3 be three constituent knots of T . Then

$$\alpha(T) \geq \max_{i \in \{1,2,3\}} \alpha(K_i) + 1$$

PROOF



Theorem

Let T be any θ -curve and K_1, K_2, K_3 be three constituent knots of T . Then

$$\alpha(T) \geq \frac{1}{2} \sum_{i=1}^3 \alpha(K_i)$$


PROOF

- A minimal arc presentation of T is given.
- $K_1 = e_1 \cup e_2$, $K_2 = e_2 \cup e_3$, and $K_3 = e_3 \cup e_1$.
- S_i be the set of half plane corresponding the edge e_i .
- $S_i \cup S_{i+1}$ form an arc presentation of the knot K_i .
- $\alpha(K_i) \leq |S_i| + |S_{i+1}|$


$$\sum_{i=1}^3 \alpha(K_i) \leq 2 \sum_{i=1}^3 |S_i| = 2\alpha(T)$$



Stacked Tangle of an θ -Curve



Stacked Tangle of a Link




Stacked Tangle of a θ -Curve

Stacked Tangle of an θ -Curve

Stacked tangle of an θ -curve is stacked disks each with the frame as boundary with following properties:

- Only two disk called **non-simple disks** contain one vertex and three line segments which joins the vertex and boundary point.
- One of the non-simple discs is at the top.
- Other disks called **simple disks** contain simple arc which joins two points on the boundary.
- When view from above
 - two arcs in different simple disks intersect at most one point(by RII)
 - arc in simple disk and tree in non-simple disk intersect at most one point(by RV)




Stacked Tangle of an θ -Curve

Simple closure of stacked tangle is a **stacked tangle** with **caps** satisfying following properties:

- A **cap** is a simple arc in outside of stacked tangle joining end points of arcs or line segments.
- When view from above any tow caps have no intersection.

Then a simple closure of a stacked tangle **without any nested caps** is corresponding to an arc presentation.



A **reduced simple closure of a stacked tangle** is




- a simple closure of a stacked tangle **without any nested caps**
- any two arcs(including line segment) joining by caps have **no intersection** when view from above

Stacked Tangle of an θ -Curve

Proposition

A reduced simple closure of a stacked tangle can be obtained a simple closure of a stacked tangle without any nested caps by applying Reidemaister Moves.

PROOF



Yamada Polynomials

Let D_T be a diagram of an θ -curve T . Then, the **Yamada Polynomial** $R(D_T) \in \mathbb{Z}[x^{\pm 1}]$ is calculated by the following properties:

- **Y6:** $R(\bigoplus) = -(x + 1 + x^{-1})(x + x^{-1}) = -x^2 - x - 2 - x^{-1} - x^{-2}$ **Y7:** $R(\bigcirc \text{---} \bigcirc) = 0$
- **Y8:** $R(T' \cup \bigcirc) = (x + 1 + x^{-1})R(T')$ for an arbitrary θ -curve diagram T'
- **Y9:** $R(\bigwedge) - R(\bigvee) = (x - x^{-1}) [R(\bigcirc \bigcirc) - R(\text{---})]$
- **Y10:** $R(\bigcirc) = x^2 R(\bigcap), \quad R(\bigcirc) = x^{-2} R(\bigcap)$
- **Y11:** $R(\bigcirc \bigcirc) = R(\bigcirc \bigcirc)$ **Y12:** $R(\bigwedge \bigvee) = R(\bigvee \bigwedge)$
- **Y13:** $R(\bigwedge \bigvee) = R(\bigvee \bigwedge), \quad R(\bigwedge \bigvee) = R(\bigvee \bigwedge)$
- **Y14:** $R(-\bigcirc) = -x R(\bigcirc), \quad R(-\bigcirc) = -x^{-1} R(\bigcirc)$

Proposition ([?])

$R(D_T)$ is an ambient isotopy invariant of T up to multiplying $(-x)^n$ for some integer n .

Theorem

Let T be any θ -curve or handcuff graph. Then

$$2 + \sqrt{\max \deg_x R(S_T) - \min \deg_x R(S_T) - 4} \leq \alpha(T)$$

where $R(T)$ is a Yamada Polynomial of T .

Lower Bounds from Yamada Polynomial

Proposition


Let S_T be a simple closure of stacked tangle of a θ -curve or handcuff graph T **without nested caps**. Then

$$\max \deg_x R(S_T) \leq c + n, \quad \min \deg_x R(S_T) \geq -(c + n),$$

where **c, n** is the number of caps and crossings in S_T , respectively.

PROOF

- Use double mathematical induction of $(c_s + c_{ss}, n)$.



Basis Step:

- If $c_S + c_{SS} = 0$, then S_T has no simple disks and is equivalent to the result of applying Y14 to \bigoplus .
 $\therefore R(S_T) = -x^{\pm 3} [-x^2 - x - 2 - x^{-1} - x^{-2}] \implies 5 \leq c + n.$
- If $n = 0$, then S_T is equivalent to $\bigcirc - \bigcirc \cup \bigcirc \cup \dots \cup \bigcirc$.
 $\therefore R(S_T) = 0 \implies 0 < 2 \leq c + n.$

All base cases satisfy the inequality.


Proof of Theorem

Inductive Step:

Assume that it holds for any $(c'_s + c'_{ss}, n') < (c_s + c_{ss}, n)$, and $c_s + c_{ss} > 0$.

Let S_T be a **simple closure of stacked tangle** of a θ -curve or handcuff graph T such that the number of simple caps, semi-simple caps, and crossings are c_s, c_{ss}, n , respectively.

Take the topmost **simple disk** D_s connected to the top disk, and a **disk** D directly above D_s .



Proof of Theorem

CASE 1. Suppose that there is no cap between D_S and D .

① Suppose that there is no intersection between D_S and D in S_T .

- D_S and D do not affect each other.
- We can swap the position of D_S and D without affecting the rest of the diagram.

② Suppose that there is an intersection between D_S and D in S_T .

- Let S_T^- , S_T^0 and S_T^∞ be the simple closure of stacked tangle which is obtained by replacing \times with \times , $\rangle\langle$ and \asymp , respectively.
- The simple caps, semi-simple caps, and crossings of the both are $c_s, c_{ss}, n - 1$.
- Applying Y9

$$R(\times) - R(\times) = (x - x^{-1}) [R(\rangle\langle) - R(\asymp)],$$

then

$$R(S_T) - R(S_T^-) = (x - x^{-1})(R(S_T^0) - R(S_T^\infty)).$$


- Then, it is sufficient to show that the interchanged one holds.

Proof of Theorem


CASE 2. Suppose that there is a cap between D_s and D .

① Suppose that D is a simple disk.

- When view from above, there are three cases:



- After applying **Y10**, the second and third cases can be regarded as the first case, and the cap can be reduced.



- S'_T has $c - 1$ caps, $c - 1$ simple caps, $c - 1$ semi-simple caps and $n - 1$ crossings

- By induction hypothesis,

$$\begin{aligned}\max \deg_x R(S_T) &= \max \deg_x R(S'_T) \pm 2 \\ &\leq [(c-1) + (n-1)] \pm 2 \\ &\leq c + n, \\ \min \deg_x R(S_T) &= \min \deg_x R(S'_T) \pm 2 \\ &\geq -[(c-1) + (n-1)] \pm 2 \\ &\geq -(c+n).\end{aligned}$$

② D is not a simple disk.

- When viewed from above, all the cases can be reduced as follows.



- $R(S_T) = -x^{\pm 1}R(S'_T)$ and $R(S_T) = x^{\pm 2}R(S''_T)$ by **Y14** and **Y10**, respectively.
- Both of S'_T and S''_T have $c - 1$ caps, c_s simple caps, $c_{ss} - 1$ semi-simple caps, and $n - 1$ crossing.

Proof of Theorem

- By induction hypothesis, in the first case,

$$\begin{aligned}\max \deg_x R(S_T) &= \max \deg_x R(S'_T) \pm 1 \\ &\leq [(c-1) + (n-1)] \pm 1 \\ &\leq c + n.\end{aligned}$$

- Similarly, in the second case,

$$\begin{aligned}\max \deg_x R(S_T) &= \max \deg_x R(S''_T) \pm 2 \\ &\leq [(c-1) + (n-1)] \pm 2 \\ &\leq c + n.\end{aligned}$$

- It holds for $\min \deg_x R(S_T)$ in the same way.



Proposition

Let S_T be a reduced simple closure of stacked tangle of a θ -curve or handcuff graph T corresponding to minimal arc presentation of T . Then

$$\max \deg_x R(S_T) - \min \deg_x R(S_T) - 2n \leq \alpha(T)$$


where n is the number of crossings in S_T .

PROOF

- S_T is a reduced simple closure of stacked tangle corresponding to minimal arc presentation.
- The number of caps c in S_T is exactly arc index of T , $\alpha(T)$.

Proof of Theorem

- Take a cap and add a positive or negative curl



- After modification of diagram as above, resulting diagram is also a simple closure of stacked tangle.
- The number of crossings is increased by 1.
- p of the caps yield a negative curl, and the remaining $c - p$ yield a positive curl.
- $S_T^{neg}(S_T^{pos})$ is the diagram obtained by inserting the p negative($c - p$ positive) curls.

Proof of Theorem

	S_T^{neg}	S_T^{pos}
Number of Caps	c	c
Number of Crossings	$n + p$	$n + (c - p)$

$$\bullet R(S_T^{neg}) = x^{-2p}R(S_T) \text{ and } R(S_T^{pos}) = x^{2(c-p)}R(S_T)$$

$$\begin{aligned} \min \deg_x R(S_T) - 2p &= \min \deg_x R(S_T^{neg}) \\ &\geq -c + -(n + p) \end{aligned}$$

$$\begin{aligned} \max \deg_x R(S_T) + 2(c - p) &= \max \deg_x R(S_T^{pos}) \\ &\leq c + [n + (c - p)] \end{aligned}$$

$$\min \deg_x R(S_T) \geq -c - n + p$$

$$\max \deg_x R(S_T) \leq n + p$$

$$\max \deg_x R(S_T) - \min \deg_x R(S_T) \leq c + 2n$$



Proof of Theorem

Theorem

Let T be any θ -curve or handcuff graph. Then

$$2 + \sqrt{\max \deg_x R(S_T) - \min \deg_x R(S_T) - 4} \leq \alpha(T)$$

where $R(T)$ is a Yamada Polynomial of T .

PROOF

Let S_T be a reduce simple closure of stacked tangle of a θ -curve or handcuff graph T corresponding to minimal arc presentation of T .

- The number of caps : $\alpha(T)$
- The number of non-simple disks : 2
- The number of simple disks : $\alpha(T) - 3$

Proof of Theorem

① Let T be any θ -curve.

Consider the maximum number of crossings in S_T .

- number of crossings by two simple disks : $\binom{\alpha(T)-3}{2} = \frac{1}{2} (\alpha(T) - 3) (\alpha(T) - 4)$
- number of crossings by a simple disk and non-simple disk : $2 (\alpha(T) - 3)$
- number of crossings by two non-simple disks : 2
- number of crossings counted by disks joined by cap : $\alpha(T) - 2$

Thus

$$\begin{aligned} n &\leq \frac{1}{2} (\alpha(T) - 3) (\alpha(T) - 4) + 2 (\alpha(T) - 3) + 2 - (\alpha(T) - 2) \\ &= \frac{1}{2} [(\alpha(T))^2 - 5\alpha(T) + 8] \end{aligned}$$

By Lemma,

$$\begin{aligned} \max \deg_x R(S_T) - \min \deg_x R(S_T) &\leq 2n + \alpha(T) \leq \alpha(T)^2 - 4\alpha(T) + 8 \\ 2 + \sqrt{\max \deg_x R(S_T) - \min \deg_x R(S_T) - 4} &\leq \alpha(T) \end{aligned}$$

Proof of Theorem

② Let T be any handcuff graph.

Consider the maximum number of crossings in S_T .

Proof of Theorem

- number of crossings by two simple disks : $\binom{\alpha(T)-3}{2} = \frac{1}{2} (\alpha(T) - 3) (\alpha(T) - 4)$
- number of crossings by a simple disk and non-simple disk : $2 (\alpha(T) - 3)$
- number of crossings by two non-simple disks : 1
- number of crossings counted by disks joined by cap : $\alpha(T) - 1 - 2 = \alpha(T) - 3$

Thus

$$\begin{aligned} n &\leq \frac{1}{2} (\alpha(T) - 3) (\alpha(T) - 4) + 2 (\alpha(T) - 3) + 1 - (\alpha(T) - 3) \\ &= \frac{1}{2} [(\alpha(T))^2 - 5\alpha(T) + 8] \end{aligned}$$

By Lemma,

$$\max \deg_x R(S_T) - \min \deg_x R(S_T) \leq 2n + \alpha(T) \leq \alpha(T)^2 - 4\alpha(T) + 8$$

Result

Grid Diagrams of θ -Curves



Grid Diagrams of Handcuff-Graphs



Further Research

Further Research

- We tried to use the Python for determining arc indices. For the Python program, we used the Topoly package, but the package had an error. We would like to use another tool to completely determine the arc indices of θ -curves and handcuff-graphs, such as Yamada package.
- Applying the result of our research to θ -curves or handcuff-graphs of higher crossings or to other spatial graphs can be researched further.