

Mini-course:  
Numerical Methods for Mathematical Finance  
Projects list

Silvia Lavagnini

May 23, 2024

- You can choose one project from the list below, to be solved in Python.
- Whenever not specified, you can use the same model parameters as seen in class, or choose your own.
- For the course *Numerical Methods for Mathematical Finance* (6 credits): it is required to solve Problem 1, 3 and one between Problem 4 and Problem 5.

1. **The Milstein scheme for SDE simulations** (requires stochastic calculus)

- (a) Derive, via Itô-Taylor expansion, the Milstein Scheme;
- (b) Compose the code for the Milstein scheme to simulate an Ornstein-Uhlenbeck process of the form

$$dX(t) = \theta(\mu - X(t))dt + \sigma dW(t),$$

for  $\mu \in \mathbb{R}$ ,  $\theta > 0$  and  $\sigma > 0$ ;

- (c) For  $\mu = 10$ ,  $\theta = 1$  and  $\sigma = 0.5$ , apply the scheme for  $\Delta t = 2^{-2}$  and  $\Delta t = 2^{-4}$  and plot the results against the exact simulation with  $\Delta t = 2^{-9}$ ;
- (d) For different values of  $\Delta t$ , compute the weak and the strong error for the scheme. Plot the results against  $\Delta t$ .

2. **Pricing vanilla options under the B&S model via PDE solver**

Consider the PDE for pricing a European call option under the Black and Scholes model:

- (a) Compose the code for a finite difference method to solve the PDE;
- (b) Choose several strike values  $K$ , and apply the scheme with temporal grid size  $\Delta t = 0.05$  and spatial grid size  $\Delta S = 1$ . Plot the result against the exact solution (Black and Scholes formula);
- (c) Repeat the experiments for different values of  $\Delta t$  (fixed  $\Delta S$ ) and plot the relative error at initial time against  $\Delta t$ . Repeat again by, this time, changing the spatial grid size  $\Delta S$  (fixed  $\Delta t$ ).

3. **Pricing vanilla options under the Ornstein-Uhlenbeck model with the COS method**

- (a) Consider an Ornstein-Uhlenbeck process of the form

$$dX(t) = \theta(\mu - X(t))dt + \sigma dW(t),$$

for  $\mu \in \mathbb{R}$ ,  $\theta > 0$  and  $\sigma > 0$ , and derive its characteristic function;

- (b) Compose the code for the COS method to price call options under the Ornstein-Uhlenbeck model;
- (c) For  $\mu = 30$ ,  $\theta = 1$  and  $\sigma = 0.5$ , apply the COS method for  $N = 16$  and  $N = 32$  for different strike values, and plot the results against the "exact solution" obtained by the same method with  $N = 5000$ ;
- (d) Repeat the approximation for different truncation parameters  $N$ . Compute and plot the maximum absolute error against  $N$ .

#### 4. Pricing Asian option via Monte Carlo

Consider an arithmetic Asian option with call-type payoff function of the form

$$\max\left(\frac{1}{T}\int_0^T S(t)dt - K, 0\right),$$

and consider a GMB for the underlying process  $S$ :

- (a) Use a Monte Carlo method with Euler scheme to approximate the option value;
- (b) Apply the scheme for  $\Delta t = 2^{-2}$  and  $\Delta t = 2^{-4}$  and plot the results against the "exact simulation" obtained by the same method with  $\Delta t = 2^{-9}$  and  $N = 10000$ ;
- (c) For different values of  $\Delta t$ , compute the relative error at initial time for the scheme. Plot the results against  $\Delta t$ .

#### 5. Pricing of options depending on two underlying processes

Let  $\mathbf{X}(t) = (X(t), Y(t))^\top$  where  $X$  satisfies the following SDE under the pricing measure  $\mathbb{Q}$

$$dX(t) = 0.06X(t)dt + 0.38X(t)dW^1(t),$$

and  $Y$  satisfies under the pricing measure  $\mathbb{Q}$

$$dY(t) = 0.06Y(t)dt + 0.15Y(t)dW^2(t),$$

where  $W^1$  and  $W^2$  are two correlated Brownian motion with correlation  $\rho = 0.7$ . Consider the option with strike price  $K$  defined by

$$V(t, \mathbf{X}) = \mathbb{E}^{\mathbb{Q}} \left[ e^{-r(T-t)} \max\left(\frac{1}{2}X(T) - \frac{1}{2}Y(T), K\right) \middle| \mathcal{F}_t \right].$$

- (a) Consider  $r = 0.06$  and  $T = 5$ . Use a Monte Carlo method with Euler scheme to approximate the option value at time  $t = 0$  for strike values  $K = 0$  to  $K = 10$ , with interval  $\Delta K = 0.5$ . Plot the results againsts  $K$ ;
- (b) Apply the scheme for  $\Delta t = 2^{-2}$  and  $\Delta t = 2^{-4}$  and plot the results against the "exact simulation" obtained by the same method with  $\Delta t = 2^{-9}$  and  $N = 10000$ ;
- (c) For different values of  $\Delta t$ , compute the relative error at initial time for the scheme. Plot the results against  $\Delta t$ .