

Signal Hierarchy Theory for Biological Petri Nets: Formal Semantics of Hierarchical Regulatory Control

Eugênio Simão^{1,*}

¹ Department of Computer Science, Universidade Federal de Santa Catarina (UFSC), Araranguá, Santa Catarina, 88906-072, Brazil

* eugenio.simao@ufsc.br

Abstract

Predicting cellular commitment thresholds—such as ATP concentrations triggering irreversible developmental transitions—requires modeling regulatory signal consumption during decision-making. Biological Petri net theory lacks formal semantics for such hierarchical regulatory signals. Classical test arcs model catalysis (non-consuming reads) but cannot express signal consumption essential for commitment finality and pathway preemption (Table 1). We introduce *signal hierarchy theory*, a formal extension to Bio-PN providing signal places as information channels distinct from metabolic mass transfer, with signal flow arcs enabling consumption semantics. Hierarchical layer assignment formalizes multi-scale organization where higher regulatory layers control lower metabolic layers through regulatory consumption. The formalism bridges the expressiveness gap between metabolic mass transfer (normal arcs) and regulatory information flow (signal flow arcs), enabling unified modeling where energy status functions simultaneously as metabolite and regulatory signal. We provide complete formal definitions of the 13-tuple extension (Eq. 5, Table 2) including signal places (Ψ), signal flow arc semantics, hierarchical layer functions, and enablement thresholds. We demonstrate the formalism’s applicability through *Bacillus subtilis* sporulation, an illustrative example: signal hierarchy analysis computes ATP commitment threshold at 2.38 mM, matching experimental measurements (2.21 mM, 7% error)—a prediction inexpressible in classical Bio-PN lacking consumption semantics. This proof-of-concept demonstrates how the formalism enables quantitative threshold determination, basin geometry quantification, and commitment reliability analysis, providing theoretical foundation for modeling irreversible cellular decisions under energy constraints.

Keywords: Systems Biology, Cellular Decision-Making, Metabolic Networks, Quantitative Modeling, Cell Fate, Biological Petri Nets

Author Summary

Cells make irreversible decisions—like committing to division, differentiation, or death—based on energy availability. Current mathematical models

treat energy molecules like ATP simply as passive participants in reactions, missing their role as active regulatory signals that determine which pathways activate. We developed a new theoretical framework that formally distinguishes between ATP as a metabolite (consumed in reactions) and ATP as a regulatory signal (controlling pathway choice). Our theory introduces “signal flow arcs” that consume regulatory information during commitment, making decisions irreversible and enabling higher-level pathways to preempt lower-level ones. We demonstrate this framework by analyzing bacterial sporulation, a life-or-death decision where ATP availability determines cell fate. Our mathematical model correctly predicted the ATP threshold for commitment within 7% of experimental values—a calculation impossible with existing models that cannot represent signal consumption. This work provides new mathematical machinery for understanding how cells make reliable decisions under changing energy conditions. The formalism has potential applications in synthetic biology (designing robust decision circuits), drug discovery (understanding ATP-dependent pathways), and precision medicine (predicting cellular responses to metabolic interventions).

1 Introduction

1.1 Motivation

Cellular decision-making requires integrated models capturing metabolic and regulatory information flow. Biological systems exhibit hierarchical organization where metabolic state (e.g., energy status) controls regulatory networks (e.g., gene expression), which in turn modulate metabolic flux. This bidirectional coupling demands unified formalisms treating metabolism and regulation as inseparable components of cellular control, not separate modeling domains. For instance, ATP in *Bacillus subtilis* simultaneously fuels metabolism (mass transfer) and gates developmental commitment (regulatory signal)—when ATP falls below 2.21 mM, cells irreversibly commit to sporulation, a threshold current formalisms cannot compute.

Biological Petri nets [1, 2] extended classical formalism with continuous places for metabolites, stochastic semantics for molecular noise, and test arcs for catalysis. However, a critical limitation remains: metabolism and regulation are separated. Normal arcs model mass transfer (substrates consumed, products generated), while test arcs model catalytic reads (enzymes unchanged). This dichotomy cannot capture regulatory molecules *consumed* during decision-making—signal consumption that collapses decision spaces and ensures commitment finality. Growth factor binding depletes extracellular signal, ATP expenditure in *B. subtilis* sporulation gates developmental commitment, and similar consumption occurs in lambda phage CI repressor binding. Recent advances address resource constraints [3], but no framework provides formal hierarchical information channels with consumption semantics necessary for modeling integrated metabolic-regulatory control.

1.2 Biological Motivation for Signal Hierarchy

Cellular regulation exhibits multi-scale hierarchical organization: **Stress response**—Environmental signals (UV damage, heat shock) trigger de-

cision points that reconfigure metabolic pathways through hierarchical preemption where regulatory layers control metabolic execution. **Energy status**—ATP/ADP ratio serves as a master regulatory signal, hierarchically controlling anabolic versus catabolic flux through multiple pathways simultaneously. **Developmental decisions**—Nutrient depletion in *Bacillus subtilis* triggers ATP-gated commitment to sporulation versus competence pathways, exemplifying metabolic state controlling gene regulatory networks. **Quorum sensing**—Autoinducer concentration represents population-level information that preempts individual cell decision-making, enabling coordinated multicellular behavior.

Existing Bio-PNs model catalysis via test arcs (non-consuming reads). However, regulatory signals are often *consumed* during decision-making, changing system state irreversibly. For example, ATP consumption in *B. subtilis* sporulation creates distinct attractor basins for developmental commitment, and growth factor binding to receptors depletes extracellular signal. Lambda phage provides another example where CI repressor binding to operators consumes regulatory signal concentration. We formalize signal places (Ψ) with signal flow arcs that consume tokens, enabling hierarchical preemption mechanisms absent in current Bio-PN theory.

1.3 Contributions

This work makes two primary contributions:

(1) Signal Hierarchy Theory—We introduced signal places (Ψ) as formal information channels distinct from metabolic mass transfer. Signal flow arcs consume tokens (unlike test arcs which read without consumption), enabling hierarchical preemption where higher regulatory layers control lower metabolic layers. The formalism provides: (i) Signal place partition ($\Psi \subseteq P$) distinguishing regulatory information channels from metabolic places, (ii) Signal flow arc semantics with consumption, (iii) Hierarchical layer assignment ($L : \Psi \rightarrow \mathbb{N}$) formalizing multi-scale organization, (iv) Preemption mechanism where higher-layer signal consumption collapses decision spaces at lower layers. This bridges the fundamental gap between metabolic networks (mass transfer) and regulatory networks (information flow) within unified Bio-PN formalism.

(2) Proof-of-Concept Application—We demonstrated the formalism’s applicability through *Bacillus subtilis* sporulation, a canonical ATP-gated developmental decision. Signal hierarchy analysis computes ATP commitment threshold at 2.38 mM, matching experimental measurements (2.21 mM, 7% error). This quantitative prediction is inexpressible in classical Bio-PN lacking consumption semantics for regulatory signals. Similar analytical results were obtained for other biological systems (lambda phage lysis-lysogeny, MAPK signaling cascades), confirming the approach’s generalizability. The application illustrates: *Threshold determination*—Computing energy concentrations triggering pathway commitment, *Basin geometry quantification*—Reachability analysis quantifying cell fate reliability and attractor convergence, *Decision finality analysis*—Formal distinction between irreversible commitment (signal consumption) and reversible regulation (test arcs). This illustrative example establishes proof-of-concept for the formalism’s predictive capability and demonstrates advantages over classical Bio-PN approaches.

Together, these contributions provide theoretical foundation for a new class of analyses about commitment reliability, basin geometry, and decision finality that lack formal semantics in classical Bio-PN theory.

1.4 Organization

Section 2 reviews classical Petri net theory, biological Petri nets, and identifies the expressiveness gap. Section 3 develops signal hierarchy theory with formal semantics. Section 4 presents the unified formalism integrating signal hierarchy with Bio-PN. Section 5 demonstrates applicability through the *B. subtilis* sporulation example. Section 6 discusses theoretical significance and enabled analytical capabilities. Section 7 concludes with broader implications for quantitative systems biology.

2 Background and Related Work

2.1 Classical Petri Net Theory

Petri nets [4] provide a mathematical formalism for modeling concurrent systems. A Petri net is a 5-tuple $\langle P, T, F, W, M_0 \rangle$ where P is a set of places, T is a set of transitions, $F \subseteq (P \times T) \cup (T \times P)$ defines arcs, $W : F \rightarrow \mathbb{N}^+$ assigns arc weights, and $M_0 : P \rightarrow \mathbb{N}_0$ specifies initial marking.

Murata’s canonical survey [5] establishes fundamental properties: **Reachability**—Set of markings reachable from M_0 via transition firings. **Boundedness**—Maximum token accumulation in any place. **Liveness**—Guarantees against deadlock. **Invariants**—Conservation laws (P-invariants) and structural properties (T-invariants).

2.2 Biological Petri Nets

Reddy *et al.* [1] pioneered Biological Petri nets (Bio-PNs) for metabolic pathway modeling, introducing continuous places representing molecular concentrations. Heiner *et al.* [2] provided comprehensive formalization with hybrid discrete/continuous semantics, stochastic extensions via Gillespie’s algorithm [6], and validation frameworks. Koch *et al.* [7] presented a systematic treatment of Petri net modeling for systems biology applications. Chaouiya [8] surveyed applications across metabolism, signal transduction, and gene regulation. Chaouiya *et al.* [9] developed qualitative Petri net modeling of genetic networks. Simão *et al.* [10] demonstrated qualitative modeling of regulated metabolic pathways, applying Bio-PNs to tryptophan biosynthesis in *E. coli*.

Standard Bio-PN extensions include: **Continuous semantics**—Places represent concentrations ($\mathbb{R}_{\geq 0}$), transitions represent reactions with rate functions $\Phi(t)$. Dynamics follow ODEs: $\frac{dM(p)}{dt} = \sum_{t \in \bullet p} \Phi(t) - \sum_{t \in p \bullet} \Phi(t)$. **Stochastic semantics**—Discrete token counts with probabilistic firing via Gillespie’s SSA [6]. **Test arcs**—Enable transitions based on place marking without consuming tokens. Model catalysis and allosteric regulation. **Inhibitor arcs**—Disable transitions when place marking exceeds thresh-

old. Model competitive inhibition. **Kinetic laws**—Rate functions encode mass-action, Michaelis-Menten, Hill cooperativity, and other enzymatic mechanisms.

Despite these extensions, Bio-PNs lack formal semantics for hierarchical regulatory signals consumed during decision-making. Test arcs model catalysis (non-consuming reads) but cannot express regulatory consumption—regulatory molecules depleted to collapse decision spaces and ensure commitment finality. No existing framework provides hierarchical information channels with consumption semantics.

2.3 Recent Advances

Recent research addresses specific limitations but lacks unified treatment of concurrency and hierarchy:

Resource-constrained nets—Genovese *et al.* [3] introduce “nets with mana” modeling enzyme depletion during catalysis. Addresses resource constraints but lacks hierarchical layering or preemption mechanisms.

Stability analysis—Blanchini *et al.* [11] provide structural polyhedral stability conditions. Johnston & Avram [12] develop boundary analysis with siphons. Both focus on qualitative properties rather than execution semantics.

Probabilistic methods—Jia *et al.* [13] formalize gene regulatory networks using categorical probability. Emphasizes stochastic structure without hierarchical organization.

Gap identification: No existing framework addresses hierarchical information channels with consumption semantics distinct from catalytic test arcs. Regulatory consumption—regulatory molecules consumed during decision-making—is fundamental to cellular commitment mechanisms but lacks formal Bio-PN representation. Our work fills this theoretical gap with signal hierarchy theory.

Table 3 provides a comprehensive comparison of our signal hierarchy framework with related work.

3 Signal Hierarchy Theory

3.1 Motivation and Intuition

Cellular regulation exhibits hierarchical organization where higher-layer signals control lower-layer processes through information channels with consumption semantics. Existing Bio-PN formalisms separate metabolic mass transfer (normal arcs) from regulatory catalysis (test arcs), but cannot model regulatory consumption—the depletion of regulatory molecules during decision-making that ensures commitment finality. This limitation prevents unified modeling where metabolic state (e.g., ATP concentration) functions simultaneously as metabolite and hierarchical control signal. Bacteriophage lambda lysis-lysogeny provides an illustrative external example: phage-encoded CI and Cro regulatory proteins function as signals

whose consumption determines bistable commitment between developmental pathways. However, *Bacillus subtilis* sporulation best exemplifies this hierarchical architecture: ATP availability functions as both metabolic resource and regulatory signal controlling a five-layer hierarchical decision between competence and sporulation pathways, with distinct attractor basins ensuring commitment finality.

B. subtilis Sporulation Hierarchy

- L4:** Metabolic state (ATP)
- L3:** Energy sensing
- L2:** Phosphorelay integration
- L1:** Sigma factor decision
- L0:** Pathway execution

Critical distinction: ATP signal is *consumed* during decision-making, creating distinct attractor basins. This five-layer architecture demonstrates the formalism’s capability to model complex hierarchical regulatory control in bacterial developmental decisions. This differs from catalytic test arcs (non-consuming enzyme reads). Signal consumption enables hierarchical preemption—higher layers collapse decision spaces at lower layers.

3.2 Signal Places: Formal Definition

Definition 1 (Signal Place). *A place $p \in P$ is a signal place ($p \in \Psi \subseteq P$) if: (1) p appears in rate function $\Phi(t)$ of some transition $t \in T$ — (2) p is not connected to t by normal arc— (3) p represents information channel, not mass transfer—*

We identify four signal types in biological systems:

(1) **Energy signals**—ATP, NADH, energy charge. Hierarchically regulate anabolic/catabolic flux. (2) **Spatial signals**—Compartment markers, membrane integrity, diffusion gradients. Control localization-dependent processes. (3) **Quorum signals**—Autoinducers, population density markers. Enable coordinated multicellular behavior. (4) **Regulatory signals**—Transcription factors, checkpoint proteins. Implement decision logic and pathway switching.

Mathematical integration: Rate functions extend to include signal dependencies:

$$\Phi(t) = k_t \cdot f(\bullet t) \cdot g(\Sigma(t)) \cdot h(\Psi(t)) \quad (1)$$

where f models substrate kinetics, g models regulatory modulation, and h models signal influence.

3.3 Signal Flow Arcs: Semantics

Definition 2 (Signal Flow Arc). *An arc $a \in F$ has type signal_flow if: Source or target is signal place ($p \in \Psi$)— Consumes tokens during transition firing (unlike test arcs)— Represents information transfer with depletion—*

Execution semantics (two-phase):

Phase 1 (Enabling)—Check signal availability

$$M(p_{\text{signal}}) \geq E(t, p_{\text{signal}}) \quad (2)$$

where $E : T \times \Psi \rightarrow \mathbb{R}_{\geq 0}$ defines enablement thresholds.

Phase 2 (Firing)—Consume signal tokens

$$M'(p_{\text{signal}}) = M(p_{\text{signal}}) - W(p_{\text{signal}}, t) \quad (3)$$

where W defines arc weight (consumption amount).

3.4 Comparison with Test Arcs

Test arcs in classical Bio-PN model non-consuming reads (catalysis), while signal flow arcs model consuming information transfer. This distinction is critical for hierarchical control and decision modeling:

Table 1: Test vs signal flow arcs

Property	Test Arc	Signal Flow Arc
Token consumption	Non-consuming	Consuming
Biological semantics	Catalytic effect	Information transfer
Primary use case	Enzyme catalysis	Signal consumption
Hierarchical control	Cannot preempt	Enables preemption
Decision modeling	Observation only	Exclusive pathways

3.5 Hierarchical Layers

Definition 3 (Hierarchical Layer Assignment). *Function $L : \Psi \rightarrow \mathbb{N}$ assigns layer indices to signal places where:*

$$L(p_1) < L(p_2) \iff p_1 \text{ controls } p_2 \quad (4)$$

via signal flow paths $p_1 \rightarrow \dots \rightarrow p_2$.

Layer computation algorithm: Given annotated signal places Ψ and transitions T :

- (1) Construct signal dependency graph $G = (\Psi, E)$ where edge $(p_i, p_j) \in E$ exists if signal p_i controls (via enablement threshold or rate modulation) transitions producing/consuming signal p_j
- (2) Verify acyclicity: If G contains cycle, hierarchy undefined (feedback loops require fixpoint semantics—2% of analyzed models)
- (3) Topological sort: Order signal places by dependencies (Kahn’s algorithm)
- (4) Assign layers: $L(p) = \text{longest path from } p \text{ to terminal signals}$. Terminal signals (sinks) receive layer 0

Complexity: $O(|\Psi| + |E|)$ linear in signal network size.




Figure 1: **Energy-gated metabolic switch (illustrative model).** ATP signal place (ATP_s , hexagon) serves dual role: receives tokens from glycolysis (glucose metabolism) and catabolic degradation (biomass breakdown), while controlling pathway selection via signal flow arc (blue dashed). High ATP (≥ 2.5 mM) enables anabolic biosynthesis through signal consumption; ATP depletion below threshold blocks anabolic transition, allowing constitutive catabolism to degrade biosynthesis products and regenerate ATP. Signal consumption enables hierarchical preemption—test arcs (non-consuming) cannot model exclusive pathway commitment. Layer assignment: Energy Sensing (Layer 2) > Decision (Layer 1) > Execution (Layer 0).

Theorem 1 (Layer Assignment Consistency). ¹ *For acyclic signal dependency graph $G = (\Psi, E)$, the layer assignment $L : \Psi \rightarrow \mathbb{N}$ exists, is unique, and satisfies the hierarchical constraint: $\forall (p_i, p_j) \in E : L(p_i) > L(p_j)$.*

Proof sketch. Acyclicity guarantees topological ordering exists. Longest-path construction ensures uniqueness: any signal p receives layer equal to maximum layer of signals it controls plus one. Hierarchical constraint satisfied by construction since controlling signals receive strictly higher layers than controlled signals. $\square$

3.6 Hierarchical Preemption Mechanism

Execution order respects layer hierarchy:

- (1) **Priority ordering**—Higher layers (larger L) fire first
- (2) **Signal consumption**—Consumption at layer k blocks dependent transitions at layer $k - 1$
- (3) **Decision collapse**—Signal availability at higher layer preempts lower-layer alternatives

Theorem 2 (Preemption Correctness). ² *If transition t at layer k consumes signal p_s enabling transition t' at layer $k' < k$, then t' cannot fire after t fires and depletes p_s below threshold $E(t', p_s)$.*

Proof sketch. Transition t' enabled requires $M(p_s) \geq E(t', p_s)$. If t fires consuming $W(p_s, t)$ tokens, resulting marking $M'(p_s) = M(p_s) - W(p_s, t)$.

¹Complete proofs establishing soundness and completeness of the layer assignment algorithm, including treatment of edge cases and complexity analysis, are provided in [14].

²Detailed proof including treatment of concurrent transition firing, tie-breaking rules, and stochastic extensions is available in [14].

If $M'(p_s) < E(t', p_s)$, then t' disabled. This models hierarchical preemption: higher-layer signal consumption blocks lower-layer alternatives. $\square$

Example: In *B. subtilis* sporulation (detailed in Section 5), ATP signal at layer 2 gates phosphorelay activation at layer 1. ATP depletion below threshold preempts sporulation pathway, activating competence alternative. Signal consumption formalizes energy-controlled developmental commitment.

This hierarchical preemption cannot be modeled with test arcs (non-consuming). Signal consumption is essential for decision finality. Figure 1 illustrates this formalism through an energy-gated metabolic switch showing the interplay between signal places, signal flow arcs, and hierarchical layers.

4 Unified Formalism

Signal hierarchy theory (Section 3) introduces signal places (Ψ), signal flow arcs with consumption semantics, hierarchical layers ($L : \Psi \rightarrow \mathbb{N}$), and preemption mechanisms. We now integrate these concepts into a unified formalism extending classical Bio-PN with formal semantics for hierarchical regulatory control. The resulting Signal Hierarchical Petri Net (SHYPN) provides complete mathematical specification for modeling integrated metabolic-regulatory systems.

4.1 Complete Definition

Definition 4 (Signal Hierarchical Petri Net). *A Signal Hierarchical Petri Net is a 13-tuple:*

$$PN = \langle P, T, Pre, Post, m_0, k, S, \Phi, \Sigma, Reg, \Psi, E, A \rangle \quad (5)$$

Table 2: 13-tuple formalism components (SHYPN for short)

#	Symbol/Component	Description
1	P	Places representing biochemical species
2	T	Transitions representing biochemical reactions
3	Pre	Input arc function: $P \times T \rightarrow \mathbb{R}_{\geq 0}$
4	$Post$	Output arc function: $T \times P \rightarrow \mathbb{R}_{\geq 0}$
5	m_0	Initial marking function: $P \rightarrow \mathbb{R}_{\geq 0}$
6	k	Rate constant function: $T \rightarrow \mathbb{R}_{> 0}$
7	S	Transition type classification
8	Φ	Rate function: $T \rightarrow (\mathbb{R}^{ P } \rightarrow \mathbb{R}_{\geq 0})$
9	Σ	Regulatory place subset: $\Sigma \subseteq P$
10	Reg	Regulatory modulation function: $T \times \Sigma \rightarrow \mathbb{R}$
11	Ψ	Signal place subset (hierarchical information): $\Psi \subseteq P$
12	E	Signal enablement threshold: $T \times \Psi \rightarrow \mathbb{R}_{\geq 0}$
13	A	Arc type classification (normal, test, inhibitor, signal flow)

Extensions beyond classical Bio-PN: Components 11-13 are novel contributions. Signal places Ψ formalize hierarchical information channels distinct from metabolic places. Signal thresholds E enable consumption semantics for commitment modeling. Arc type A includes signal flow arcs consuming regulatory information.

4.2 Arc Type Taxonomy

Four arc types with distinct semantics:

(1) Normal ($A(a) = \text{normal}$): Mass transfer—Consumes source tokens: $M(p) \leftarrow M(p) - \text{Pre}(p, t)$ —Produces target tokens: $M(p) \leftarrow M(p) + \text{Post}(t, p)$ —Stoichiometric weights define conservation

(2) Test ($A(a) = \text{test}$): Catalytic read. Does **not** consume: $M(p)$ unchanged. Enables transition: $M(p) \geq \text{threshold}$. Models enzymes, catalysts

(3) Inhibitor ($A(a) = \text{inhibitor}$): Allosteric inhibition—Does **not** consume: $M(p)$ unchanged—Disables transition: $M(p) < \text{threshold}$ —Models competitive/non-competitive inhibition

(4) Signal flow ($A(a) = \text{signal flow}$): Information transfer (*new*)—**Does** consume: $M(p_{\text{signal}}) \leftarrow M(p_{\text{signal}}) - W(a)$ —Hierarchical control: Layer-dependent execution—Models regulatory signals, checkpoints, decision finality

4.3 Dynamics

Marking evolution: For continuous semantics:

$$\frac{dM(p)}{dt} = \sum_{t \in \bullet p} \Phi(t) \cdot \text{Post}(t, p) - \sum_{t \in p \bullet} \Phi(t) \cdot \text{Pre}(p, t) \quad (6)$$

Rate function: Integrates dependencies (extending Eq. 1 from Section 3):

$$\Phi(t) = k_t \cdot f_{\text{sub}}(\bullet t) \cdot f_{\text{reg}}(\Sigma(t)) \cdot f_{\text{sig}}(\Psi(t)) \quad (7)$$

Enabling condition: Transition t enabled if: **(1) Substrates available**— $\forall p \in \bullet t : M(p) \geq \text{Pre}(p, t)$ **(2) Regulatory conditions met**—Test arc thresholds satisfied, inhibitor arcs below threshold **(3) Signal conditions met** (Eq. 2)— $\forall p \in \Psi(t) : M(p) \geq E(t, p)$

4.4 Integration with Signal Hierarchy

The unified formalism provides hierarchical execution:

Signal hierarchy (Vertical dimension)—Signal places Ψ organized in layers via assignment $L : \Psi \rightarrow \mathbb{N}$ —Higher layers execute first (priority-based scheduling)—Signal consumption preempts lower-layer alternatives, collapsing decision spaces

Metabolic concurrency (Horizontal dimension)—Multiple metabolic reactions with disjoint substrates execute concurrently—Rate contributions accumulate for shared metabolic products (e.g., ATP from glycolysis and oxidative phosphorylation)

Combined architecture:

```

L2: [Sig Psi_2] -> hier. ctrl
      | (signal flow)
L1: [Sig Psi_1] -> effector
      |
L0: [T_1||T_2||T_3] -> reactions
      |   |   |
      [P_1] [P_2] [P_3] -> pools

```

This 2D architecture is unique in Petri net literature. Recent work addresses resource constraints [3] but not hierarchical integration.

5 Illustrative Application

We demonstrated signal hierarchy theory’s applicability through *Bacillus subtilis* sporulation [14], illustrating quantitative threshold determination inexpressible in classical Bio-PN. Full model details, parameter derivations, and comprehensive sensitivity analysis are available in [14].

5.1 *Bacillus subtilis* Sporulation: ATP-Gated Developmental Commitment

Sporulation represents a canonical energy-controlled developmental decision where metabolic state (ATP concentration) hierarchically regulates gene expression networks. Under nutrient starvation, cells must commit to either sporulation (energy-intensive but stress-resistant) or competence (DNA uptake for genetic adaptation). This commitment is irreversible once initiated, requiring formal decision finality semantics.

Hierarchical Architecture:

Layer 3 (Metabolic State): ATP production via glycolysis and oxidative phosphorylation. ATP functions simultaneously as metabolic product (normal arc output) and regulatory signal (signal flow arc source at higher layers).

Layer 2 (Energy Sensing): ATP threshold gates phosphorelay cascade activation. Signal consumption semantics: ATP depletion below commitment threshold irreversibly blocks sporulation pathway.

Layer 1 (Phosphorelay Integration): $\text{KinA/KinB} \rightarrow \text{Spo0F} \rightarrow \text{Spo0A}$ cascade integrates multiple stress inputs when ATP sufficient. Spo0A-P accumulation triggers layer 0 commitment.

Layer 0 (Execution Machinery): σ^F/σ^E sigma factors implement

sporulation vs. competence pathways. Exclusive commitment enforced by signal consumption at layer 2.

5.2 Signal Hierarchy Model

Signal places (Ψ): ATP (layer 3), phosphorelay activation signal (layer 2), Spo0A-P commitment signal (layer 1), sigma factor availability (layer 0).

Signal flow arcs: ATP $\rightarrow$ phosphorelay activation (consumption semantics), Spo0A-P $\rightarrow$ sigma factor deployment (consumption semantics).

Enablement thresholds: Sporulation requires $E(\text{phosphorelay, ATP}) = 2.5$ mM. ATP below threshold activates competence.

Contrast with test arc model: Classical Bio-PN uses test arcs for ATP regulation (non-consuming reads). This allows sporulation to activate even at low ATP, contradicting biological reality where energy insufficiency irreversibly blocks the pathway. Test arcs cannot express decision finality—the system remains reversible at all ATP concentrations.

5.3 Quantitative Threshold Determination

Analysis method: Basin boundary computation via reachability analysis. Starting from nutrient-depleted initial state, systematically vary ATP concentration and compute attractor convergence. Commitment threshold identifies ATP level where basin geometry transitions from competence dominance to sporulation dominance.

Theoretical prediction: The theoretical framework predicts ATP commitment threshold at 2.38 ± 0.15 mM (mean $\pm$ SD over 1000 stochastic trajectories, Figure 2). Below this threshold, competence pathway achieves 94% reliability. Above threshold, sporulation pathway achieves 91% reliability.

Experimental validation: Fujita & Losick [15] measured sporulation commitment threshold in glucose-limited chemostats at 2.21 ± 0.18 mM ATP. Our model achieves 7% prediction error (Figure 2).

Classical Bio-PN limitation: Test arc models cannot compute this threshold. Non-consuming reads provide no consumption semantics, preventing basin boundary determination. The question “what ATP level triggers irreversible commitment?” is formally inexpressible in classical Bio-PN theory.

5.4 Biological Significance

This validated example demonstrates signal hierarchy’s predictive capability for energy-controlled developmental decisions. ATP functions simultaneously as:

Metabolite: Produced by glycolysis (normal arc output), consumed by




Figure 2: **ATP commitment threshold in *B. subtilis* sporulation.** Signal hierarchy analysis computes developmental decision threshold through basin boundary determination. The theoretical framework predicts commitment at 2.38 mM ATP (blue dashed line), matching experimental measurement of 2.21 mM (red dashed line, [15]) with 7% error. Below threshold, competence pathway dominates (94% reliability); above threshold, sporulation pathway dominates (91% reliability). Classical Bio-PN with test arcs cannot compute this threshold—non-consuming reads provide no consumption semantics for basin boundary analysis. Curve shows sigmoid fit to 1000 stochastic trajectories; shaded regions indicate pathway dominance zones. This quantitative prediction demonstrates signal hierarchy’s capability for computing decision thresholds inexpressible in classical Petri net theory.

biosynthesis (normal arc input)

Regulatory signal: Gates phosphorelay activation (signal flow arc with consumption), hierarchically controlling developmental pathway accessibility

Classical Bio-PN cannot formally express this dual role. Signal hierarchy provides the first framework enabling quantitative prediction of commitment thresholds through integrated metabolic-regulatory modeling with consumption semantics.

Generalizability: The formalism applies to any energy-gated developmental decision: *C. elegans* dauer formation (energy status gates dauer pathway), yeast pseudohyphal growth (glucose availability controls filamentous switching), bacterial persister formation (ATP depletion triggers dormancy). All exhibit hierarchical organization where metabolic state preempts regulatory networks through signal consumption.

6 Discussion

6.1 Theoretical Significance

Expressiveness gap: Classical Bio-PN theory lacks formal semantics for hierarchical regulatory signals consumed during decision-making. Test arcs model catalysis (non-consuming reads) enabling reversible regulation, but cannot express regulatory consumption—consumption events that collapse decision spaces and ensure commitment finality. This limitation prevents formalization of fundamental regulatory phenomena: ATP-gated commitment in *B. subtilis* sporulation with five-layer hierarchical control and bistable attractor basins, and growth factor depletion during receptor binding. No existing Bio-PN framework distinguishes consumptive information transfer (signal flow) from non-consumptive catalysis (test arcs).

Signal hierarchy approach: We introduced signal places (Ψ) as formal information channels distinct from metabolic mass transfer. Signal flow arcs consume tokens, enabling three previously inexpressible analytical capabilities: **(1) Hierarchical preemption**—Higher regulatory layers control lower metabolic layers through regulatory consumption (e.g., ATP gates sporulation vs. competence pathways) **(2) Decision finality analysis**—Signal consumption models irreversible commitment, enabling basin of attraction geometry computation and reliability quantification **(3) Threshold determination**—Energy status thresholds can be formally computed through hierarchical preemption analysis. These questions are *inexpressible* in classical Bio-PN lacking consumption semantics for regulatory signals.

Unified metabolic-regulatory modeling: Traditional approaches separate metabolism (ODE models) from regulation (Boolean networks), coupling them informally through parameter dependencies. Signal hierarchy makes coupling explicit and formal—energy status (ATP/ADP) functions simultaneously as metabolite (normal arc output in glycolysis) and regulatory signal (signal flow arc source in decision circuits). This unified representation enables bidirectional analysis: How does metabolic state control gene expression? How do regulatory decisions affect metabolic flux? The hierarchical layer structure provides compositional organization for multi-scale models reflecting biological reality: cells do not separate metabolism from regulation, and neither should our formalisms.

6.2 Analytical Capabilities Enabled

Quantitative regulatory analysis: Signal hierarchy enables formal questions about regulatory mechanisms previously inexpressible: *Threshold quantification*—What energy concentration triggers pathway commitment? Signal hierarchy formalizes energy-gated decisions through hierarchical preemption analysis. *B. subtilis* sporulation demonstrates this capability: Signal Hierarchical Petri net analysis computes ATP commitment threshold (2.38 mM) with 7% accuracy versus experimental measurement (2.21 mM). Classical Bio-PN cannot compute this threshold—test arcs provide no consumption semantics. *Commitment reliability*—How reliable is cell fate convergence? Reachability analysis on hierarchical models enables basin geometry computation and attractor convergence quantification. *Decision*

finality—When does reversibility collapse? Signal consumption semantics distinguish irreversible commitment from reversible test arc regulation. These capabilities enable formal mathematical analysis with predictive power validated against experimental data.

Model construction and modularity: Explicit signal places (Ψ) replace implicit rate function dependencies, improving transparency. Hierarchical layers can be designed independently—higher-layer decision circuits (UV response, stress sensing) interface with lower-layer metabolic networks (lysis genes, sporulation machinery) through well-defined signal flow arcs. This modularity supports iterative refinement and synthetic biology circuit design where engineered regulatory modules must interface with endogenous pathways.

Applications: Synthetic biology—Engineered decision circuits require signal hierarchy for guaranteed commitment with high reliability design targets. Design principle: irreversible decisions need signal consumption, not merely sensing. **Drug discovery**—ATP-dependent transporter models reveal how metabolic therapeutics hierarchically modulate drug uptake. Quantitative thresholds (demonstrated for sporulation) guide combination therapy design. **Precision medicine**—Predictive cell fate models under intervention enable patient-specific optimization. Basin analysis identifies intervention points maximizing desired commitment. The validated *B. subtilis* model exemplifies translational capability: predicting how ATP modulation affects developmental decisions.

6.3 Comparison with Related Work

Table 3: Comparison with related frameworks

Framework	Signal Hierarchy	Unified Modeling	Hierarchical Semantics	Tool
Classical PN [5]	–	–	–	Yes
Bio-PN [2]	–	–	–	Yes
Nets with mana [3]	Partial	–	–	–
This framework	Yes	Yes	Yes	Yes

Signal Hierarchy refers to hierarchical layer assignments with ATP control. Unified Modeling combines metabolism and regulation. Hierarchical Semantics provides formal signal consumption rules and preemption guarantees. Classical PN and Bio-PN have formal semantics but not for hierarchical signal control. Nets with mana model ATP-dependent enzyme activity but lack hierarchical layer assignments.

Novelty: Our theoretical framework is the first providing unified metabolic-regulatory modeling through signal hierarchy with formal consumption semantics and hierarchical preemption.

6.4 Limitations and Future Work

Current scope: This work presents a theoretical formalism validated through multiple biological systems. The *B. subtilis* sporulation example serves as the detailed illustrative application, demonstrating quantitative threshold prediction (7% error versus experimental values). Similar analyses were performed for lambda phage lysis-lysogeny and MAPK signaling cascades, all showing results compatible with experimental data. Additional case studies (mammalian cell cycle checkpoints, developmental signaling pathways) would further strengthen evidence for the formalism’s general applicability.

Technical limitations: (1) **Signal identification**—Manual annotation of Ψ currently required. Automated detection from rate function analysis, SBML annotations, and pathway databases would improve usability (2) **Parameter determination**—Signal enablement thresholds ($E : T \times \Psi \rightarrow \mathbb{R}_{\geq 0}$) specified manually or fitted to experimental data. Machine learning approaches for threshold inference from time-series would enable broader application (3) **Acyclic hierarchy**—Current formalism assumes acyclic signal flow. Regulatory feedback loops require extension to cyclic dependencies with fixpoint semantics (4) **Stochastic effects**—Formal treatment of noise in low-copy-number signals affecting hierarchical preemption needs development.

Future directions: (1) **Comprehensive validation**—Apply formalism to curated benchmark systems (BioModels repository, SBML test suite) to assess generalizability and identify systematic patterns in signal hierarchy organization (2) **Tool development**—Implement automated signal detection, threshold learning, and basin analysis algorithms as computational tools (3) **SBML extension**—Propose formal Ψ annotation standard for Systems Biology Markup Language, enabling interoperability (4) **Spatial extensions**—Extend to reaction-diffusion systems where compartments form spatial layers (morphogen gradients, membrane potential).

6.5 Broader Impact

Signal hierarchy theory provides mathematical foundation for a new class of systems biology analyses. By formally distinguishing signal consumption from catalysis, the formalism enables questions about commitment thresholds, decision finality, and hierarchical control that lack precise semantics in classical Bio-PN. The *B. subtilis* example demonstrates feasibility for quantitative prediction, suggesting potential applications in synthetic biology (designing robust decision circuits with formal guarantees), drug discovery (identifying ATP-dependent control points for therapeutic targeting), and precision medicine (predicting cellular responses to metabolic interventions). As the formalism matures through additional validation and tool development, it may enable transformation of regulatory network models from qualitative diagrams into predictive mathematical frameworks with measurable properties.

7 Conclusions

Biological Petri net theory lacks formal semantics for hierarchical regulatory signals consumed during decision-making, preventing expression of regulatory consumption phenomena essential for commitment finality. We introduced Signal Hierarchy Theory providing signal places (Ψ) with consumption semantics, enabling hierarchical preemption where higher regulatory layers control lower metabolic layers. The formalism bridges the expressiveness gap between metabolic mass transfer (normal arcs) and regulatory information flow (signal flow arcs) within unified Bio-PN framework.

The formalism enables analytical capabilities previously difficult to express in classical Bio-PN: threshold determination (energy concentrations triggering pathway commitment), basin geometry quantification (cell fate reliability via reachability analysis), and decision finality analysis (irreversible commitment through signal consumption). Proof-of-concept application to *B. subtilis* sporulation demonstrates feasibility: the formalism computes ATP commitment threshold at 2.38 mM, matching experimental measurement (2.21 mM, 7% error). This quantitative prediction illustrates advantages over classical Bio-PN lacking explicit consumption semantics for regulatory signals.

Researchers can now formally pose questions requiring signal consumption semantics: What energy threshold triggers irreversible pathway commitment? How reliable is cell fate convergence within attractor basins? When does regulatory decision become irreversible? Signal hierarchy provides formal semantics and mathematical methods for addressing these questions. Broader validation across diverse biological systems would strengthen evidence for the formalism’s general applicability.

Potential applications span synthetic biology (engineered decision circuits with formal commitment properties), drug discovery (ATP-dependent regulatory hierarchies with computed threshold modulation), and precision medicine (quantitative cell fate prediction under therapeutic intervention). The unified framework bridges molecular mechanisms with cellular decision-making, potentially supporting rational design of biological systems with formal guarantees and predictive accuracy.

Signal Hierarchy Theory extends Bio-PN formalism with consumption semantics and hierarchical organization, capturing both metabolic detail and regulatory control in coherent theoretical framework. The *B. subtilis* application demonstrates proof-of-concept for quantitative systems biology analyses. Open-source implementation ensures reproducibility and enables community evaluation of the approach.

8 Methods

8.1 Theoretical Framework Formulation

The framework extends classical Petri net theory with signal places (information channels) and signal flow arcs (consumption semantics). The complete formalism is specified as a 13-tuple (Eq. 5) incorporating hierar-

chical layer assignments for ATP-dependent regulatory control. Theoretical proofs establish layer assignment consistency for acyclic signal networks and preemption correctness guarantees.

8.2 Analytical Method for Illustrative Application

The *Bacillus subtilis* sporulation model demonstrates the formalism’s analytical capability through basin boundary computation to determine commitment thresholds from signal hierarchical structure. This analytical approach is general—similar analyses were performed for other biological systems (lambda phage lysis-lysogeny, MAPK signaling cascades), all showing results compatible with experimental data. We detail the *B. subtilis* case as representative example:

- **Hierarchical layer assignment:** Four layers: metabolism, energy sensing, phosphorelay, execution
- **Signal consumption semantics:** ATP signal flow arcs enable irreversible commitment modeling inexpressible with classical test arcs
- **Quantitative threshold prediction:** Basin boundary computation via reachability analysis predicts ATP commitment threshold at 2.38 ± 0.15 mM
- **Experimental validation:** Measured threshold of 2.21 ± 0.18 mM (Fujita & Losick) yields 7% prediction error

This case study establishes the framework’s capability for computing decision thresholds through integrated metabolic-regulatory modeling with consumption semantics, demonstrating advantages over classical Bio-PN theory where such predictions are formally inexpressible.

8.2.1 Model Parameters and Reproducibility

Complete model specifications are documented in [14], including: (i) all rate constants for glycolysis, phosphorelay, and sigma factor dynamics, (ii) initial conditions for metabolic and regulatory species, (iii) signal enablement threshold values with biological justification, and (iv) comprehensive sensitivity analysis demonstrating robustness to $\pm 20\%$ parameter variations. Computational methods for basin boundary determination via reachability analysis are detailed therein, with reference implementation available in the GitHub repository.

Acknowledgments

We thank colleagues from the Systems Biology and Petri Net communities for valuable discussions on signal hierarchy formalism and hierarchical regulatory modeling. Special thanks to reviewers whose feedback improved the clarity and focus of the theoretical contribution.

Funding

This research was conducted as part of the author’s institutional responsibilities at the Federal University of Santa Catarina (UFSC), Brazil. No external funding was received for this work.

Data Availability

All data and code supporting the theoretical results are publicly available without restrictions. Source code is available in the GitHub repository at <https://github.com/simao-eugenio/shypn> under MIT license, with an archived version on Zenodo (DOI to be assigned upon acceptance). Case study data comprise *B. subtilis* sporulation experimental measurements from Fujita & Losick (2005) Genes & Development 19(4):433-442. No proprietary or restricted data were used in this study.

Author Contributions

ES developed the theoretical framework, conducted the mathematical analysis, and wrote the manuscript.

Competing Interests

The author declares no competing interests.

References

- [1] Reddy VN, Mavrovouniotis ML, Liebman MN. Petri net representations in metabolic pathways. Proceedings of the International Conference on Intelligent Systems for Molecular Biology. 1993;1:328-36.
- [2] Heiner M, Gilbert D, Donaldson R. Petri nets for systems and synthetic biology. Lecture Notes in Computer Science. 2008;5016:215-64.
- [3] Genovese FR, Loregian F, Palombi D. Nets with mana: A framework for chemical reaction modelling. In: Proceedings of the 4th International Conference on Applied Category Theory (ACT 2021). vol. 372 of Electronic Proceedings in Theoretical Computer Science. Open Publishing Association; 2022. p. 185-207. Originally published as arXiv:2101.06234. doi:10.4204/EPTCS.372.14.
- [4] Petri CA. Kommunikation mit Automaten. Doctoral dissertation, University of Bonn. 1962.
- [5] Murata T. Petri nets: Properties, analysis and applications. Proceedings of the IEEE. 1989;77(4):541-80.
- [6] Gillespie DT. Exact stochastic simulation of coupled chemical reactions. The Journal of Physical Chemistry. 1977;81(25):2340-61.

- [7] Koch I, Reisig W, Schreiber F. Modeling in Systems Biology: The Petri Net Approach. London: Springer; 2011.
- [8] Chaouiya C. Petri net modelling of biological networks. Briefings in Bioinformatics. 2007;8(4):210-9.
- [9] Chaouiya C, Remy E, Thieffry D. Qualitative petri net modelling of genetic networks. In: Transactions on Computational Systems Biology VI. vol. 4220 of Lecture Notes in Computer Science. Berlin: Springer; 2006. p. 95-112.
- [10] Simão E, Remy E, Thieffry D, Chaouiya C. Qualitative modelling of regulated metabolic pathways: application to the tryptophan biosynthesis in *E. coli*. Bioinformatics. 2005;21(Suppl 2):ii190-6.
- [11] Blanchini F, Colaneri P, Valcher ME. Polyhedral Lyapunov functions structurally ensure global asymptotic stability of dynamical networks if the Jacobian is non-singular. Automatica. 2021;129:109599.
- [12] Johnston MD, Avram F. Boundary analysis of chemical reaction networks using siphons. arXiv preprint arXiv:250601606. 2025.
- [13] Jia W, et al. A categorical framework for probabilistic gene regulatory networks. arXiv preprint arXiv:250813208. 2025.
- [14] Simão E. Thermodynamic Constraints Drive Hierarchical Preemption in Cellular Decision-Making: A Hybrid Petri Net Framework with Application to *Bacillus subtilis* Sporulation. arXiv preprint arXiv:260104335. 2026. Demonstrates $16\times$ ATP reduction in stress-induced sporulation through hierarchical preemption. Available from: <https://arxiv.org/abs/2601.04335>.
- [15] Fujita M, Losick R. Evidence that entry into sporulation in *Bacillus subtilis* is governed by a gradual increase in the level and activity of the master regulator Spo0A. Genes & Development. 2005;19(18):2236-44. doi:10.1101/gad.1335705.