

Binomial coefficients

Mathematics for Computer Science

Jean-Marc Vincent¹

¹Laboratoire LIG
Equipe-Projet MESCAL
Jean-Marc.Vincent@imag.fr

These notes are only the sketch of the lecture : the aim is to apply the basic counting techniques to the binomial coefficients and establish combinatorial equalities.

References : Concrete Mathematics : A Foundation for Computer Science
Ronald L. Graham, Donald E. Knuth and Oren Patashnik Addison-Wesley
1989 (chapter 5)



Definition

$\binom{n}{k}$ is the number of ways to choose k elements among n elements



<http://www-history.mcs.st-and.ac.uk/Biographies/Pascal.html>

For all integers $0 \leq k \leq n$

$$\binom{n}{k} = \frac{n(n-1) \cdots (n-k+1)}{k!} \quad (1)$$

Hint : Prove it by a combinatorial argument Hint : the number of sequences of k different elements among n is $n(n-1) \cdots (n-k+1)$ and the number of orderings of a set of size k is $k!$.

Basic properties

$$\binom{n}{k} = \frac{n!}{k!(n-k)!} \quad (2)$$

Prove it directly from Equation 1

For all integers $0 \leq k \leq n$

$$\binom{n}{k} = \binom{n}{n-k} \quad (3)$$

Prove it directly from 2

Prove it by a combinatorial argument *Hint : bijection between the set of subsets of size k and ???.*

Exercise

Give a combinatorial argument to prove that for all integers $0 \leq k \leq n$:

$$k \binom{n}{k} = n \binom{n-1}{k-1} \quad (4)$$



Pascal's triangle


Recurrence equation

The binomial coefficients satisfy

$$\binom{n}{k} = \binom{n-1}{k-1} + \binom{n-1}{k} \quad (5)$$

Prove it directly from Equation 1

Prove it by a combinatorial argument *Hint : partition in two parts the set of subsets of size k ; those containing a given element and those not.*



Thanks to Elisha

The binomial theorem

For all integer n and a formal parameter X

$$(1 + X)^n = \sum_{k=0}^n \binom{n}{k} X^k \text{ (Newton 1666)} \quad (6)$$

Prove it by a combinatorial argument *Hint : write*

$(1 + X)^n = \underbrace{(1 + X)(1 + X) \cdots (1 + X)}_{n \text{ terms}}$ in each term chose 1 or X , what is the coefficient of X^k in the result (think "vector of n bits").

Exercises

Use a combinatorial argument to prove :

$$\sum_{k=0}^n \binom{n}{k} = 2^n$$

Use the binomial theorem to prove (give also a combinatorial argument)

$$\sum_{k=0}^n \binom{n}{k} = \sum_{k=0}^n \binom{n}{k} = 2^{n-1}$$

Summations and Decompositions

The Vandermonde Convolution

For all integers m, n, k

$$\sum_{j=0}^k \binom{m}{j} \binom{n}{k-j} = \binom{m+n}{k} \quad (7)$$

Prove it by a combinatorial argument *Hint : choose k elements in two sets one of size m and the other n .*

Exercise

Prove that

$$\sum_{k=0}^n \binom{n}{k}^2 = \binom{2n}{n} \quad (8)$$

Hint : Specify Equation 7

Summations and Decompositions (2)

Upper summation

For all integers $p \leq n$

$$\sum_{k=p}^n \binom{k}{p} = \binom{n+1}{p+1} \quad (9)$$

Exercises

Establish the so classical result

$$\sum_{k=1}^n \binom{k}{1}$$

Compute

$$\sum_{k=2}^n \binom{k}{2}$$

and deduce the value of $\sum_{k=1}^n k^2$

The main rules in combinatorics (I)

Bijection rule

Let A and B be two finite sets if there exists a bijection between A and B then

$$|A| = |B|.$$

Summation rule

Let A and B be two **disjoint** finite sets then

$$|A \cup B| = |A| + |B|.$$

Moreover if $\{A_1, \dots, A_n\}$ is a partition of A (for all $i \neq j$, $A_i \cap A_j = \emptyset$ and $\bigcup_{i=1}^n A_i = A$)

$$|A| = \sum_{i=1}^n |A_i|.$$

The main rules in combinatorics (II)

Product rule

Let A and B be two finite sets then

$$|A \times B| = |A| \cdot |B|.$$

Inclusion/Exclusion principle

Let A_1, A_2, \dots, A_n be sets

$$|A_1 \cup \dots \cup A_n| = \sum_{k=1}^n (-1)^{k+1} \sum_{S \subseteq \{1, \dots, n\}, |S|=k} \left| \bigcap_{i \in S} A_i \right|.$$

Exercises

Illustrate these rules by the previous examples, giving the sets on which the rule apply.