

Mathematics for computer science

Homework 1

SID-LAKHDAR Riyane

15/10/2015

I understand what plagiarism entails and I declare that this report is my own, original work SID-LAKHDAR Riyane october the 15 th, 2015



Abstract

This report summarise, explains and refers to the answers we have designed for the first Mathematics for computer science homework. In this report, we will use the following notations:

- N and K are two finite sets of size n and k .
- N will be represented as

$$N = \{n_i | i \text{ belongs to } [0, n-1]\} \quad (1)$$

- The set of expected f function will include the partial functions: a function f belonging to this set may be undefined on a specific point of its input set N .
If a function f is undefined on a point x belonging to N , we will always write

$$f(x) = \epsilon \quad (2)$$

1 Question 1

A function

$$f : N \rightarrow K \quad (3)$$

is an application which associates to each element n , belonging to N , at most one element k belonging to K .

Thus, building such a function f is equivalent to build a word

$$w = k_0 \dots k_{n-1} \text{ where } \forall i \in [0, n-1] k_i \in K \cup \{\epsilon\} \quad (4)$$

As we have no restriction on f , we have $k+1$ different choices to choose any one of the k_i . This choice is independant from the choice of any k_j where j is different from i . Thus, the number of different word

$$w = k_0 \dots k_{n-1} \quad (5)$$

is

$$\prod_{i=0}^{n-1} (k+1) = (k+1)^n \quad (6)$$

Thus, the number of different functions

$$f : K \rightarrow N \quad (7)$$

with no restriction on f is

$$(k+1)^n \quad (8)$$

This result would be

$$k^n \quad (9)$$

if we only consider the non partial functions.

2 Question 2

Using the same arguments as previously, we can say that our problem is equivalent to find all the different words

$$w = k_0 \dots k_{n-1} \text{ where } \forall i \in [0, n-1] k_i \in K \cup \{\epsilon\} \quad (10)$$

Let w_0 such a word, and f_{w_0} the corresponding injective function. We have:

$$\forall i, j \in [0, n-1] \text{ with } i \neq j, f_{w_0}(n_i) = f_{w_0}(n_j) \implies n_i = n_j \quad (11)$$

which is absurde by definition of n_i and n_j . So

$$\forall i, j \in [0, n-1], i \neq j \implies f_{w_0}(n_i) \neq f_{w_0}(n_j) \quad (12)$$

Using this condition we can conclude that

- To chose the character k_0 of w_0 among K union epsilon, we have $(k+1)$ choices.
- To chose the character k_1 of w_0 among K union epsilon excluding K_0 , we have $(k+1 - 1)$ choices for each k_0 .

- To chose the character k_2 of w_0 among K union epsilon excluding K_0, K_1 , we have $(k+1 - 2)$ choices for each k_0, k_1 .
- To chose the character k_i of w_0 among K union epsilon excluding K_0, K_1, \dots, k_{i-1} , we have $(k+1 - i)$ choices for each k_0, k_1, \dots, k_{i-1} .

Thus, the number of different words $w = k_0 \dots k_{n-1}$ which respect the condition (12), and the number of injective function is:

$$\prod_{i=0}^{n-1} k+1-i = \frac{(k+1)!}{(k-n+1)!} \text{ (as } k \geq n \text{ by definition of an injective function)} \quad (13)$$

This result would be

$$\frac{k!}{(k-n)!} \quad (14)$$

if we only consider the non partial functions.

3 Question 3

Let's f a surjection between N and K . So:

- Each element form K is mapped to at least 1 element from N
- Each element from N is mapped to at most 1 element from K

Thus, according to pigeonhole principle, n and k must to respect the rule:

$$n \geq k \quad (15)$$

Let f such a surjective function, and w its corresponding world:

$w = w_0, \dots, w_{n-1}$ where $w_i = f(n_i)$.

According to the definition of a surjective function, the word w should contain, at least, once every k_i bellonging to K . Thus, a word w must be a combination of

$$k_0, \dots, k_{k-1} \quad c_0, \dots, c_{n-k-1} \text{ where } c_i \in K \cup \{\epsilon\} \quad (16)$$

But we know that the number of different

$$k_0, \dots, k_{k-1} \text{ is } k! \quad (17)$$

and the number of different

$$c_0, \dots, c_{n-k-1} \text{ where } c_i \in K \cup \{\epsilon\} \text{ is } (k+1)^{n-k} \quad (18)$$

Thus, the number of different word w , and the number of surjective functions is

$$(k+1)^{(n-k)} k! \text{ with } n \geq k \quad (19)$$

4 Question 4

By definition of an bijection, the input and output sets must have the same cardinals. Thus, in the following answer, we will consider the condition $k = n$ respected.

Building a bijection between N and K is equivalent to build a word

$$\omega = k_0, k_1, \dots, k_{n-1} \text{ where } \forall i, j \in [0, n-1] k_i \neq k_j \quad (20)$$

Thus,

- To chose the character k_0 among K , we have k choices.
- To chose the character k_1 among K excluding K_0 , we have $(k - 1)$ choices for each k_0 .
- To chose the character k_2 among K excluding K_0, K_1 , we have $(k - 2)$ choices for each k_0, k_1 .
- To chose the character k_i among K excluding K_0, K_1, \dots, k_{i-1} , we have $(k - i)$ choices for each k_0, k_1, \dots, k_{i-1} .

Thus, the number of different bijective function from N to K is

$$\prod_{i=0}^{n-1} k - i = \prod_{i=0}^{n-1} n - i = n! \quad (21)$$