Week 1: Complex numbers and differential equations

Simon Elias Schrader

August 23th 2024

 Q1.1 Why is there an upper limit to the photon energy that can be observed in the emission spectrum of the hydrogen atom?


- Q1.1 Why is there an upper limit to the photon energy that can be observed in the emission spectrum of the hydrogen atom?
- Q1.5 Which of the experimental results for the photoelectric effect suggests that light can display particle-like behavior?

- Q1.1 Why is there an upper limit to the photon energy that can be observed in the emission spectrum of the hydrogen atom?
- Q1.5 Which of the experimental results for the photoelectric effect suggests that light can display particle-like behavior?
- Q1.9 In the double-slit experiment, researchers found that an equal amount of energy passes through each slit. Does this result allow you to distinguish between purely particle-like and purely wave-like behavior?

- Q1.1 Why is there an upper limit to the photon energy that can be observed in the emission spectrum of the hydrogen atom?
- Q1.5 Which of the experimental results for the photoelectric effect suggests that light can display particle-like behavior?
- Q1.9 In the double-slit experiment, researchers found that an equal amount of energy passes through each slit. Does this result allow you to distinguish between purely particle-like and purely wave-like behavior?
- Q1.10 The inability of classical theory to explain the spectral density distribution of a blackbody was called the ultraviolet catastrophe. Why is this name appropriate?

- Q1.1 Why is there an upper limit to the photon energy that can be observed in the emission spectrum of the hydrogen atom?
- Q1.5 Which of the experimental results for the photoelectric effect suggests that light can display particle-like behavior?
- Q1.9 In the double-slit experiment, researchers found that an equal amount of energy passes through each slit. Does this result allow you to distinguish between purely particle-like and purely wave-like behavior?
- Q1.10 The inability of classical theory to explain the spectral density distribution of a blackbody was called the ultraviolet catastrophe. Why is this name appropriate?
- Q1.12 Why is a diffraction pattern generated by an electron gun formed by electrons interfering with themselves rather than with one another?

- Every complex number $z \in \mathbb{C}$ can be written as z = a + ib, for $a, b \in \mathbb{R}$.
- We can also write z as $z=re^{i\theta}$, where $r,\theta\in\mathbb{R}$ and $r\geq 0$, $\theta\in[0,2\pi)$.



• We have the relation (Euler's equation)

$$e^{i\theta} = \cos(\theta) + i\sin(\theta)$$

We have the relation (Euler's equation)

$$e^{i\theta} = \cos(\theta) + i\sin(\theta)$$

• The complex conjugate of a complex number $z = a + ib = re^{i\theta}$ is

$$z^* = a - ib = re^{-i\theta}$$

We have the relation (Euler's equation)

$$e^{i\theta} = \cos(\theta) + i\sin(\theta)$$

• The complex conjugate of a complex number $z = a + ib = re^{i\theta}$ is

$$z^* = a - ib = re^{-i\theta}$$

 The length/modulus/magnitude/absolute value of a complex number is

$$|z|=r=\sqrt{a^2+b^2}=\sqrt{zz^*}$$

We have the relation (Euler's equation)

$$e^{i\theta} = \cos(\theta) + i\sin(\theta)$$

• The complex conjugate of a complex number $z = a + ib = re^{i\theta}$ is

$$z^* = a - ib = re^{-i\theta}$$

 The length/modulus/magnitude/absolute value of a complex number is

$$|z|=r=\sqrt{a^2+b^2}=\sqrt{zz^*}$$

• A note on notation: Sometimes, z^* is written as \bar{z} . We have $a = \text{Re}(z) = \Re(z)$, $b = \text{Im}(z) = \Im(z)$.

Example exercises

- Calculate $z_3 = z_1/z_2$, where $z_1 = r_1e^{i\theta_1}$, $z_2 = r_2e^{i\theta_2}$.
- Write z = 3 + 4i in the form $re^{i\theta}$.
- Write the following number in the form a + ib:

$$z=\frac{3+5i}{2-4i}$$

• For two complex numbers z_1, z_2 , show that

$$(z_1z_2)^*=z_1^*z_2^*$$

 We consider the first order differential equation (related to the Schrödinger equation)

$$\frac{d}{dt}f(t) = -iE/\hbar f(t)$$

Can you solve it (see eq. 2.23 in the book)?

 We consider the first order differential equation (related to the Schrödinger equation)

$$\frac{d}{dt}f(t) = -iE/\hbar f(t)$$

Can you solve it (see eq. 2.23 in the book)?

• What happens in the previous exercise if f(t) is a function of two variables, f(x,t) (and d/dt becomes $\partial/\partial t$)?

 We consider the first order differential equation (related to the Schrödinger equation)

$$\frac{d}{dt}f(t) = -iE/\hbar f(t)$$

Can you solve it (see eq. 2.23 in the book)?

- What happens in the previous exercise if f(t) is a function of two variables, f(x,t) (and d/dt becomes $\partial/\partial t$)?
- How do you solve the differential equation

$$\frac{d^2f(x)}{dx^2} + B\frac{df(x)}{dx} + Cf(x) = 0$$

 We consider the first order differential equation (related to the Schrödinger equation)

$$\frac{d}{dt}f(t) = -iE/\hbar f(t)$$

Can you solve it (see eq. 2.23 in the book)?

- What happens in the previous exercise if f(t) is a function of two variables, f(x,t) (and d/dt becomes $\partial/\partial t$)?
- How do you solve the differential equation

$$\frac{d^2f(x)}{dx^2} + B\frac{df(x)}{dx} + Cf(x) = 0$$

• Consider now the partial differential equation

$$\frac{\partial^2 f(x,y)}{\partial x^2} + \frac{\partial^2 f(x,y)}{\partial y^2} + f(x,y) = 0$$

How do you solve it?