

# Slow Component of $\dot{V}O_2$ Kinetics: Mechanistic Bases and Practical Applications

ANDREW M. JONES<sup>1</sup>, BRUNO GRASSI<sup>2</sup>, PETER M. CHRISTENSEN<sup>3</sup>, PETER KRUSTRUP<sup>1,3</sup>, JENS BANGSBO<sup>3</sup>, and DAVID C. POOLE<sup>4</sup>

<sup>1</sup>*Sport and Health Sciences, College of Life and Environmental Sciences, St. Luke's Campus, University of Exeter, Exeter, UNITED KINGDOM;* <sup>2</sup>*Department of Medical and Biological Sciences, School of Medicine, University of Udine, Udine, ITALY;* <sup>3</sup>*Department of Exercise and Sport Sciences, The August Krogh Building, University of Copenhagen, Copenhagen, DENMARK;* and <sup>4</sup>*Departments of Kinesiology, Anatomy and Physiology, Kansas State University, Manhattan, KS*

## ABSTRACT

JONES, A. M., B. GRASSI, P. M. CHRISTENSEN, P. KRUSTRUP, J. BANGSBO, and D. C. POOLE. Slow Component of  $\dot{V}O_2$  Kinetics: Mechanistic Bases and Practical Applications. *Med. Sci. Sports Exerc.*, Vol. 43, No. 11, pp. 2046–2062, 2011. The  $\dot{V}O_2$  slow component, a slowly developing increase in  $\dot{V}O_2$  during constant-work-rate exercise performed above the lactate threshold, represents a progressive loss of skeletal muscle contractile efficiency and is associated with the fatigue process. This brief review outlines the current state of knowledge concerning the mechanistic bases of the  $\dot{V}O_2$  slow component and describes practical interventions that can attenuate the slow component and thus enhance exercise tolerance. There is strong evidence that, during constant-work-rate exercise, the development of the  $\dot{V}O_2$  slow component is associated with the progressive recruitment of additional (type II) muscle fibers that are presumed to have lower efficiency. Recent studies, however, indicate that muscle efficiency is also lowered (resulting in a “mirror-image”  $\dot{V}O_2$  slow component) during fatiguing, high-intensity exercise in which additional fiber recruitment is unlikely or impossible. Therefore, it seems that muscle fatigue underpins the  $\dot{V}O_2$  slow component, although the greater fatigue sensitivity of recruited type II fibers might still play a crucial role in the loss of muscle efficiency in both situations. Several interventions can reduce the magnitude of the  $\dot{V}O_2$  slow component, and these are typically associated with an enhanced exercise tolerance. These include endurance training, inspiratory muscle training, priming exercise, dietary nitrate supplementation, and the inspiration of hyperoxic gas. All of these interventions reduce muscle fatigue development either by improving muscle oxidative capacity and thus metabolic stability or by enhancing bulk muscle  $O_2$  delivery or local  $\dot{Q}O_2$ -to- $\dot{V}O_2$  matching. Future honing of these interventions to maximize their impact on the  $\dot{V}O_2$  slow component might improve sports performance in athletes and exercise tolerance in the elderly or in patient populations. **Key Words:** SKELETAL MUSCLE, METABOLISM, ENERGETICS, EFFICIENCY, FATIGUE, EXERCISE TOLERANCE

After the commencement of constant-work-rate (CWR) exercise situated below the so-called lactate threshold (LT) or gas exchange threshold (GET), pulmonary  $O_2$  uptake ( $\dot{V}O_2$ ) rises relatively rapidly to attain a new steady state within a few minutes of exercise onset (90,128). If the work rate is above the LT, however, the attainment of a steady state is at least delayed (102,128,132) owing to the emergence of a supplementary, slowly developing component of the  $\dot{V}O_2$  response (14,95). When the work rate is below the “critical

power” (CP) (68), the eventual  $\dot{V}O_2$  steady state is greater than the value that would be predicted from the sub-LT  $\dot{V}O_2$ –work rate relationship; when the work rate is above the CP, no steady state is achievable but, rather,  $\dot{V}O_2$  continues to rise with time until the  $\dot{V}O_{2\max}$  is reached, heralding the imminent termination of exercise (102,126,128) (Fig. 1). These characteristic  $\dot{V}O_2$  profiles have been used to classify the various exercise intensity domains, namely, “moderate” (<LT), “heavy” (>LT but <CP), and “severe” (>CP) (100,102,132). Whereas the  $\dot{V}O_2$  response to <LT exercise is well described, after the exclusion of phase 1, by a monoexponential function, the  $\dot{V}O_2$  response to >LT exercise has been shown to be better described by biexponential processes with the second term being of delayed onset (14). This suggests a time-dependent, as well as intensity-dependent, loss of muscle efficiency as >LT exercise proceeds.

This “slow component” of  $\dot{V}O_2$  is not trivial: in the severe domain, its magnitude can exceed  $1 \text{ L} \cdot \text{min}^{-1}$  and represent  $\geq 25\%$  of the total increase in  $\dot{V}O_2$  above the preexercise baseline (98). The  $\dot{V}O_2$  slow component is therefore distinct from the relatively modest “ $O_2$  drift” (<200 mL  $O_2$ ) that may attend moderate exercise of a prolonged duration (i.e.,  $\geq 60$  min). On a purely academic level, the  $\dot{V}O_2$  slow component phenomenon is of interest because its study is likely

This review is based on the symposium “The Slow Component of  $\dot{V}O_2$  Kinetics: Mechanistic Bases and Practical Applications” presented at the 2010 ACSM Annual Meeting in Baltimore, Maryland.

Address for correspondence: Andrew M. Jones, Ph.D., Sport and Health Sciences, College of Life and Environmental Sciences, St. Luke's Campus, University of Exeter, Heavitree Road, Exeter, EX1 2LU, United Kingdom; E-mail: a.m.jones@exeter.ac.uk.

Submitted for publication January 2011.

Accepted for publication April 2011.

0195-9131/11/4311-2046/0

MEDICINE & SCIENCE IN SPORTS & EXERCISE®

Copyright © 2011 by the American College of Sports Medicine

DOI: 10.1249/MSS.0b013e31821f6fc1




FIGURE 1—Schematic framework illustrating the approximate location of key parameters of aerobic function (LT/GET, CP, and  $\dot{V}O_{2max}$ ) on the incremental/ramp exercise test (upper left panel) and demonstrating the profiles of  $\dot{V}O_2$  (upper middle panel) and blood lactate concentration (upper right panel) to CWR exercise in the moderate (<LT/GET), heavy (>LT/GET<CP), and severe (>CP to  $\dot{V}O_{2max}$ ) exercise intensity domains. Arrows depict point of exhaustion during severe-intensity exercise. Hatched areas denote  $\dot{V}O_2$  slow component that increases  $\dot{V}O_2$  above that predicted for the work rate from the  $\dot{V}O_2$ –work rate relation in the upper left panel (i.e., unloaded cycling  $\dot{V}O_2$  + ~10 mL O<sub>2</sub>·min<sup>-1</sup>·W<sup>-1</sup>). Note that, for heavy exercise (>LT/GET<CP), the  $\dot{V}O_2$  slow component can be stabilized at a submaximal  $\dot{V}O_2$ ; however, for severe exercise (>CP), the slow component drives  $\dot{V}O_2$  to  $\dot{V}O_{2max}$ . Lower panel illustrates the intensity domain dependency of the  $\dot{V}O_2$  slow component with its maximum value being achieved just above CP. For higher work rates, the  $\dot{V}O_2$  slow component decreases because the  $\dot{V}O_{2max}$  represents the upper limit for its expression.

to enhance our basic understanding of muscle energetics, metabolic control, and the determinants of the efficiency of skeletal muscle contraction. From a more functional perspective, the  $\dot{V}O_2$  slow component is of importance because it seems to be closely related to the progressive loss of muscle homeostasis and the development of fatigue that is evident during >LT exercise (23,68,111). Study of the  $\dot{V}O_2$  slow component may therefore provide insights into the determinants of exercise (in)tolerance in both healthy and patient populations; this, in turn, might aid the development of pharmacological, nutritional or exercise-related interventions with the potential to enhance human performance, health and well-being.

This invited review reflects the proceedings of a symposium of the same title that was presented at the 2010 American College of Sports Medicine meeting in Baltimore. Surprisingly, the last American College of Sports Medicine symposium that focused specifically on the  $\dot{V}O_2$  slow component was presented in 1993 (98). More than 250 scientific articles on this topic have been published since that date, and hence, the purpose of this review was to provide a contemporary overview of what is known, and what is still not known, about the  $\dot{V}O_2$  slow component. The review consists of four interrelated parts. First, Dr. David C. Poole provides the historical background to the discovery of the  $\dot{V}O_2$  slow component and its eventual recognition as an essential feature of the physiological response to high-intensity exercise.

Second, Dr. Bruno Grassi outlines key findings concerning the mechanistic bases of the  $\dot{V}O_2$  slow component emanating from experiments using the isolated *in situ* dog gastrocnemius preparation. Third, Peter M. Christensen, M.Sc., and Drs. Peter Krstrup and Jens Bangsbo review evidence from invasive studies in human volunteers, which provide insight into the role of muscle fiber type and fiber recruitment on the  $\dot{V}O_2$  slow component. Finally, Dr. Andrew M. Jones discusses the practical interventions that can attenuate the  $\dot{V}O_2$  slow component and thus predispose to enhanced exercise tolerance and considers the implications of these effects for understanding the physiological underpinnings of the  $\dot{V}O_2$  slow component.

## $\dot{V}O_2$ SLOW COMPONENT: HISTORY AND SIGNIFICANCE

When one reads research papers and standard texts from much of the 20th century, the control of pulmonary  $\dot{V}O_2$  (and, by inference, muscle  $\dot{V}O_2$ ) kinetics is assumed to be a linear first-order system. This characterization implies that ATP– $\dot{V}O_2$  coupling is rate-limited by a single first-order reaction (i.e., the mitochondrial creatine kinase reaction) (126) such that  $\dot{V}O_2$  kinetics is not rate-limited by O<sub>2</sub> transport *per se*. Several “fundamental” tenets of exercise physiology are founded on this notion. Specifically: 1)  $\dot{V}O_2$  increases as a unitary function of work rate (i.e., the gain = 9–11 mL

$\text{O}_2 \cdot \text{min}^{-1} \cdot \text{W}^{-1}$  for cycle exercise) across the range of achievable  $\dot{V}\text{O}_2$  values. 2) Irrespective of work intensity, the  $\dot{V}\text{O}_2$  profile ascribes to a single exponential process. 3) The time constant ( $\tau$ , time-to-reach 63% of the response as well as the mean response time, MRT) is invariant with work rate. 4) The  $\text{O}_2$  deficit can simply be calculated as  $\Delta\dot{V}\text{O}_2\tau$ . An important consequence of the acceptance of item 1) is the almost-ubiquitous tradition of characterizing exercise “intensity” as percent  $\dot{V}\text{O}_{2\text{max}}$ .

There is much support for the first-order linearity of  $\dot{V}\text{O}_2$  from muscles contracting *in vitro* (frog sartorius) (60,92) and *in situ* (dog gastrocnemius) (97). Furthermore, Riggs’s (108) concept of “superposition” provides a rigorous test for the first-order linearity of  $\dot{V}\text{O}_2$ . Considering the pulse/impulse as the input function, for which the first integral is the step and the second is the ramp, across these different work-forcing functions (i.e., pulse, step, ramp), the parameters of the  $\dot{V}\text{O}_2$  (output) (i.e.,  $\tau$ , MRT, gain) should be identical. However, empirical evidence has only supported the first-order linearity of  $\dot{V}\text{O}_2$  for exercise within the moderate-intensity domain (i.e., below the LT or GET). Crucially, for higher work rates in the heavy (>LT/GET) and severe (>CP) domains, a slow component of the  $\dot{V}\text{O}_2$  kinetics emerges, which becomes superimposed on the underlying and faster primary or fundamental response (Fig. 1). Thus, the  $\dot{V}\text{O}_2$  slow component, which may amount to as much as 1000–1500  $\text{mL O}_2 \cdot \text{min}^{-1}$ , challenges our understanding of muscle energetics and the foundational tenets of exercise physiology listed above. First, for CWR exercise in the heavy or severe domains, the end-exercise gain may increase considerably above 9–11  $\text{mL O}_2 \cdot \text{min}^{-1} \cdot \text{W}^{-1}$  (57,112). Second, the  $\dot{V}\text{O}_2$  kinetics are no longer well fit by a single exponential response. Third, the MRT increases. Fourth, confidence in the calculation of the  $\text{O}_2$  deficit is eroded, in part, by lack of formal characterization (i.e., linear, exponential, or other) (14,95,129) of the  $\dot{V}\text{O}_2$  slow component and the presumption that the end-exercise or steady-state  $\dot{V}\text{O}_2$  is the appropriate frame of reference (15,127). Finally, as  $\dot{V}\text{O}_2$  increases as a function of time during CWR exercise, the practice of describing a given exercise intensity relative to  $\dot{V}\text{O}_{2\text{max}}$  is fatuous (see also Jones et al. [68]).

It is extraordinary that the technology to measure  $\dot{V}\text{O}_2$  and thus identify the  $\dot{V}\text{O}_2$  slow component has been in existence for more than a century. However, because the  $\dot{V}\text{O}_2$  slow component has been inexplicable based on our conventional understanding of muscle energetics, its presence has been either systematically ignored or explained away. Perhaps the first evidence for the existence of the  $\dot{V}\text{O}_2$  slow component emerged in 1913 in the data of Krogh and Lindhard (79); however, it escaped detailed scientific scrutiny at that time. A decade later, in 1923, Hill and Lupton (61) reported that, for one subject,  $\dot{V}\text{O}_2$  increased by more than 0.5  $\text{L} \cdot \text{min}^{-1}$  between the 4th and 27th minute of treadmill running. Possibly because this diverged from contemporary models of energetics, it was explained away as the result of “a painful blister on the foot causing inefficient

movement” (61). Moreover, published in 1961, Astrand and Saltin’s (4) profiles of  $\dot{V}\text{O}_2$  during fatiguing high-intensity exercise demonstrate clearly  $\dot{V}\text{O}_2$  slow components that drive  $\dot{V}\text{O}_2$  to  $\dot{V}\text{O}_{2\text{max}}$  and portend exhaustion. Unfortunately, in the *Textbook of Work Physiology* (3), just two pages before reproduction of those actual  $\dot{V}\text{O}_2$  profiles,  $\dot{V}\text{O}_2$  is depicted fictitiously as rising in a strictly linear fashion with work rate to  $\dot{V}\text{O}_{2\text{max}}$  (3, p. 300–302). While this profile may be observed with the rapidly incremented or ramp test (Fig. 1, upper left panel), it is certainly not evident in any situation where the  $\dot{V}\text{O}_2$  slow component has time to emerge (Fig. 1, middle upper and lower panels). Another important consequence of the  $\dot{V}\text{O}_2$  slow component evident from Astrand and Saltin’s (4) and subsequent publications (e.g., Åstrand and Rodahl (3) and Poole et al. [102]) is that its capacity to drive  $\dot{V}\text{O}_2$  to  $\dot{V}\text{O}_{2\text{max}}$  extends the range of work rates at which  $\dot{V}\text{O}_{2\text{max}}$  may be achieved, provided that exercise is continued either to exhaustion or at least for a sufficient duration for the  $\dot{V}\text{O}_2$  slow component to develop fully. This property of >CP exercise has defined the extremities of the severe exercise intensity domain. Specifically, CP is the highest power output (or, more correctly, metabolic rate) (12) for which  $\dot{V}\text{O}_2$  can be stabilized below  $\dot{V}\text{O}_{2\text{max}}$  (68). By definition, all severe-intensity work rates (i.e., >CP) drive  $\dot{V}\text{O}_2$  to  $\dot{V}\text{O}_{2\text{max}}$ . Beyond the upper limit of the severe domain, what has been termed *extreme* (62,132) is characterized as exercise that is so intense that exhaustion intervenes before the kinetics of  $\dot{V}\text{O}_2$  allows  $\dot{V}\text{O}_{2\text{max}}$  to be achieved (this domain is omitted from Fig. 1 for clarity).


Despite some scientists’ contention that the  $\dot{V}\text{O}_2$  response could and should be described by a single exponential process that is invariant with exercise intensity (33,93), by the mid 1970s, the compelling weight of evidence demonstrated that, for heavy and severe exercise,  $\dot{V}\text{O}_2$  kinetics (on- and off-transient) become more complex, and the overall response is slowed in comparison with moderate exercise (Fig. 1) (58,90,128). Recognition that the  $\dot{V}\text{O}_2$  slow component was a reproducible event and of sufficient magnitude to contribute importantly to overall exercise energetics and possibly exercise tolerance led Hagberg et al. (51) to investigate its mechanistic bases. Using presumptive estimates of the  $\text{O}_2$  cost of the respiratory muscles and elevations in body temperature (the so-called “ $\text{Q}_{10}$  effect”), they concluded that these two mechanisms could account for essentially all of the  $\dot{V}\text{O}_2$  slow component (review Poole et al. [98,102]). Irrespective of this conclusion, recognition that the  $\dot{V}\text{O}_2$  slow component occurs only above LT/GET (57,112,129) and that many other potentially calorogenic processes are manifested simultaneously justified consideration of multiple additional putative mediators. These included lactate itself (via its stimulation of gluconeogenesis and other mechanisms); exercising muscle temperature (the principal site of any  $\text{Q}_{10}$  effect) (129); catecholamines; respiratory, cardiac, and auxiliary muscle work (e.g., arms for stabilization during cycle ergometry); and reduced contractile

efficiency of higher-order fibers (fast twitch, type IIa/b/d/x) (review Poole et al. [98]). Further correlative studies supported that the profile of blood lactate accumulation was more closely related to that of the  $\dot{V}O_2$  slow component than were ventilation, HR, body temperature or catecholamines (28,102).

A crucial advance in resolving the mechanistic bases for the  $\dot{V}O_2$  slow component was use of the constant-infusion thermodilution technique, pioneered by Andersen and Saltin (2), to make high-fidelity leg (and contracting muscle) blood flow and  $\dot{V}O_2$  measurements (101). Using that technique to compartmentalize the  $\dot{V}O_2$  slow component, Poole et al. (101) determined that the dominant portion (>80%) arose from within the contracting muscles, thereby truncating the list of candidate mechanisms to those extant at that site (Fig. 2). As a result, focused subsequent experiments disqualified muscle temperature (75), catecholamines (42), and increased muscle or blood lactate concentration (99) as viable mediators of the  $\dot{V}O_2$  slow component and emplaced muscle fiber energetics and recruitment patterns center stage (Fig. 2). Subsequent studies have confirmed that the dominant portion of the  $\dot{V}O_2$  slow component derives from within the contracting muscles (74,83). Using  $^{31}\text{P}$  magnetic resonance spectroscopy, Rossiter et al. (111) established that the  $\dot{V}O_2$  slow component was associated with a slow component of muscle phosphocreatine concentration ([PCr]), indicating that the slow component is linked to a greater ATP cost of force production rather than an elevated  $\dot{V}O_2$  cost of ATP production (i.e.,  $\Delta\text{P}/\text{O}$  ratio) *per se*.

There is substantial evidence in the animal (rat [44,106, 125] and mouse [32]) and human (30,52,118) literature that fast-twitch fibers (type II), compared with slow-twitch (type I) fibers, have a greater ATP cost for contractile activ-

ity owing to different chemical-to-mechanical coupling efficiencies, more rapid actomyosin turnover, far faster calcium pump activity (120), and less efficient FAD versus NAD-linked  $\alpha$ -glycerolphosphate shuttle activity (135). Thus, while extrapolations from *in vitro* to *in vivo* energetics should be made with caution (55), there is a solid foundation for the hypothesis that exercise, which recruits a greater proportion of type II fibers, will require a greater ATP (and thus  $\dot{V}O_2$ ) cost. Empirical testing of this hypothesis has been hampered by limitations including poor signal-to-noise fidelity and spatial resolution issues with existing technologies (e.g., EMG, magnetic resonance imaging (MRI)). Notwithstanding these considerations, beginning with the observation of a weak correlation between increased iEMG activity and  $\dot{V}O_2$  slow component amplitude by Shinohara and Moritani (117), there is support, using experimental perturbations such as priming exercise (21,35), manipulations of pedal frequency (104), glycogen depletion (27,86), preferential blockade of slow-twitch fiber recruitment (84), and exercise training (114), for greater (presumably type II) fiber recruitment occurring in synchrony with the  $\dot{V}O_2$  slow component. Other EMG (19,94,96) and MRI-based studies (37,115) support this thesis. It is also pertinent that the  $\dot{V}O_2$  slow component is greater in individuals with a higher proportion of type II versus type I fibers in the musculus vastus lateralis (13,67,103). Moreover, in the rat, the  $\dot{V}O_2$  slow component is associated with a preferential increase in blood flow and presumably greater recruitment of low oxidative fast-twitch muscle fibers (type IIb/d/x) (29). It is becoming increasingly evident, however, that the  $\dot{V}O_2$  slow component need not be dependent on recruitment of more fibers *per se* but, rather, may be driven by metabolic processes occurring within fibers that are already recruited (124,138).



**FIGURE 2—An array of putative mediators of the  $\dot{V}O_2$  slow component.** Note that, at least in healthy individuals during cycle ergometry, the bulk of the  $\dot{V}O_2$  slow component derives from intramuscular sites within the exercising limbs. However, insofar as remote events (e.g., increased work of breathing, altered arterial O<sub>2</sub> pressures) may affect exercising muscle blood flow and therefore O<sub>2</sub> delivery distribution, they may affect the  $\dot{V}O_2$  slow component to a surprising degree. See text for further details.

## $\dot{V}O_2$ SLOW COMPONENT: LESSONS FROM ANIMAL MODELS


As discussed above, the “slow component” of  $\dot{V}O_2$  kinetics is often considered to be mainly caused by a progressive recruitment of fibers as exercise proceeds and those fibers initially recruited become fatigued. The main experimental evidence in favor of this concept derives from EMG (19) and MRI (37) studies, as well as from experiments in which glycogen depletion (86), and selective neural blockade of type I fibers (84) was obtained. As discussed in Jones et al. (67), an alternative (or concurrent) explanation for the  $\dot{V}O_2$  slow component could involve an increase in metabolic demands within the already recruited fibers that are fatiguing (see, e.g., Pringle et al. [104] and Scheuermann et al. [116]).

An excellent experimental model to discriminate between these hypotheses is presented by the isolated dog gastrocnemius preparation *in situ*, originally used by Piiper et al. (97) to study skeletal muscle  $\dot{V}O_2$  kinetics. Using this model, Grassi et al. have investigated the roles of convective (45,48) and diffusive (46)  $O_2$  delivery as limiting factors for the  $\dot{V}O_2$  kinetics, as well as the putative roles of pyruvate dehydrogenase activation (47), nitric oxide synthase (NOS) inhibition (49), and CK inhibition (50) in the regulation of  $\dot{V}O_2$  kinetics. In this preparation, all muscle fibers are maximally activated (tetanic contractions) from the beginning of the contraction period, by direct electrical stimulation of the motor nerve. Thus, a sequential activation of motor units (the mechanism supposedly responsible for the  $\dot{V}O_2$  slow component, see above) is, by definition, impossible. In this preparation, a “modulation” of the metabolic intensity of contractions (in terms of percentages of  $\dot{V}O_2$  peak) can be accomplished by modulating the frequency of the contractions: with one contraction every 2 s (45,46),  $\dot{V}O_2$  should correspond at “steady state” to  $\sim 60\%$  of  $\dot{V}O_2$  peak, whereas with one contraction every 1 s (48),  $\dot{V}O_2$  peak was reached.

Was a  $\dot{V}O_2$  slow component observed in these experiments? The answer is not straightforward. If we consider the “classic” slow component (that is a further increase in  $\dot{V}O_2$  with time above the expected steady state), there was no slow component in the experiments at a relatively low metabolic intensity (45,46), there was a slow component in a minority of the experiments at an intermediate metabolic intensity (47,49), and there was a slow component in the majority of the experiments at peak  $\dot{V}O_2$  (48). Interestingly, a  $\dot{V}O_2$  slow component was not observed in the experiments carried out at an intermediate metabolic intensity, but in which the muscle was pump-perfused with a markedly elevated constant blood flow (50). This may represent indirect evidence that the  $\dot{V}O_2$  slow component is related to  $O_2$  availability. It is possible that, under these experimental conditions, increased bulk muscle blood flow and/or a better local matching of blood flow to metabolic rate increased “metabolic stability” (139) and reduced fatigue development and the associated loss of muscle effi-

ciency. In contrast, in conditions of spontaneous blood flow adjustment, such as during exercise in humans, blood flow distribution to active fibers may be suboptimal and may contribute to fatigue development and the loss of muscle efficiency, which is reflected in the  $\dot{V}O_2$  slow component.

In the previously mentioned studies, the fact that the muscles significantly fatigued during the 3-min (or 4-min) contraction period was neglected. The values of the fatigue index (the ratio between force at the end of the contraction period and the initial force) ranged between 0.84 (45) and 0.64 (48). In other words, there was no  $\dot{V}O_2$  slow component or only a small  $\dot{V}O_2$  slow component, but in the presence of a significant fall (by  $\sim 15\%$ – $35\%$ ) in force output. When the  $\dot{V}O_2$  data from one of these studies (49) were taken and “normalized” per unit of force, there was a clear  $\dot{V}O_2$  slow component, the amplitude of which corresponded on average to  $\sim 20\%$  of the total amplitude of the  $\dot{V}O_2$  increase (138). Thus, in the “classic”  $\dot{V}O_2$  slow component, described in exercising humans during CWR exercise, the external work rate is maintained (possibly by recruiting additional fibers) at the expense of an increasing  $\dot{V}O_2$ . In the isolated muscle *in situ* model, on the other hand, the muscle cannot recruit additional fibers, force decreases as a consequence of fatigue, and  $\dot{V}O_2$  remains essentially constant (Fig. 3). This phenomenon was termed a *mirror image* of the slow component (138). The two scenarios have a common denominator, namely, a reduced efficiency of muscle contractions: constant mechanical power output, with an increasing  $\dot{V}O_2$ ; or, conversely, a substantially constant  $\dot{V}O_2$ , but with a decreasing force. The study of Zoladz et al. (138) therefore demonstrated that the reduced efficiency of muscle




**FIGURE 3**—In the “classic” slow component of  $\dot{V}O_2$  kinetics (*upper panels*), CWR exercise is maintained (possibly by recruiting additional fibers) at the expense of an increasing  $\dot{V}O_2$ . In the isolated muscle *in situ* model (138), the muscle cannot recruit additional fibers, force decreases as a consequence of fatigue and  $\dot{V}O_2$  is essentially constant (*lower panels*). The two scenarios have a common denominator, that is, a fatigue-induced reduced efficiency of muscle contractions.

contraction, and thus the putative mechanism responsible for the  $\dot{V}O_2$  slow component, is not necessarily related (or due) to a *progressive* recruitment of muscle fibers.

A reduced efficiency of muscle contractions is, in fact, typically associated with fatigue (11,136). The two phenomena have several common denominators, such as a decrease in the Gibbs free energy of ATP hydrolysis, a decrease in PCr and glycogen concentrations, as well as increases in  $[H^+]$ ,  $[ADP]$ ,  $[P_i]$ ,  $[IMP]$ ,  $[NH_3]$ , etc. (1,40). The  $\dot{V}O_2$  slow component, or its mirror image, would then be associated with (or be a consequence of) a lower level of “metabolic stability” (139). Good metabolic stability in skeletal muscles during rest-to-work transitions is associated with good exercise tolerance, and results, for a given increase in  $\dot{V}O_2$ , in a less pronounced decrease in PCr and the cytosolic phosphorylation potential, as well as in a less pronounced increase in  $[P_i]$ ,  $[ADP_{free}]$ ,  $[AMP_{free}]$ , and  $[IMP_{free}]$  (139).

The slow component of  $\dot{V}O_2$  kinetics is associated with a slow component of PCr kinetics (111), that is with an increased “phosphate cost” of force production, which would explain the reduced contractile efficiency. Thus, the  $\dot{V}O_2$  slow component would be associated with (or caused by) a decreased efficiency of the contractile machinery (increase of the ATP/power output ratio) rather than by a decreased efficiency of the ATP production system (increase in the  $\dot{V}O_2$ /ATP ratio). This means that a “CWR” exercise, that is, an exercise characterized by a constant external mechanical power output, would be associated, in the presence of progressive fatigue, with a progressively higher ATP turnover rate.


Is anything like the “mirror image” of the  $\dot{V}O_2$  slow component, which was observed in the isolated muscle preparation *in situ*, observed in exercising humans as well? The answer is yes. In the study by Ribeiro et al. (107), for example,




**FIGURE 4**—Mean  $\pm$  SEM power output, iEMG, and  $\dot{V}O_2$  profiles during 3 min of all-out exercise (open symbols) and work-matched severe-intensity CWR exercise performed to the limit of tolerance (closed symbols) on a cycle ergometer. A. Mean power profile during 3 min of all-out cycle exercise evidences an early peak before falling to attain a value that is not different from the CP (dashed horizontal line); the mean power profile for the CWR test is shown as a solid horizontal line. The corresponding  $\dot{V}O_2$  profiles are shown in B. Note that the  $\dot{V}O_{2max}$  (dashed horizontal line) is attained in both tests but is reached rapidly in the all-out test but only after the development of the  $\dot{V}O_2$  slow component during CWR exercise. When transformed into a “gain” (i.e.,  $\dot{V}O_2$ /power) (C), a  $\dot{V}O_2$  slow component phenomenon is observed for both modes of exercise and the gain is higher for all-out exercise. This  $\dot{V}O_2$  slow component-like response during all-out exercise occurs despite a substantial fall in iEMG over time, in contrast to the progressive increase in iEMG during CWR exercise (D). Data from Vanhatalo et al. (124).

the subjects had to keep their pulmonary  $\dot{V}O_2$  constant during cycling exercise carried out for 40 min at  $\sim 55\%$ ,  $\sim 65\%$ , and  $\sim 75\%$  of the previously determined  $\dot{V}O_{2max}$ . To do so, the subjects had to decrease their mechanical power output. The decrease was linearly related to exercise intensity, ranging from  $\sim 5\%$  at a  $\dot{V}O_2$  corresponding to  $\sim 55\%$  of  $\dot{V}O_{2max}$ , to  $\sim 15\%$  at a  $\dot{V}O_2$  corresponding to  $\sim 75\%$  of  $\dot{V}O_{2max}$ . In another study by Stoudemire et al. (119), the subjects had to maintain, during 30 min of running on a treadmill, pulmonary  $\dot{V}O_2$  at a constant level corresponding to that associated (during an incremental test) with a blood lactate concentration of  $\sim 4$  mM. To do so, the subjects had to progressively reduce (by  $\sim 15\%$  at the end of the 30-min period) their running speed. Most recently and perhaps most convincingly, Vanhatalo et al. (124) reported that the  $\dot{V}O_2$  slow component was greater in a 3-min all-out cycle test, in which peak power is attained within the first 10 s before progressively falling with time to attain the CP, compared with a 3-min maximal CWR test (124) (Fig. 4). The iEMG declined by 26% during the all-out test and increased by 60% during the CWR test from the first 30 s to the last 30 s of exercise (Fig. 4). The considerable reduction in muscle efficiency in the all-out test in the face of a progressively falling iEMG indicates that increased muscle activation including progressive fiber recruitment is not requisite for the  $\dot{V}O_2$  slow component to develop during voluntary exercise in humans. These effects are essentially identical to the “mirror image” of the  $\dot{V}O_2$  slow component, which was observed in the canine skeletal muscle preparation *in situ* (138).

The strategy of maintaining the highest work rate associated with steady-state levels of relevant variables, such as pulmonary  $\dot{V}O_2$ , HR, muscle and blood pH, and blood lactate concentration, is based on the “maximal lactate steady-state” (16) or “critical power” (68) concepts. In this



**FIGURE 5**—Muscle oxygen uptake at selected time points during 30 min of knee extensor exercise at 30 W with an untrained (filled bars) and a trained leg (open bars). Note that a pronounced slow component of muscle  $\dot{V}O_2$  is apparent for the untrained leg (12% from 2 to 3 min and a further 11% increase from 3 to 30 min) but is absent in the trained leg. Values are means  $\pm$  SEM ( $n = 6$ ). Data from Krstrup (80).



**FIGURE 6**—Schematic representation of the most likely determinants of muscular efficiency during dynamic exercise. The results obtained in knee extensor exercise and cycle studies (39,82,84–86) suggest that muscle fiber recruitment influences the  $O_2$  cost as well as ATP turnover during dynamic exercise, whereas muscle temperature, lactate, and pH in the investigated range have little or no effect. Further studies are required to elucidate the precise effects of dynamic exercise on the P/O ratio.

scenario, the strategy of maintaining the highest work rate associated with reasonably good metabolic stability and efficiency of muscle contractions seems to represent a key factor for successful endurance performance. High-class marathon runners, for example, perform their marathon race at the highest possible running velocity at which they can maintain, throughout the race, unchanged blood pH (140) and presumably the lowest energy cost of running. It seems to represent “common sense,” in biological terms, that a system tries to maintain, as far as is possible, conditions of metabolic steady states to prevent (or minimize) the occurrence of fatigue. In this respect, the slow component of  $\dot{V}O_2$  kinetics (but also of other variables) represents a deviation from a steady state. The more pronounced this deviation, the more marked the consequences on fatigue and exercise (in)tolerance would likely be (23). As an example, Salvadego et al. (113) recently observed that, in obese adolescents exercising at 80%  $\dot{V}O_{2max}$ , the amplitude of the  $\dot{V}O_2$  slow component was linearly and inversely related to the time to exhaustion during the test.

## $\dot{V}O_2$ SLOW COMPONENT: ADDRESSING THE MECHANISTIC BASES IN HUMANS


Multiple studies have demonstrated that the contracting muscles are contributing to the pulmonary  $\dot{V}O_2$  slow component (74,83,101). It is also evident that the muscle  $\dot{V}O_2$  slow component represents a decline in muscular efficiency, which is ameliorated with training (80) (Fig. 5). It is, however, unclear what is causing the muscle  $\dot{V}O_2$  slow component and thus the lowering of muscle work efficiency. As illustrated in Figure 6, it has been suggested to be caused by a progressive increase in muscle temperature, acidosis, or changes in fiber-type recruitment and/or mitochondrial P/O ratio. These possibilities will be discussed in this section.

As mentioned previously, an increase in muscle temperature does not seem to influence the muscle  $\dot{V}O_2$  slow component to a significant extent (39,85). In one study, submaximal knee extensor exercise (43 W) was performed for 10 min with and without prior passive heating, and no differences were observed in muscle  $\dot{V}O_2$  or total energy turnover despite a 1.5°C higher mean quadriceps muscle temperature in the heated trial compared with control (37.9°C vs 36.4°C) (39). In another study, subjects cycled for 20 min at moderate and heavy exercise intensities on separate days eliciting 50% and 80%  $\dot{V}O_{2max}$ , respectively (85). A  $\dot{V}O_2$  slow component was only observed during the intense work with an increase in  $\dot{V}O_2$  from 2.62 L·min<sup>-1</sup> after 3 min of exercise to 2.76 L·min<sup>-1</sup> after 6 min of exercise with a further increase to 2.87 L·min<sup>-1</sup> after 20 min of cycling. However, there was no difference in the change of muscle temperature between the moderate and heavy exercise, being

1.0°C versus 1.1°C, respectively, between 3 and 6 min and 1.1°C versus 1.3°C from 6 to 20 min.

Muscle pH was only lowered during the intense cycling in which the  $\dot{V}O_2$  slow component was observed, indicating that muscular acidosis may be causing the decrease in efficiency. A number of studies, however, suggest that the relationship is not causal. Raising the lactate concentration through direct infusion of lactate in working dogs did not alter  $\dot{V}O_2$  (99). Likewise, no effect on the  $\dot{V}O_2$  slow component was observed after infusion of adrenaline in humans, which increased blood lactate concentration and reduced pH (42).

Change in the muscle fiber recruitment pattern has been studied by measurements of single muscle fiber content of PCr and glycogen. In the study by Krstrup et al. (85), the glycogen content in type I fibers decreased both during the moderate and heavy exercise bout, whereas the content in type II fibers only decreased during the intense exercise bout where the  $\dot{V}O_2$  slow component was observed. The PCr levels in individual fibers from muscle biopsies taken before and after 3, 6, and 20 min of exercise provided additional insight into the fiber recruitment pattern. At rest, approximately 10% of both the type I and type II fibers were below mean resting values minus 1 SD, and after moderate exercise, 25% of the type I fibers were below the mean resting value minus 1SD, whereas little change was observed for the type II fibers. After 3, 6, and 20 min of the intense exercise, 37%, 70%, and 74% of the type I fibers, respectively, and 45%, 83%, and 74% of the type II fibers, respectively, had PCr levels below mean resting values minus 1SD. These findings indicate that type I fiber recruitment dominated during moderate-intensity exercise whereas



**FIGURE 7—A,** Thigh  $\dot{V}O_2$  before and during 10 min of knee extensor exercise at 30 W with (open symbols) and without (full symbols) femoral arterial injections of a neuromuscular blocking agent (CUR). **B,** type I (ST) and type II (FT) fiber phosphocreatine (CP) breakdown during CUR and Control. Values are means  $\pm$  SEM ( $n = 6$ ). \*CUR versus CON. §Significant CP breakdown. #FT versus ST fibers. All  $P < 0.05$ . Data from Krstrup et al. (84).




both type I and type II fibers were involved during heavy-intensity exercise. Furthermore, the data demonstrate that more type II fibers were recruited from 3 to 6 min where the  $\dot{V}O_2$  slow component was apparent.

As type II fibers seem to be less efficient than the type I fibers during dynamic exercise in humans, the  $\dot{V}O_2$  slow component may be due to the progressive recruitment of type II fibers (80,84–86). Thus, Krstrup et al. (84) used the neuromuscular blocking agent cisatracurium (curare analog, CUR) to impair the activation of the type I fibers during 10 min of one-legged knee extensor exercise bouts at a moderate intensity (30 W). Measurements of PCr levels in single fibers confirmed that fewer type I fibers were active with CUR infusion compared with the control situation (Fig. 7). Muscle  $\dot{V}O_2$  was around  $100 \text{ mL} \cdot \text{min}^{-1}$  higher with infusion of CUR and mechanical efficiency was lower (16 vs. 20%) compared with control, showing

that a greater recruitment of type II fibers leads to a higher muscle  $\dot{V}O_2$  (Fig. 7).

Additional support for recruitment of type II fibers playing a crucial part in the development of the  $\dot{V}O_2$  slow component is found in a study in which the glycogen levels were manipulated before whole body exercise (86). Subjects performed 20 min of moderate-intensity cycling at 50%  $\dot{V}O_{2\text{max}}$  on three separate days. On two occasions, the subjects carried out 3 h of low-intensity cycling (40%  $\dot{V}O_{2\text{max}}$ ) the day before the experiment to lower the glycogen levels in the type I fibers. On one of these occasions, the subjects fasted overnight (CHO-DEP), and on the other, they ingested a diet with a high CHO content to examine whether the exercise the day before influenced the response (CHO-RE). In the third condition, the subjects rested the day before testing and had an overnight fast. Analysis of muscle biopsies obtained before testing revealed that the prior exercise



**FIGURE 8—**Distribution of single-fiber phosphocreatine (CP) content during 20 min of cycling at 50%  $\dot{V}O_{2\text{max}}$  for type I (ST) fibers (Panel A) and type II (FT) fibers (Panel B) in a control condition (CON) and after prior glycogen depletion of ST fibers caused by 3 h of cycling at 40%  $\dot{V}O_{2\text{max}}$  the day before testing (CHO-DEP). Panel C shows pulmonary  $\dot{V}O_2$  during cycling at 50%  $\dot{V}O_{2\text{max}}$  in CON and after CHO-DEP ( $n = 12$ ) while Panel D shows pulmonary  $\dot{V}O_2$  in CON and after refuelling of the CHO stores (CON-RE;  $n = 6$ ).

and fasting (CHO-DEP) had depleted the glycogen stores of most type I fibers. Muscle fiber recruitment pattern during the 20-min exercise bout was assessed by measuring glycogen levels before and after exercise as well as PCr levels in individual fibers after 0, 3, and 20 min of cycling. A markedly higher type II fiber recruitment was seen in CHO-DEP compared with the control condition (Fig. 8A, B). The pulmonary  $\dot{V}O_2$  after 3 min of cycling was higher in CHO-DEP than in control, and in contrast to the control situation, a  $\dot{V}O_2$  slow component ( $\sim 110 \text{ mL} \cdot \text{min}^{-1}$ ) was observed in CHO-DEP (Fig. 8C). The notion that the  $\dot{V}O_2$  slow component was caused by the exercise performed the day before testing could be dismissed because the  $\dot{V}O_2$  response after the CHO refueling strategy was the same as in the control condition (Fig. 8D). Thus, these findings underline that the  $\dot{V}O_2$  slow component during CWR exercise is related to a greater recruitment of type II fibers. In addition, muscle pH was not changed during either of the exercise bouts, demonstrating that a  $\dot{V}O_2$  slow component can be observed without changes in muscle acidity.

The effect of manipulating the glycogen content of type I and type II fibers on the  $\dot{V}O_2$  slow component has been studied by others. Carter et al. (27) had subjects cycle at a moderate (80% GET) and heavy intensity (50% of the difference between GET and  $\dot{V}O_{2\text{max}}$ ) after glycogen depletion of either primarily type I fibers (3 h of cycling at 30%  $\dot{V}O_{2\text{max}}$ ) or type II fibers ( $10 \times 1 \text{ min}$  at 120%  $\dot{V}O_{2\text{max}}$ ). After glycogen depletion of the type II fibers, the  $\dot{V}O_2$  slow component during heavy exercise was reduced compared with control, in agreement with Krstrup et al. (86). However, in contrast to the findings from Krstrup et al. (86), neither of the manipulations had any effect on  $\dot{V}O_2$  during moderate-intensity exercise. One explanation may be that, in the study of Carter et al. (27), the exercise duration was only 6 min compared with 20 min used by Krstrup et al. (86), in which the  $\dot{V}O_2$  slow component was significant after approximately 15 min of exercise. Bouckaert et al. (20) lowered the glycogen content in the type I fibers by having their subjects eat a meal without CHO (60% fat and 40% protein) and cycle for 2 h at 60%  $\dot{V}O_{2\text{max}}$ . Twelve hours after the exercise without food, the subjects exercised for 9 min at 85%  $\dot{V}O_{2\text{max}}$ . In comparison with the control condition, the efficiency was reduced after the glycogen depletion protocol with an increase in  $\dot{V}O_2$  of approximately  $140 \text{ mL} \cdot \text{min}^{-1}$ , consistent with the findings by Krstrup et al. (86). The magnitude of the  $\dot{V}O_2$  slow component, however, was not changed by the intervention.

Interestingly, when a transition to a heavy or severe-intensity work rate is initiated from a moderate-intensity work rate (as opposed to a baseline of “unloaded” cycling), the overall  $\dot{V}O_2$  kinetics is slower and the response gain is greater, i.e., efficiency is impaired (133,134); reciprocal effects are observed in the intramuscular [PCr] profiles (35,72). These effects do not seem to be related to differences in muscle  $O_2$  delivery (34,36). This “work-to-work” model is useful in exploring differences in fiber-type ener-

getics because it theoretically allows the  $\dot{V}O_2$  response of different segments of the fiber-type recruitment hierarchy to be isolated (56). The different  $\dot{V}O_2$  profiles that are elicited in these conditions (34–36,133,134) also suggest that type II fibers have slower  $\dot{V}O_2$  kinetics and lower efficiency compared with type I fibers. On this basis, a theoretical model that attempts to link fiber recruitment patterns to the different  $\dot{V}O_2$  profiles in the various exercise intensity domains, including the delayed and elevated steady state observed for heavy exercise and the lack of steady state for severe exercise has been advanced (see Fig. 6 of Wilkerson and Jones [134]).

Little is known about the mitochondrial P/O ratio during dynamic exercise in humans. However, there is indirect evidence that the mitochondria play an important role in determining muscular efficiency and it could be speculated that P/O ratio is changed over time owing to progressive increases in intracellular  $\text{Ca}^{2+}$  and other electrolytes. Under ischemic conditions, no changes were observed in muscle efficiency during a 90-s bout of knee extensor exercise at 30 W, whereas efficiency declined during a similar bout of exercise in free-flow conditions (82). A gradual lowering of P/O ratio specifically in the type II fibers, which are expected to develop higher intracellular  $\text{Ca}^{2+}$  concentration than type I fibers, may also explain why a higher  $\dot{V}O_2$  was only apparent after 2 min of exercise in the CUR experiments (84) (Fig. 7).

Overall, it seems that recruitment of additional muscle fibers and a change in fiber-type recruitment toward type II fibers can play an important role in the development of the muscle  $\dot{V}O_2$  slow component during intense CWR exercise, whereas changes in muscle temperature and acidity seem to be of little importance. *In vitro* studies using isolated mitochondria from human muscle collected after fatiguing exercise indicate that there are no long-lasting effects on the mitochondrial P/O ratio (121). However, further studies are required to elucidate the precise role of the P/O ratio for the observed difference in efficiency between fiber types as well as for the development of the  $\dot{V}O_2$  slow component during dynamic exercise in humans (105).

## **$\dot{V}O_2$ SLOW COMPONENT: PRACTICAL SIGNIFICANCE AND APPLICATIONS**

Several lines of evidence indicate that the development of the  $\dot{V}O_2$  slow component is intimately related to the muscle fatigue process and that exercise tolerance can be improved by interventions which serve to reduce or eliminate the  $\dot{V}O_2$  slow component. For example, the temporal profiles of intramuscular [ADP] and [ $P_i$ ] (increasing) and [PCr] and pH (decreasing) are exercise intensity domain-dependent and, as such, become progressively perturbed in concert with the development of the  $\dot{V}O_2$  slow component (35,71,111). Thus, during severe-intensity exercise, the  $\dot{V}O_2$  slow component drives  $\dot{V}O_2$  to  $\dot{V}O_{2\text{max}}$  and exercise is terminated as dictated by the parameters of the power–time

relationship for high-intensity exercise (i.e., CP and  $W'$ , where  $W'$  represents the finite amount of work that can be done above CP) (23,68,102). Indeed, it has been suggested that the magnitude of both the  $\dot{V}O_2$  slow component and the  $W'$  are a function of the difference between the  $\dot{V}O_2$  at CP and the  $\dot{V}O_{2max}$ ; that is, there is a reciprocal relationship between the development of the  $\dot{V}O_2$  slow component and the progressive reduction in the  $W'$  (23,122). Given this relationship between the development of the  $\dot{V}O_2$  slow component and the progressive loss of muscle metabolic homeostasis, it is important to appreciate which interventions have the potential to reduce the  $\dot{V}O_2$  slow component. This is not only of academic interest in providing insights into the mechanistic bases to the  $\dot{V}O_2$  slow component; it is also of practical import in devising strategies for enhancing performance in athletes and improving exercise tolerance in the aged or those with pulmonary, cardiovascular, or metabolic disorders.

**Training.** Fortunately, several such interventions exist, and perhaps the most potent of these is endurance exercise training. First, it is interesting to note that the  $\dot{V}O_2$  slow component tends to be relatively small in endurance athletes, presumably as a consequence of the LT and CP occurring at a high fraction of  $\dot{V}O_{2max}$  in this population (66). Second, it has been established that 6–8 wk of endurance training results in a significant reduction in the  $\dot{V}O_2$  slow component when subjects are tested at the same absolute work rate (26,28,137) (Fig. 5). This may also be related to the well-known effects of endurance training on the LT and CP and/or be linked to enhanced muscle blood flow (and/or its distribution) or muscle oxidative capacity (63). These latter adaptations, in turn, would be expected to improve metabolic stability, reduce the rate of fatigue development, and attenuate the requirement for higher-order motor units to be recruited to sustain power output. Consistent with this, it has been reported that 5 wk of endurance training results in an attenuated fall of muscle pH and a significant reduction in the amplitude of the intramuscular [PCr] slow component (69). Given that the termination of exercise in the severe domain coincides with the attainment of consistently low muscle [PCr] and pH values (71,122), these data suggest that the attenuation of the [PCr] slow component (and thus  $\dot{V}O_2$  slow component) might be mechanistically linked with enhanced exercise tolerance after endurance training.

Although it is well established that generalized endurance training reduces the  $\dot{V}O_2$  slow component, the specific type of training (i.e., volume, duration, intensity, etc.) that is optimal for this effect remains under investigation. For example, Berger et al. (18) compared the effects of endurance training (three to four sessions per week of 30-min duration at 60%  $\dot{V}O_{2max}$ ) with work-matched interval training (three to four sessions per week involving  $20 \times 1$ -min exercise bouts at 90%  $\dot{V}O_{2max}$ ) on  $\dot{V}O_2$  kinetics in previously untrained subjects and found that both types of training were similarly effective in reducing the  $\dot{V}O_2$  slow component. In contrast, Bailey et al. (8) reported that just 2 wk (six ses-

sions) of interval training involving  $4-7 \times 30$ -s cycle sprints significantly reduced the  $\dot{V}O_2$  slow component and enhanced severe-intensity exercise tolerance, whereas six sessions of work-matched, moderate-intensity cycling was ineffective. Similarly, 3 months of participation in an intense intermittent sport like football (soccer) resulted in a larger reduction of the  $\dot{V}O_2$  slow component compared with moderate-intensity running in previously untrained men (81). The efficacy of a specific training intervention will depend on factors such as initial fitness as well as the duration and other characteristics of the training program (63). Interestingly, it has recently been reported that 4 wk of pressure-threshold inspiratory muscle training, which reduced exercise-induced inspiratory muscle fatigue, significantly reduced the  $\dot{V}O_2$  slow component and enhanced exercise tolerance during severe- and extreme-intensity exercise (6). This effect may be consequent to an attenuation of the  $O_2$  and blood flow requirements of the respiratory muscles for a given rate of pulmonary ventilation, thereby attenuating the metaboreflex and enabling an increased limb  $O_2$  delivery (53). In this regard, it is of interest that reducing the work of breathing by having subjects inspire He- $O_2$  also reduces the  $\dot{V}O_2$  slow component (31).


**Priming.** In addition to the chronic interventions of endurance or inspiratory muscle training, the  $\dot{V}O_2$  slow component can also be attenuated using an acute bout of heavy or severe-intensity “priming” exercise (24,43,91). Although often termed “warm-up” exercise, the effects of prior exercise on  $\dot{V}O_2$  kinetics during a subsequent bout of high-intensity exercise cannot be attributed to the elevated muscle temperature *per se* (78). Rather, it is likely that enhancements of both bulk muscle blood flow and local matching of blood flow to  $\dot{V}O_2$ , as well as an increased activity of mitochondrial enzymes, after priming exercise retard the rate of fatigue development and the requirement for additional motor unit recruitment to maintain power output during subsequent exercise (7,21,89). It has been reported that priming exercise reduces the [PCr] slow component (41, 110), although contradictory data also exist (64). Whereas the potential for priming exercise to reduce the  $\dot{V}O_2$  slow component during subsequent exercise is well documented (e.g., 7,21,24,43,78,91,110), the effect on exercise tolerance is controversial, with some studies reporting an improvement (22,70) and others an impairment (38,131) of subsequent exercise performance.

In a comprehensive recent investigation, Bailey et al. (7) examined the interaction of previous exercise intensity (heavy and severe) and subsequent recovery interval (3, 9, and 20 min) on  $\dot{V}O_2$  kinetics and severe-intensity exercise tolerance. The effects on  $\dot{V}O_2$  kinetics were appreciably greater when severe (as opposed to heavy) priming exercise was performed. Exercise tolerance was not altered by heavy-intensity priming exercise. When the recovery interval separating the two severe-intensity exercise bouts was 3 min, exercise tolerance in the second bout was impaired by 16% relative to the control (no prior exercise) condition.

However, when the recovery interval was extended to 9 and 20 min, exercise tolerance was improved by 15% and 30%, respectively, relative to the control condition. These data indicate that prior severe-intensity exercise can enhance the tolerance to subsequent severe-intensity exercise provided that it is coupled with adequate recovery duration ( $\geq 9$  min). This combination, although almost certainly not uniquely efficacious, presumably optimizes the balance between preserving the positive effects of prior exercise on  $\dot{V}O_2$  kinetics and providing sufficient time for muscle [PCr] and pH to recover toward baseline values (38,123). It seems clear that athletes should carefully consider their precompetition preparation regimens to benefit from the potential of priming exercise to enhance performance.

The reduced  $\dot{V}O_2$  slow component observed after priming exercise in the study of Bailey et al. (7) was associated with a blunted increase of the muscle integrated EMG (iEMG) between 2 and 6 min of exercise. Other studies (21,89) have also reported changes in iEMG after priming exercise that may reflect changes in motor unit recruitment profiles. A diminished rate of fatigue development during exercise after priming (as a function, for example, of greater muscle  $O_2$  availability) might limit the requirement for additional motor unit recruitment. Alternatively, a greater iEMG in the early minutes of exercise, as has been observed in several studies (7,21,35,89), may indicate a reduced metabolic strain per recruited muscle fiber and may also be conducive to improved exercise performance. The influence of priming exercise on motor unit recruitment profiles likely reflects the balance between the potentially beneficial and detrimental effects of priming exercise on muscle function, effects which in turn will be related to the intensity of the priming exercise bout and the duration of the subsequent recovery interval (7).

**Nutrition.** Recent studies have reported that dietary nitrate supplementation in the form of beetroot juice, which increases plasma [nitrite] (an index of increased NO bioavailability), reduces the amplitude of the  $\dot{V}O_2$  slow component (5,9,88). The mechanistic bases for this effect remain unclear. The reduced  $\dot{V}O_2$  slow component is mirrored by a reduced intramuscular [PCr] slow component (9) (Fig. 9), suggesting that muscle contractile efficiency is enhanced by the intervention. Given the important role of NO in the regulation of blood flow, however, it is also possible that nitrate supplementation reduces the  $\dot{V}O_2$  slow component by improving the matching of  $O_2$  delivery to  $\dot{V}O_2$  in active motor units (5,73). Irrespective of the mechanistic bases for the effect, it seems that the attenuated muscle metabolic perturbation after nitrate supplementation allows high-intensity exercise tolerance to be enhanced (5,9,88). Interestingly, similar effects on the  $\dot{V}O_2$  slow component and exercise tolerance have been reported after ingestion of a dietary supplement containing L-arginine, the substrate for the “conventional” NO production pathway catalyzed by NOS (10), whereas the  $\dot{V}O_2$  slow component is increased when NOS is inhibited pharmacologically (73). Collectively, these



**FIGURE 9—Pulmonary  $\dot{V}O_2$  (A) and muscle [PCr] measured by  $^{31}P$  magnetic resonance spectroscopy (B) during high-intensity knee extension exercise after dietary supplementation with nitrate (closed symbols) or placebo (open symbols). Note the reduced  $\dot{V}O_2$  and [PCr] slow components after nitrate supplementation, along with the significantly extended time-to-exhaustion. Data from Bailey et al. (5).**

data suggest an important role for NO in modulating the development of the  $\dot{V}O_2$  slow component and thus affecting exercise tolerance.

Another nutritional intervention that has been reported to influence the  $\dot{V}O_2$  slow component is sodium bicarbonate ingestion: in one study, the  $\dot{V}O_2$  slow component amplitude was reduced (77), and in another, the onset of the  $\dot{V}O_2$  slow component was delayed and end-exercise  $\dot{V}O_2$  was reduced (17). This effect is presumably mediated by a greater efflux of  $H^+$  from muscle to blood, thereby reducing or delaying muscle fatigue and the recruitment of higher-order motor units. Consistent with this interpretation, infusion of the drug dichloroacetate, which activates the pyruvate dehydrogenase enzyme complex and reduces substrate-level phosphorylation, results in a reduced amplitude (109) or delayed onset (65) of the  $\dot{V}O_2$  slow component.

The previous discussion highlights that the  $\dot{V}O_2$  slow component is intimately related to intramuscular events and to the development of fatigue. In this regard, it is interesting that increasing arterial  $O_2$  partial pressure through the inspiration of hyperoxic gas mixtures markedly attenuates the  $\dot{V}O_2$  slow component (91,130) and also spares muscle PCr hydrolysis (54,122), an index of metabolic stability. In contrast, when exercise is performed in the supine position (76) and when muscle blood flow is reduced (87), the

$\dot{V}O_2$  slow component is increased. Collectively, these observations indicate that the  $\dot{V}O_2$  slow component can be manipulated by altering blood flow or  $O_2$  delivery to the active muscle(s). Again, this is presumably linked to greater metabolic stability, reduced fatigue, improved efficiency, and changes to fiber recruitment.

## CONCLUSIONS

The  $\dot{V}O_2$  slow component is a fundamental property of the metabolic response to exercise performed above the LT that has been excluded from mainstream textbooks and teaching in exercise physiology, presumably because its existence presents an inconvenient challenge to our understanding of muscle energetics. This is unfortunate given that the causes and consequences of the  $\dot{V}O_2$  slow component are so important for a more complete appreciation of metabolic control, muscle efficiency, and the determinants of exercise tolerance.

Mechanistically, the progressive loss of muscle efficiency represented by the  $\dot{V}O_2$  slow component is associated with the development of fatigue and can be offset by interventions that enhance metabolic stability (5–8, 9,18,21,26,28,91,130,137). During high-intensity CWR exercise, the “conventional”  $\dot{V}O_2$  slow component is associated with a progressive recruitment of additional higher-order (type II) muscle fibers, and the low efficiency of these fibers might well contribute to the increased  $O_2$  cost of exercise (67,84–87,133). However, fatigued fibers might also become less efficient and require a greater  $O_2$  consumption per unit of ATP turnover and/or a greater ATP turnover per unit of power output (11,136). Recent

data indicate that muscle efficiency is also lowered (resulting in a “mirror-image”  $\dot{V}O_2$  slow component) during fatiguing high-intensity exercise in which additional fiber recruitment is debarred (59,124,138). It is important to recognize, however, that the existence of a  $\dot{V}O_2$  slow component-like phenomenon in this situation does not necessarily challenge the strong evidence that the  $\dot{V}O_2$  slow component is linked in some fashion to the recruitment of type II muscle fibers (13,37,67,84–86, 103,104,134), only that the *progressive, additional* recruitment of these fibers is not obligatory for the slow component to develop (25,59,124,138). Type II fibers will be recruited from the onset of fatiguing high-intensity or all-out exercise and the greater fatigue sensitivity, and likely slower  $\dot{V}O_2$  kinetics and impaired efficiency of these fibers relative to type I fibers might still play a crucial role in the development of the  $\dot{V}O_2$  slow component (104,124,133,138).

From a practical perspective, further study of the mechanistic bases of the  $\dot{V}O_2$  slow component will be important in designing interventions for the enhancement of exercise performance. Although there is an obvious application to athletes, it is also clear that patient populations and the elderly, in whom daily mobility and the quality of life may be restricted because of a low or limited  $O_2$  transport capacity, might benefit tremendously from therapeutic strategies that reduce the  $\dot{V}O_2$  slow component. This is a worthy goal for future studies.

There was no funding received.

The authors report no conflict of interest.

The results of the present study do not constitute endorsement by the American College of Sports Medicine.

## REFERENCES

- Allen DG, Lamb GD, Westerblad H. Skeletal muscle fatigue: cellular mechanisms. *Physiol Rev*. 2008;88:287–332.
- Andersen P, Saltin B. Maximal perfusion of skeletal muscle in man. *J Physiol*. 1985;366:233–49.
- Åstrand PO, Rodahl K. *Textbook of Work Physiology: Physiological Basis of Exercise*. New York (NY): McGraw-Hill; 1986. p. 1–756.
- Åstrand PO, Saltin B. Maximal oxygen uptake and heart rate in various types of muscular activity. *J Appl Physiol*. 1961;16: 977–81.
- Bailey SJ, Fulford J, Vanhatalo A, et al. Dietary nitrate supplementation enhances muscle contractile efficiency during knee-extensor exercise in humans. *J Appl Physiol*. 2010;109: 135–48.
- Bailey SJ, Romer LM, Kelly J, Wilkerson DP, DiMenna FJ, Jones AM. Inspiratory muscle training enhances pulmonary  $O_2$  uptake kinetics and high-intensity exercise tolerance in humans. *J Appl Physiol*. 2010;109:457–68.
- Bailey SJ, Vanhatalo A, Wilkerson DP, Dimenna FJ, Jones AM. Optimizing the “priming” effect: influence of prior exercise intensity and recovery duration on  $O_2$  uptake kinetics and severe-intensity exercise tolerance. *J Appl Physiol*. 2009;107: 1743–56.
- Bailey SJ, Wilkerson DP, Dimenna FJ, Jones AM. Influence of repeated sprint training on pulmonary  $O_2$  uptake and muscle deoxygenation kinetics in humans. *J Appl Physiol*. 2009;106: 1875–87.
- Bailey SJ, Winyard P, Vanhatalo A, et al. Dietary nitrate supplementation reduces the  $O_2$  cost of low-intensity exercise and enhances tolerance to high-intensity exercise in humans. *J Appl Physiol*. 2009;107:1144–55.
- Bailey SJ, Winyard PG, Vanhatalo A, et al. Acute L-arginine supplementation reduces the  $O_2$  cost of moderate-intensity exercise and enhances high-intensity exercise tolerance. *J Appl Physiol*. 2010;109:1394–403.
- Barclay CJ. Mechanical efficiency and fatigue of fast and slow muscles of the mouse. *J Physiol*. 1996;497:587–96.
- Barker T, Poole DC, Noble ML, Barstow TJ. Human critical power–oxygen uptake relationship at different pedalling frequencies. *Exp Physiol*. 2006;91:621–32.
- Barstow TJ, Jones AM, Nguyen PH, Casaburi R. Influence of muscle fiber type and pedal frequency on oxygen uptake kinetics of heavy exercise. *J Appl Physiol*. 1996;81:1642–50.
- Barstow TJ, Molé PA. Linear and non-linear characteristics of oxygen uptake kinetics during heavy exercise. *J Appl Physiol*. 1991;71:2099–2106.
- Bearden SE, Moffatt RJ.  $\dot{V}O_2$  kinetics and the  $O_2$  deficit in heavy exercise. *J Appl Physiol*. 2000;88:1407–12.

16. Beneke R, von Duvillard SP. Determination of maximal lactate steady-state response in selected sports events. *Med Sci Sports Exerc.* 1996;28(2):241–6.
17. Berger NJ, McNaughton LR, Keatley S, Wilkerson DP, Jones AM. Sodium bicarbonate ingestion alters the slow but not the fast phase of  $\dot{V}O_2$  kinetics. *Med Sci Sports Exerc.* 2006;38(11):1909–17.
18. Berger NJ, Tolfrey K, Williams AG, Jones AM. Influence of continuous and interval training on oxygen uptake on-kinetics. *Med Sci Sports Exerc.* 2006;38(3):504–12.
19. Borrani F, Candau R, Millet GY, Perrey S, Fuchslocher J, Rouillon JD. Is the  $\dot{V}O_2$  slow component dependent on progressive recruitment of fast-twitch fibers in trained runners. *J Appl Physiol.* 2001;90:2212–20.
20. Bouckaert J, Jones AM, Koppo K. Effect of glycogen depletion on the oxygen uptake slow component in humans. *Int J Sports Med.* 2004;25:351–6.
21. Burnley M, Doust JH, Ball D, Jones AM. Effects of prior heavy exercise on  $\dot{V}O_2$  kinetics during heavy exercise are related to changes in muscle activity. *J Appl Physiol.* 2002;93:167–74.
22. Burnley M, Doust JH, Jones AM. Effects of prior warm-up regime on severe-intensity cycling performance. *Med Sci Sports Exerc.* 2005;37(5):838–45.
23. Burnley M, Jones AM. Oxygen uptake kinetics as a determinant of sports performance. *Eur J Sports Sci.* 2007;7:63–79.
24. Burnley M, Jones AM, Carter H, Doust JH. Effects of prior heavy exercise on phase II pulmonary oxygen uptake kinetics during heavy exercise. *J Appl Physiol.* 2000;89:1387–96.
25. Cannon DT, White AC, Andriano MF, Kolkhorst FW, Rossiter HB. Skeletal muscle fatigue precedes the slow component of oxygen uptake kinetics during exercise in humans. *J Physiol.* 2011;589:727–39.
26. Carter H, Jones AM, Barstow TJ, Burnley M, Williams C, Doust JH. Effect of endurance training on oxygen uptake kinetics during treadmill running. *J Appl Physiol.* 2000;89:1744–52.
27. Carter H, Pringle JS, Boobis L, Jones AM, Doust JH. Muscle glycogen depletion alters oxygen uptake kinetics during heavy exercise. *Med Sci Sports Exerc.* 2004;36(6):965–72.
28. Casaburi R, Storer TW, Ben-Dov I, Wasserman K. Effect of endurance training on possible determinants of  $\dot{V}O_2$  during heavy exercise. *J Appl Physiol.* 1987;62:199–207.
29. Copp SW, Hirai DM, Musch TI, Poole DC. Critical velocity in the rat: implications for hindlimb muscle blood flow distribution and fiber recruitment. *J Physiol.* 2010;588:5077–87.
30. Coyle EF, Sidossis LS, Horowitz JF, Beltz JD. Cycling efficiency is related to the percentage of type I muscle fibers. *Med Sci Sports Exerc.* 1992;24(7):782–8.
31. Cross TJ, Sabapathy S, Schneider DA, Haseler LJ. Breathing  $He-O_2$  attenuates the slow component of  $O_2$  uptake kinetics during exercise performed above the respiratory compensation threshold. *Exp Physiol.* 2010;95:172–83.
32. Crow MT, Kushmerick MJ. Chemical energetics of slow- and fast-twitch muscles of the mouse. *J Gen Physiol.* 1982;79:147–66.
33. di Prampero PE, Davies CTM, Cerretelli P, Margaria R. An analysis of  $O_2$  debt contracted in submaximal exercise. *J Appl Physiol.* 1970;29:547–51.
34. DiMenna FJ, Bailey SJ, Vanhatalo A, Chidnok W, Jones AM. Elevated baseline  $\dot{V}O_2$  per se does not slow  $O_2$  uptake kinetics during work-to-work exercise transitions. *J Appl Physiol.* 2010;109:1148–54.
35. DiMenna FJ, Fulford J, Bailey SJ, Vanhatalo A, Wilkerson DP, Jones AM. Influence of priming exercise on muscle [PCr] and pulmonary  $O_2$  uptake dynamics during “work-to-work” knee-extension exercise. *Respir Physiol Neurobiol.* 2010;172:15–23.
36. DiMenna FJ, Wilkerson DP, Burnley M, Jones AM. Influence of priming exercise on pulmonary  $O_2$  uptake kinetics during transitions to high-intensity exercise from an elevated baseline. *J Appl Physiol.* 2008;105:538–46.
37. Endo MY, Kobayakawa M, Kinugasa R, et al. Thigh muscle activation distribution and pulmonary  $\dot{V}O_2$  kinetics during moderate, heavy, and very heavy intensity cycling exercise in humans. *Am J Physiol Regul Integr Comp Physiol.* 2007;293:R812–20.
38. Ferguson C, Whipp BJ, Cathcart AJ, Rossiter HB, Turner AP, Ward SA. Effects of prior very-heavy intensity exercise on indices of aerobic function and high-intensity exercise tolerance. *J Appl Physiol.* 2007;103:812–22.
39. Ferguson RA, Krstrup P, Kjær M, Mohr M, Ball D, Bangsbo J. Effect of temperature on skeletal muscle energy turnover during dynamic knee-extensor exercise in humans. *J Appl Physiol.* 2006;101:47–52.
40. Fitts RH. The cross-bridge cycle and skeletal muscle fatigue. *J Appl Physiol.* 2008;104:551–8.
41. Forbes SC, Raymer GH, Kowalchuk JM, Thompson RT, Marsh GD. Effects of recovery time on phosphocreatine kinetics during repeated bouts of heavy-intensity exercise. *Eur J Appl Physiol.* 2008;103:665–75.
42. Gaesser GA, Ward SA, Baum VC, Whipp BJ. Effects of infused epinephrine on slow phase of  $O_2$  uptake kinetics during heavy exercise in humans. *J Appl Physiol.* 1994;77:2413–9.
43. Gerbino A, Ward SA, Whipp BJ. Effects of prior exercise on pulmonary gas exchange kinetics during high-intensity exercise in humans. *J Appl Physiol.* 1996;80:99–107.
44. Gibbs CL, Gibson WR. Energy production of rat soleus muscle. *Am J Physiol.* 1972;223:864–71.
45. Grassi B, Gladden LB, Samaja M, Stary CM, Hogan MC. Faster adjustment of  $O_2$  delivery does not affect  $\dot{V}O_2$  on-kinetics in isolated in situ canine muscle. *J Appl Physiol.* 1998;85:1394–403.
46. Grassi B, Gladden LB, Stary CM, Wagner PD, Hogan MC. Peripheral  $O_2$  diffusion does not affect  $\dot{V}O_2$  on-kinetics in isolated in situ canine muscle. *J Appl Physiol.* 1998;85:1404–12.
47. Grassi B, Hogan MC, Greenhaff PL, et al.  $\dot{V}O_2$  on-kinetics in dog gastrocnemius in situ following activation of pyruvate dehydrogenase by dichloroacetate. *J Physiol.* 2002;538:195–207.
48. Grassi B, Hogan MC, Kelley KM, et al. Role of convective  $O_2$  delivery in determining  $\dot{V}O_2$  on-kinetics in canine muscle contracting at peak  $\dot{V}O_2$ . *J Appl Physiol.* 2000;89:1293–301.
49. Grassi B, Hogan MC, Kelley KM, Howlett RA, Gladden LB. Effects of NOS inhibition by L-NAME on oxygen uptake kinetics in isolated canine muscle in situ. *J Physiol.* 2005;568:1021–33.
50. Grassi B, Rossiter HB, Hogan MC, et al. Faster  $O_2$  uptake kinetics in canine skeletal muscle *in situ* after acute creatine kinase inhibition. *J Physiol.* 2011;589:221–3.
51. Hagberg JM, Mullin JP, Nagle FJ. Oxygen consumption during constant-load exercise. *J Appl Physiol.* 1978;45:381–4.
52. Han YS, Geiger PC, Cody MJ, Macken RL, Sieck GC. ATP consumption rate per cross bridge depends on myosin heavy chain isoform. *J Appl Physiol.* 2003;94:2188–96.
53. Harms CA, Babcock MA, McClaran SR, et al. Respiratory muscle work compromises leg blood flow during maximal exercise. *J Appl Physiol.* 1997;82:1573–83.
54. Haseler LJ, Kindig CA, Richardson RS, Hogan MC. The role of oxygen in determining phosphocreatine onset kinetics in exercising humans. *J Physiol.* 2004;558:985–92.
55. He ZH, Bottinelli R, Pellegrino MA, Ferenczi MA, Reggiani C. ATP consumption and efficiency of human single muscle fibers with different myosin isoform composition. *Biophys J.* 2000;79:945–61.

56. Henneman E, Mendell LM. Functional organization of motoneuron pool and its inputs. In: *Handbook of Physiology. The Nervous System. Motor Control*. Bethesda (MD): American Physiological Society. Sect. 1, Vol. II, Pt. 1, Chap. 11. 1981. p. 423–507.
57. Henson LC, Poole DC, Whipp BJ. Fitness as a determinant of oxygen uptake response to constant-load exercise. *Eur J Appl Physiol*. 1989;59:21–8.
58. Henry FM, DeMoor JC. Lactic and alactic oxygen consumption in moderate exercise of graded intensity. *J Appl Physiol*. 1956;8: 608–14.
59. Hepple RT, Howlett RA, Kindig CA, Sary CM, Hogan MC. The O<sub>2</sub> cost of the tension-time integral in isolated single myocytes during fatigue. *Am J Physiol Regul Integr Comp Physiol*. 2010; 298:R983–8.
60. Hill AV. The time course of the oxygen consumption of stimulated frog's muscle. *J Physiol*. 1940;98:207–27.
61. Hill AV, Lupton H. Muscular exercise, lactic acid, and the supply and utilization of oxygen. *Q J Med*. 1923;16:135–71.
62. Hill DW, Poole DC, Smith JC. The relationship between power and the time to achieve  $\dot{V}O_{2max}$ . *Med Sci Sports Exerc*. 2002; 34(4):709–14.
63. Jones AM, Carter H. The effect of endurance training on parameters of aerobic fitness. *Sports Med*. 2000;29:373–86.
64. Jones AM, Fulford J, Wilkerson DP. Influence of prior exercise on muscle [phosphorylcreatine] and deoxygenation kinetics during high-intensity exercise in men. *Exp Physiol*. 2008;93: 468–78.
65. Jones AM, Koppo K, Wilkerson DP, Wilmshurst S, Campbell IT. Dichloroacetate does not speed phase-II pulmonary  $\dot{V}O_2$  kinetics following the onset of heavy intensity cycle exercise. *Pflugers Arch*. 2004;447:867–74.
66. Jones AM, Poole DC. Physiological demands of endurance exercise. In: Maughan RJ, editor. *Olympic Textbook of Science in Sport*. Oxford (UK): IOC, Blackwell; 2008. p. 43–55.
67. Jones AM, Pringle JSM, Carter H. Influence of muscle fiber type and motor unit recruitment on  $\dot{V}O_2$  kinetics. In: Jones AM, Poole DC, editors. *Oxygen Uptake Kinetics in Sport, Exercise and Medicine*. London (UK): Routledge; 2005. p. 261–93.
68. Jones AM, Vanhatalo A, Burnley M, Morton RH, Poole DC. Critical power: implications for the determination of  $\dot{V}O_{2max}$  and exercise tolerance. *Med Sci Sports Exerc*. 2010;42(10):1876–90.
69. Jones AM, Wilkerson DP, Berger NJ, Fulford J. Influence of endurance training on muscle [PCr] kinetics during high-intensity exercise. *Am J Physiol Regul Integr Comp Physiol*. 2007;293: R392–401.
70. Jones AM, Wilkerson DP, Burnley M, Koppo K. Prior heavy exercise enhances performance during subsequent perimaximal exercise. *Med Sci Sports Exerc*. 2003;35(12):2085–92.
71. Jones AM, Wilkerson DP, DiMenna F, Fulford J, Poole DC. Muscle metabolic responses to exercise above and below the “critical power” assessed using <sup>31</sup>P-MRS. *Am J Physiol Regul Integr Comp Physiol*. 2008;294:R585–93.
72. Jones AM, Wilkerson DP, Fulford J. Muscle [phosphocreatine] dynamics following the onset of exercise in humans: the influence of baseline work-rate. *J Physiol*. 2008;586:889–98.
73. Jones AM, Wilkerson DP, Wilmshurst S, Campbell IT. Influence of L-NAME on pulmonary O<sub>2</sub> uptake kinetics during heavy-intensity cycle exercise. *J Appl Physiol*. 2004;96:1033–8.
74. Koga S, Poole DC, Shiojiri T, et al. Comparison of oxygen uptake kinetics during knee-extension and cycle exercise. *Am J Physiol Regul Integr Comp Physiol*. 2005;288:R212–20.
75. Koga S, Shiojiri T, Kondo N, Barstow TJ. Effect of increased muscle temperature on oxygen uptake kinetics during exercise. *J Appl Physiol*. 1997;83:1333–8.
76. Koga S, Shiojiri T, Shibasaki M, Kondo N, Fukuba Y, Barstow TJ. Kinetics of oxygen uptake during supine and upright heavy exercise. *J Appl Physiol*. 1999;87:253–60.
77. Kolkhorst FW, Rezende RS, Levy SS, Buono MJ. Effects of sodium bicarbonate on  $\dot{V}O_2$  kinetics during heavy exercise. *Med Sci Sports Exerc*. 2004;36(11):1895–9.
78. Koppo K, Jones AM, VandenBossche L, Bouckaert J. Effect of prior exercise on  $\dot{V}O_2$  slow component is not related to muscle temperature. *Med Sci Sports Exerc*. 2002;34(10):1600–4.
79. Krogh A, Lindhard J. The regulation of respiration and circulation during the initial stages of muscular work. *J Physiol*. 1913; 47:112–36.
80. Krstrup P. Muscle oxygen uptake and energy turnover during dynamic exercise in humans. [PhD thesis]. Copenhagen (Denmark): University of Copenhagen. 2004. ISBN: 87-89361-87-3, p. 1–241.
81. Krstrup P, Christensen JF, Randers MB, et al. Muscle adaptations and performance enhancements of soccer training for untrained men. *Eur J Appl Physiol*. 2010;108:1247–58.
82. Krstrup P, Ferguson RA, Kjær M, Bangsbo J. ATP and heat production in human skeletal muscle during dynamic exercise: higher efficiency of anaerobic than aerobic ATP resynthesis. *J Physiol*. 2003;549:255–69.
83. Krstrup P, Jones AM, Wilkerson DP, Calbet JA, Bangsbo J. Muscular and pulmonary O<sub>2</sub> uptake kinetics during moderate- and high-intensity sub-maximal knee-extensor exercise in humans. *J Physiol*. 2009;587:1843–56.
84. Krstrup P, Secher NH, Relu MU, Hellsten Y, Söderlund K, Bangsbo J. Neuromuscular blockade of slow twitch muscle fibres elevates muscle oxygen uptake and energy turnover during sub-maximal exercise in humans. *J Physiol*. 2008;586:6037–48.
85. Krstrup P, Söderlund K, Mohr M, Bangsbo J. The slow component of oxygen uptake during intense, sub-maximal exercise in man is associated with additional fibre recruitment. *Pflugers Arch*. 2004;447:855–66.
86. Krstrup P, Söderlund K, Mohr M, Bangsbo J. Slow-twitch fiber glycogen depletion elevates moderate-exercise fast-twitch fiber activity and O<sub>2</sub> uptake. *Med Sci Sports Exerc*. 2004;36(6): 973–82.
87. Krstrup P, Söderlund K, Relu MU, Ferguson RA, Bangsbo J. Heterogeneous recruitment of quadriceps muscle portions and fibre types during moderate intensity knee-extensor exercise: effect of thigh occlusion. *Scand J Med Sci Sports*. 2009;19:576–84.
88. Lansley KE, Winyard PG, Fulford J, et al. Dietary nitrate supplementation reduces the O<sub>2</sub> cost of walking and running: a placebo-controlled study. *J Appl Physiol*. 2010;110:591–600.
89. Layec G, Bringard A, Le Fur Y, et al. Effects of a prior high-intensity knee-extension exercise on muscle recruitment and energy cost: a combined local and global investigation in humans. *Exp Physiol*. 2009;94:704–19.
90. Linnarsson D. Dynamics of pulmonary gas exchange and heart rate changes at start and end of exercise. *Acta Physiol Scand*. 1974;415:1–68.
91. MacDonald M, Pedersen PK, Hughson RL. Acceleration of  $\dot{V}O_2$  kinetics in heavy submaximal exercise by hyperoxia and prior high-intensity exercise. *J Appl Physiol*. 1997;83:1318–25.
92. Mahler M. Kinetics of oxygen consumption after a single isometric tetanus of frog sartorius muscle at 20°C. *J Gen Physiol*. 1978;71:559–80.
93. Margaria R, Mangili F, Cuttica F, Cerretelli P. The kinetics of oxygen consumption at the onset of muscular exercise in man. *Ergonomics*. 1965;8:49–54.
94. Osborne MA, Schneider DA. Muscle glycogen reduction in man: relationship between surface EMG activity and oxygen uptake kinetics during heavy exercise. *Exp Physiol*. 2006;91:179–89.
95. Paterson DH, Whipp BJ. Asymmetries of oxygen uptake transients at the on- and offset of heavy exercise in humans. *J Physiol*. 1991;443:575–86.
96. Perrey S, Betik A, Candau R, Rouillon JD, Hughson RL. Comparison of oxygen uptake kinetics during concentric and eccentric cycle exercise. *J Appl Physiol*. 2001;91:2135–42.

97. Piiper J, di Prampero PE, Cerretelli P. Oxygen debt and high-energy phosphates in gastrocnemius muscle of the dog. *Am J Physiol.* 1968;215:523–31.
98. Poole DC, Barstow TJ, Gaesser GA, Willis WT, Whipp BJ.  $\dot{V}O_2$  slow component: physiological and functional significance. *Med Sci Sports Exerc.* 1994;26(11):1354–8.
99. Poole DC, Gladden LB, Kurdak S, Hogan MC. L-(+)-Lactate infusion into working dog gastrocnemius: no evidence lactate *per se* mediates  $\dot{V}O_2$  slow component. *J Appl Physiol.* 1994;76:787–92.
100. Poole DC, Jones AM. Towards an understanding of the mechanistic bases of  $\dot{V}O_2$  kinetics. In: Jones AM, Poole DC, editors. *Oxygen Uptake Kinetics in Sport, Exercise and Medicine*. London (UK): Routledge; 2005. p. 294–328.
101. Poole DC, Schaffartzik W, Knight DR, et al. Contribution of excising legs to the slow component of oxygen uptake kinetics in humans. *J Appl Physiol.* 1991;71:1245–60.
102. Poole DC, Ward SA, Gardner GW, Whipp BJ. Metabolic and respiratory profile of the upper limit for prolonged exercise in man. *Ergonomics.* 1988;31:1265–79.
103. Pringle JS, Doust JH, Carter H, et al. Oxygen uptake kinetics during moderate, heavy and severe intensity “submaximal” exercise in humans: the influence of muscle fibre type and capillarisation. *Eur J Appl Physiol.* 2003;89:289–300.
104. Pringle JS, Doust JH, Carter H, Tolfrey K, Jones AM. Effect of pedal rate on primary and slow-component oxygen uptake responses during heavy-cycle exercise. *J Appl Physiol.* 2003;94:1501–7.
105. Rasmussen UF, Krstrup P, Bangsbo J, Rasmussen HN. The effect of high-intensity exhaustive exercise studied in isolated mitochondria from human skeletal muscle. *Pflügers Arch Eur J Physiol.* 2001;443:180–7.
106. Reggiani C, Potma EJ, Bottinelli R, Canepari M, Pellegrino MA, Steinen GJM. Chemo-mechanical energy transduction in relation to myosin isoform composition in skeletal muscle fibres of the rat. *J Physiol.* 1997;502:449–60.
107. Ribeiro JP, Hughes V, Fielding RA, Holden W, Evans W, Knuttgen HG. Metabolic and ventilatory responses to steady state exercise relative to lactate thresholds. *Eur J Appl Physiol.* 1986;55:215–21.
108. Riggs DS. *Control Theory and Physiological Feedback Mechanisms*. New York (NY): Krieger, Huntington; 1976. p. 1–599.
109. Rossiter HB, Ward SA, Howe FA, et al. Effects of dichloroacetate on  $\dot{V}O_2$  and intramuscular  $^{31}\text{P}$  metabolite kinetics during high-intensity exercise in humans. *J Appl Physiol.* 2003;95:1105–15.
110. Rossiter HB, Ward SA, Kowalchuk JM, Howe FA, Griffiths JR, Whipp BJ. Effects of prior exercise on oxygen uptake and phosphocreatine kinetics during high-intensity knee-extension exercise in humans. *J Physiol.* 2001;537:291–303.
111. Rossiter HB, Ward SA, Kowalchuk JM, Howe FA, Griffiths JR, Whipp BJ. Dynamic asymmetry of phosphocreatine concentration and  $O_2$  uptake between the on- and off-transients of moderate- and high-intensity exercise in humans. *J Physiol.* 2002;541:991–1002.
112. Roston WL, Whipp BJ, Davis JA, Cunningham DA, Effros RM, Wasserman K. Oxygen uptake kinetics and lactate concentration during exercise in humans. *Am Rev Respir Dis.* 1987;135:1080–4.
113. Salvadego D, Lazzer S, Busti C, et al. Gas exchange kinetics in obese adolescents. Inferences on exercise tolerance and prescription. *Am J Physiol Regul Integr Comp Physiol.* 2010;299:R1298–305.
114. Saunders MJ, Evans EM, Arngimsson SA, Allison JD, Cureton KJ. Endurance training reduces end-exercise  $\dot{V}O_2$  and muscle use during submaximal cycling. *Med Sci Sports Exerc.* 2003;35(2):257–62.
115. Saunders MJ, Evans EM, Arngimsson SA, Allison JD, Warren GL, Cureton KJ. Muscle activation and the slow component rise in oxygen uptake during cycling. *Med Sci Sports Exerc.* 2000;32(12):2040–5.
116. Scheuermann B, Hoelting BD, Noble ML, Barstow TJ. The slow component of  $O_2$  uptake is not accompanied by changes in muscle EMG during repeated bouts of heavy exercise in humans. *J Physiol.* 2001;531:245–56.
117. Shinohara M, Moritani T. Increase in neuromuscular activity and oxygen uptake during heavy exercise. *Ann Physiol Anthropol.* 1992;11:257–62.
118. Stienen GJ, Kiers JL, Bottinelli R, Reggiani C. Myofibrillar ATPase activity in skinned human skeletal muscle fibres: fibre type and temperature dependence. *J Physiol.* 1996;493:299–307.
119. Stoudemire NM, Wideman L, Pass KA, McGinnes CL, Gaesser GA, Weltman A. The validity of regulating blood lactate concentration during running by ratings of perceived exertion. *Med Sci Sports Exerc.* 1996;28(4):490–95.
120. Szentesi P, Zaremba R, van Mechelen W, Stienen GJ. ATP utilization for calcium uptake and force production in different types of human skeletal muscle fibres. *J Physiol.* 2001;531:393–403.
121. Tonkonogi M, Walsh B, Tiivelt T, Saks V, Sahlin K. Mitochondrial function in human skeletal muscle is not impaired by high intensity exercise. *Pflügers Arch.* 1999;437:562–8.
122. Vanhatalo A, Fulford J, DiMenna FJ, Jones AM. Influence of hyperoxia on muscle metabolic responses and the power-duration relationship during severe-intensity exercise in humans: a  $^{31}\text{P}$  magnetic resonance spectroscopy study. *Exp Physiol.* 2010;95:528–40.
123. Vanhatalo A, Jones AM. Influence of prior sprint exercise on the parameters of the “all-out critical power test” in men. *Exp Physiol.* 2009;94:255–63.
124. Vanhatalo A, Poole DC, DiMenna FJ, Bailey SJ, Jones AM. Muscle fiber recruitment and the slow component of  $O_2$  uptake: constant work rate vs. all-out sprint exercise. *Am J Physiol Regul Integr Comp Physiol.* 2011;300:700–7.
125. Wendt IR, Gibbs CL. Energy production of rat extensor digitorum longus muscle. *Am J Physiol.* 1973;224:1081–6.
126. Whipp BJ, Mahler M. Dynamics of pulmonary gas exchange during exercise. In: West JB, editor. *Pulmonary Gas Exchange*. Vol. II. London (UK): Organism and Environment, Academic Press; 1980. p. 33–96.
127. Whipp BJ, Rossiter HB. The kinetics of oxygen uptake: Physiological inferences from the parameters. In: Jones AM, Poole DC, editors. *Oxygen Uptake Kinetics in Sport, Exercise and Medicine*. London (UK): Routledge; 2005. p. 62–94.
128. Whipp BJ, Wasserman K. Oxygen uptake kinetics for various intensities of constant-load work. *J Appl Physiol.* 1972;33:351–6.
129. Whipp BJ, Wasserman K. Effect of anaerobiosis on the kinetics of  $O_2$  uptake during exercise. *Fed Proc.* 1986;45:2942–7.
130. Wilkerson DP, Berger NJ, Jones AM. Influence of hyperoxia on pulmonary  $O_2$  uptake kinetics following the onset of exercise in humans. *Respir Physiol Neurobiol.* 2006;153:92–106.
131. Wilkerson DP, Kopko K, Barstow TJ, Jones AM. Effect of prior multiple-sprint exercise on pulmonary  $O_2$  uptake kinetics following the onset of perimaximal exercise. *J Appl Physiol.* 2004;97:1227–36.
132. Wilkerson DP, Kopko K, Barstow TJ, Jones AM. Effect of work rate on the functional “gain” of phase II pulmonary  $O_2$  uptake response to exercise. *Respir Physiol Neurobiol.* 2004;142:211–23.
133. Wilkerson DP, Jones AM. Influence of initial metabolic rate on pulmonary  $O_2$  uptake on-kinetics during severe intensity exercise. *Respir Physiol Neurobiol.* 2006;152:204–19.



134. Wilkerson DP, Jones AM. Effects of baseline metabolic rate on pulmonary O<sub>2</sub> uptake on-kinetics during heavy-intensity exercise in humans. *Respir Physiol Neurobiol.* 2007;156:203–11.
135. Willis WT, Jackman MR. Mitochondrial function during heavy exercise. *Med Sci Sports Exerc.* 1994;26(11):1347–54.
136. Woledge RC. Possible effects of fatigue on muscle efficiency. *Acta Physiol Scand.* 1998;162:267–73.
137. Womack CJ, Davis SE, Blumer JL, Barrett E, Weltman AL, Gaesser GA. Slow component of O<sub>2</sub> uptake during heavy exercise: adaptation to endurance training. *J Appl Physiol.* 1995;79:838–45.
138. Zoladz JA, Gladden LB, Hogan MC, Nieckarz Z, Grassi B. Progressive recruitment of muscle fibers is not necessary for the slow component of  $\dot{V}O_2$  kinetics. *J Appl Physiol.* 2008;105:575–80.
139. Zoladz JA, Korzeniewski B, Grassi B. Training-induced acceleration of oxygen uptake kinetics in skeletal muscle: the underlying mechanisms. *J Physiol Pharmacol.* 2006;57:67–84.
140. Zoladz JA, Sargeant AJ, Emmerich J, Stoklosa J, Zychowski A. Changes in acid–base status of marathon runners during an incremental field test. Relationship to mean competitive marathon velocity. *Eur J Appl Physiol.* 1993;67:71–6.