

Dose-Finding Clinical Trials in Stan




Kristian Brock, StanCon Helsinki, 30-Aug-18



What dose?

Not too toxic...but


High enough to be effective



What dose?

Not too toxic...but

High enough to be effective



Is it any good?

Does it do something desirable?


What dose?

Not too toxic...but

High enough to be effective

Pivotal study

The challenger vs the standard of care



Is it any good?

Does it do something desirable?


What dose?

Not too toxic...but

High enough to be effective

Pivotal study

The challenger vs the standard of care



Is it any good?

Does it do something desirable?


Post-marketing

We'll use it...

But let's keep an eye on it


**Dose-finding by
toxicity**

Dose Finding and the 3+3 Method




Surely folk use statistical models these days?

- Chiuzan *et al.* (2017) conducted systematic review of dose-finding methods
- Cancer trials published between 2008 and 2014




“adoption of [model-based] designs continues to remain low”


Continual Reassessment Method (CRM)



Continual Reassessment Method (CRM)




Continual Reassessment Method (CRM)




Levy et al. - ssHHT in leukaemia

Prob(Toxicity) after all 18 patients in Levy, et al.



Levy, et al. (2006) A Phase I Dose-Finding and Pharmacokinetic Study of Subcutaneous Semisynthetic Homoharringtonine (ssHHT) in Patients with Advanced Acute Myeloid Leukaemia. BJC. <https://doi.org/10.1038/sj.bjc.6603265>.



Levy *et al.* - ssHHT in leukaemia




Levy, et al.(2006) A Phase I Dose-Finding and Pharmacokinetic Study of Subcutaneous Semisynthetic Homoharringtonine (ssHHT) in Patients with Advanced Acute Myeloid Leukaemia. BJC. <https://doi.org/10.1038/sj.bjc.6603265>.

**Dose-finding by
toxicity & efficacy**

Dose-toxicity & efficacy in modern therapies



EffTox - dose-finding based on efficacy & toxicity trade-offs



Posterior beliefs on attractiveness

Prob(“dose in column is more attractive than dose in row”)

Dose	1	2	3	4	5
1		0.92	0.86	0.82	0.8
2			0.75	0.68	0.64
3				0.58	0.54
4					0.52
5					

trialr - clinical trial designs in R & Stan

```
1
2 # Install|
3 devtools::install_github('brockk/trialr')
4 # Or
5 install.packages('trialr')
6
7 # CRM
8 levy <- '1NNN 3NNT 4NNT 4NNN 4NTN 4TNT'
9 skeleton <- c(0.05, 0.10, 0.15, 0.33, 0.5)
10 target <- 0.33
11 levy_mod <- stan_crm(levy, skeleton = skeleton, target = target,
12                     model = 'logistic_gamma', a0 = 4,
13                     beta_shape = 1, beta_inverse_scale = 1,
14                     seed = 123, control = list(adapt_delta = 0.95))
15
16 # EffTox
17 outcomes <- '1NEN 2NBE'
18 mod2 <- stan_efftox_demo(outcomes, seed = 123)
19 # That is short-hand for
20 mod2 <- stan_efftox(outcomes,
21                     real_doses = c(1.0, 2.0, 4.0, 6.6, 10.0),
22                     efficacy_hurdle = 0.5, toxicity_hurdle = 0.3,
23                     p_e = 0.1, p_t = 0.1,
24                     eff0 = 0.5, tox1 = 0.65,
25                     eff_star = 0.7, tox_star = 0.25,
26                     alpha_mean = -7.9593, alpha_sd = 3.5487,
27                     beta_mean = 1.5482, beta_sd = 3.5018,
28                     gamma_mean = 0.7367, gamma_sd = 2.5423,
29                     zeta_mean = 3.4181, zeta_sd = 2.4406,
30                     eta_mean = 0, eta_sd = 0.2,
31                     psi_mean = 0, psi_sd = 1,
32                     seed = 123)
33
```

Future work:

- Add more trial designs :-)
- Plumb it to work with tidybayes
- More visualisation via bayesplot
- Scrutinise fit by shinystan, etc
- Automated documents via Flexdashboards?

The End

Kristian Brock

Senior Biostatistician

Cancer Research UK Clinical Trials Unit

University of Birmingham

k.brock@bham.ac.uk or

kristian.brock@gmail.com



Institute of
Translational
Medicine

Birmingham



UNIVERSITY OF
BIRMINGHAM