

Alternative splicing regulates mouse embryonic stem cell pluripotency and differentiation

Nathan Salomonis^{a,b,1}, Christopher R. Schlieve^a, Laura Pereira^c, Christine Wahlquist^d, Alexandre Colas^d, Alexander C. Zambon^e, Karen Vranizan^{f,2}, Matthew J. Spindler^{a,b}, Alexander R. Pico^a, Melissa S. Cline^g, Tyson A. Clark^g, Alan Williams^g, John E. Blume^g, Eva Samal^a, Mark Mercola^d, Bradley J. Merrill^e, and Bruce R. Conklin^{a,b,h,1}

^aGladstone Institute of Cardiovascular Disease, San Francisco, CA 94158; ^bPharmaceutical Sciences and Pharmacogenomics Graduate Program, University of California, San Francisco, CA 94143; ^cDepartment of Biochemistry and Molecular Genetics, University of Illinois, Chicago, IL 60607; ^dSanford-Burnham Institute for Medical Research, La Jolla, CA 92037; ^eDepartment of Pharmacology, University of California, La Jolla, CA 92093; ^fFunctional Genomics Laboratory, University of California, Berkeley, CA 94720; ^gAffymetrix, Inc., Santa Clara, CA 95051; and ^hDepartments of Medicine and Cellular and Molecular Pharmacology, University of California, San Francisco, CA 94143

Edited by Elaine Fuchs, The Rockefeller University, New York, NY, and approved April 27, 2010 (received for review October 27, 2009)

Two major goals of regenerative medicine are to reproducibly transform adult somatic cells into a pluripotent state and to control their differentiation into specific cell fates. Progress toward these goals would be greatly helped by obtaining a complete picture of the RNA isoforms produced by these cells due to alternative splicing (AS) and alternative promoter selection (APS). To investigate the roles of AS and APS, reciprocal exon–exon junctions were interrogated on a genome-wide scale in differentiating mouse embryonic stem (ES) cells with a prototype Affymetrix microarray. Using a recently released open-source software package named AltAnalyze, we identified 144 genes for 170 putative isoform variants, the majority (67%) of which were predicted to alter protein sequence and domain composition. Verified alternative exons were largely associated with pathways of Wnt signaling and cell-cycle control, and most were conserved between mouse and human. To examine the functional impact of AS, we characterized isoforms for two genes. As predicted by AltAnalyze, we found that alternative isoforms of the gene *Serca2* were targeted by distinct microRNAs (miRNA-200b, miRNA-214), suggesting a critical role for AS in cardiac development. Analysis of the Wnt transcription factor *Tcf3*, using selective knockdown of an ES cell-enriched and characterized isoform, revealed several distinct targets for transcriptional repression (*Stmn2*, *Ccnd2*, *Atf3*, *Klf4*, *Nodal*, and *Jun*) as well as distinct differentiation outcomes in ES cells. The findings herein illustrate a critical role for AS in the specification of ES cells with differentiation, and highlight the utility of global functional analyses of AS.

AltAnalyze | microRNA | splice isoforms | *Atp2a2* | *Tcf7l1*

Embryonic stem (ES) cells are a vital tool for studying the regulation of early embryonic propagation and cell-fate decisions. Research in this area has led to the development of new technologies for adult somatic cell reprogramming and insights into the steps required for lineage commitment (1, 2). Several factors critical for the self-renewal of pluripotent cells have been identified in both conventional biochemical screens and genome-wide expression analyses. These include the transcription factors Oct4, Sox2, and Nanog, which interact with a common set of promoters to promote self-renewal and pluripotency (3). Recently, Tcf3, a β -catenin-responsive transcription factor, was implicated in this core transcriptional network as a direct transcriptional repressor of Oct4 and Nanog (4, 5), and is itself a target of these factors (6, 7).

In higher eukaryotes, alternative splicing (AS) and alternative promoter selection (APS) contribute to proteomic diversity by increasing the number of distinct mRNAs from a single gene locus. In different tissues and cellular states, transcript variation can alter protein interaction networks by removing or inserting protein domains, changing subcellular localization, or regulating gene expression (8). AS can also remove binding sites for translational repression by microRNAs (miRNAs) (9). AS and

APS appear to increase transcript diversity in ES cells and thus are likely to affect differentiation to distinct tissue lineages (10–13). A better understanding of how AS regulates protein diversity and translational repression during ES cell differentiation may provide critical insights into this process.

Results

AS and APS Are Prominent Features of Mouse ES Cell Differentiation. ES cells and embryoid bodies (EBs) were profiled with a prototype Affymetrix exon–exon junction microarray, which interrogates the mRNAs of ~7,500 genes and more than 40,000 putative exon–exon junctions. For this study, we developed additional analytical methods for a free open-source program named AltAnalyze (<http://www.altanalyze.org/>) (13) (*SI Materials and Methods*). To identify alternative exons (AEs) that might indicate AS or APS, we added a previously described linear regression-based method (14) to AltAnalyze to compare the expression of reciprocal exon–exon junctions (pairs of exon–exon junctions measuring the inclusion of one or more exons). In addition to scoring AEs, this software determines the likelihood of an AE score occurring by chance, assigns protein associations to regulated exon–exon junctions, and identifies functional sequence elements (domains, motifs, and miRNA binding sites) differing between aligning alternate mRNAs and their corresponding proteins (Fig. 1A) (13).

Using AltAnalyze to analyze ES cell differentiation, we identified 170 unique AEs corresponding to 144 genes of 4,269 expressed genes (*Dataset S1*). Pathway analysis of these genes with the program GO-Elite (15), from within AltAnalyze, showed enrichment in Wnt and TGF- β receptor signaling pathways, actin cytoskeleton, lipid transport, muscle contraction, mRNA metabolism, and embryonic development, among others (*Table S1*).

Of the 170 AEs, 108 had evidence of AS, and 20 had evidence of APS; the remainder did not associate with a known AS or APS event. The majority of these annotated AS events aligned to pre-

Author contributions: N.S., C.R.S., L.P., C.W., A.C., A.C.Z., K.V., M.J.S., M.S.C., T.A.C., A.W., J.E.B., E.S., M.M., B.J.M., and B.R.C. designed research; N.S., C.R.S., L.P., C.W., A.C., A.C.Z., M.J.S., and E.S. performed research; N.S., C.W., A.C., M.J.S., A.R.P., M.S.C., T.A.C., A.W., J.E.B., and M.M. contributed new reagents/analytic tools; N.S., C.R.S., L.P., C.W., A.C., K.V., and E.S. analyzed data; and N.S. wrote the paper.

The authors declare no conflict of interest.

This article is a PNAS Direct Submission.


Freely available online through the PNAS open access option.

Data deposition: The data reported in this paper have been deposited in the Gene Expression Omnibus (GEO) database, www.ncbi.nlm.nih.gov/geo/ (accession no. GSE17021).

¹To whom correspondence should be addressed. E-mail: nsalomonis@gladstone.ucsf.edu or bconklin@gladstone.ucsf.edu.

²Deceased June 8, 2009.

This article contains supporting information online at www.pnas.org/lookup/suppl/doi:10.1073/pnas.0912260107/-DCSupplemental.



short form of Tcf3 [Tcf3(s)], our analysis highlights a longer isoform of Tcf3 [Tcf3(l)], enriched in ES cells and down-regulated upon differentiation.

The two Tcf3 isoforms detected by our exon–exon junction microarray analysis differ in the inclusion of a 42-bp cassette exon, which encodes an additional 14 amino acids [i.e., Tcf3(l)] that overlap with the Groucho binding domain (41, 42). The Groucho binding domain is necessary for Tcf3 to repress a Nanog reporter (4). As shown by RT-PCR and qPCR, Tcf3(l) is up-regulated 2-fold in ES cells vs. EBs, and Tcf3(s) is expressed at roughly equivalent levels (Fig. S4A and B). Proteins for both isoforms were detected in ES cells, as were cDNAs encoding each isoform in a Tcf3-null ($-/-$) ES cell line (4) (Fig. 6A).

To assess their ability to regulate the transcription of known targets, the Tcf3 isoforms were transiently expressed in Tcf3 $^{-/-}$ ES cells and assayed for expression of Tcf/Lef- β -catenin transcriptional (TOPFlash) (43) or Nanog promoter-driven luciferase reporter plasmids (4). The isoforms equally inhibited the TOPFlash and Nanog reporters in a concentration-dependent manner (Fig. 6B and C). Thus, Tcf3(l) retains transcriptional repressive activity for Tcf/Lef targets and the Nanog promoter-driven luciferase.

Tcf3(l) Knockdown Preferentially Increases Known and Novel Target Gene Expression. Given that Tcf3 isoforms show similar activity on episomal promoters, we used RNAi to examine the effects of these isoforms on endogenous promoters. Stable knockdown of Tcf3(s), Tcf3(l), and all Tcf3 isoforms [Tcf3(a)] was achieved in ES cells with the pSicoR-Ef1 α -mCh-puro lentiviral construct with isoforms-specific short hairpin RNAs (shRNAs; Fig. 6D). This strategy yielded up to 90% knockdown (KD) of the targeted isoforms, with minimal or no reduction in the nontargeted isoform in ES cell and EB RNA (Fig. S4C). When analyzed by Western blot, no Tcf3 protein was detected in Tcf3(a) KD ES cells (Fig. S4D).

KD of any Tcf3 isoform increased Nanog expression, with maximal up-regulation by Tcf3(a) KD (4-fold), followed by Tcf3(l) (3.5-fold), and Tcf3(s) (1.8-fold) KD (Fig. 6E). Interestingly, whereas Oct4 was up-regulated with Tcf3(l) KD (2-fold), Oct4 expression levels were unchanged with either Tcf3(s) or Tcf3(a) KD relative to wild-type ES cells (t test $P < 0.05$; Fig. 6F). Although Tcf3(l) KD produced a greater up-regulation of Oct4 and Nanog than Tcf3(s) KD, this effect might be due to the higher expression of Tcf3(l) in undifferentiated ES cells.

To expand the search for differentially responsive Tcf3 transcriptional targets, we examined the expression of a panel of 17 known and 17 novel candidate TCF target genes. Novel Tcf3 target genes were identified as being affected by Tcf3 knockout, KD, induction, and promoter occupancy microarray studies (6, 40, 44). Of the 34 genes examined, 20 genes were differentially expressed with KD of at least one of Tcf3 isoform ($P < 0.05$; Fig. S5). Among these differentially expressed genes, six (Atf3, Ccnd2, Jun, Klf4, Nodal, and Stmn2) exhibited a 2-fold or greater difference in expression between Tcf3(s) and Tcf3(l) KD (Fig. 6G). These patterns of expression persisted throughout ES cell differentiation (Fig. S64). These factors are implicated in both the regulation cell-cycle progression and a broad range of developmental pathways (Fig. 2) (30, 32, 33, 45). Tcf3 promoter occupancy was previously observed for Atf3, Ccnd2, Jun, Nodal, and Stmn2, as was opposing regulation by Tcf3 knockout and induction for Atf3, Jun, Klf4, and Stmn2 (6, 40, 44). These data suggest that the two Tcf3 isoforms regulate both an overlapping set as well as distinct sets of target genes.

Delayed Differentiation of Tcf3 Knockdown Lines. Isoform-specific Tcf3 KD lines displayed unique cell-colony morphologies in the presence and absence of LIF (Fig. 7 and Fig. S7). With LIF, wild-type E14 ES cells typically distribute into a monolayer when grown on gelatinized culture plates (without mouse feeder cells). Tcf3(a),




Fig. 6. Selective Tcf3 isoform expression and transcriptional activities in ES cells. (A) Tcf3 $^{-/-}$ ES cells were transfected with full-length cDNAs for both isoforms expressed under the control of a cytomegalovirus promoter and compared with wild-type ES cells. Expression of Tcf3(s) and Tcf3(l) isoforms resulted in a shift in the detection of Tcf3 protein on a polyacrylamide gel with a Tcf3 antibody common to both isoforms. To assess activity of these isoforms, Tcf3 $^{-/-}$ ES cells were transfected with (B) a Tcf/Lef transcriptional reporter construct (TOPFlash) and a stable form of β -catenin (β -cat) or (C) Nanog promoter reporter, driving luciferase expression, with or without increasing amounts of transfected cDNA for the Tcf3 isoforms. (D) To achieve isoform-specific KD, Tcf3 regions unique to or in common with each isoform were targeted using shRNAs directed against either exon junctions [E3–E4 for all or E4–E5 for Tcf3(s)] or exons [exon E4a for Tcf3(l)]. Quantitative expression levels for (E) Oct4, (F) Nanog, and (G) putative Tcf3 transcriptional targets are reported for undifferentiated wild-type (wt) and isoform-specific Tcf3 KD lines cultured with LIF. Fold induction is relative to wild-type ES cells. Values are mean \pm SEM of biological triplicates. An asterisk indicates t test $P < 0.05$.

Tcf3(s), and Tcf3(l) KD ES cells resulted in clustered, rounded colonies when maintained in LIF. When wild-type ES cells were removed from LIF, they displayed a differentiated morphology (cobblestone-like) by day 3 and could not be passaged further, whereas all of the Tcf3-KD lines could be further passaged to day 6 and did not have a predominantly differentiated morphology. In




Fig. 7. Altered morphology of ES cells and differentiated EBs with Tcf3 isoform knockdown. Cell morphology, clustering, and differentiation were observed for wild-type, Tcf3(s), Tcf3(l), and Tcf3(a) KD ES cells before and after removal of LIF (–LIF). See Fig. S7 for additional time points.

addition to confirming that complete loss of Tcf3 allows for LIF-independent ES cell propagation (5, 40), these data suggest that loss of either Tcf3 isoform enhances self-renewal.

Tcf3 Knockdowns Inhibit Distinct Differentiation Pathways. To assess differentiation outcomes with loss of individual Tcf3 isoforms, wild-type and Tcf3 KD lines were derived for EBs at 1, 3, 6, and 9 d of differentiation. qPCR analysis of the time-course data revealed that Tcf3(s) KD had the most profound overall effect on differentiation marker expression, blunting the expression of most lineage (ectoderm, endoderm, mesoderm) and tissue-specific markers examined (Fig. 8 and Fig. S6B). This effect was most similar to Tcf3(a) KD, except for the ectodermal markers Fgf5 and Nestin, the endodermal markers FoxA2 and Sox17, and the osteoblast marker Sp7, which were expressed at similar or greater levels than in wild-type ES cells. Tcf3(l) KD largely did not affect expression of lineage markers, but blunted expression of the examined cardiac and neural markers. These data suggest that Tcf3 isoforms have distinct roles in certain, but not all, lineage pathways.

Discussion

Our study uncovered over 100 putative AS events predicted to differ substantially in ES cells and EBs. These events appear to be unique to ES cell differentiation and occur in genes essential to the maintenance of pluripotency, lineage specification, and cell-cycle progression. In many cases, the predicted AS events appear to affect the putative domain structure of the resulting proteins, suggesting such proteins have a functional role in differentiating ES cells. Analysis of Serca2 confirmed removal of targeted miRNA binding sites upon ES cell differentiation in both a candidate approach and an unbiased screen. One targeted miRNA binding site, miR-200b, is restricted to the cardiac progenitor lineage, where the nontargeted isoform, Serca2a, is also most highly expressed in this cell type (38, 39). This observation has broad implications for the ability of AS to regulate protein expression, without the need to regulate target gene or miRNA transcription. In the case of Tcf3, although selective expression of Tcf3 isoforms similarly limited self-renewal of ES cells, a handful of transcriptional targets were regulated by one but not both isoforms. Interestingly, these target genes are associated with a broad range of self-renewal and differentiation pathways, raising the interesting hypothesis that isoform-specific regulation of these targets affects distinct lineage commitment decisions. Further analysis of Tcf3 isoforms will be critical to determine how these isoforms interact differently with target promoters and binding partners, and in which cells they are restricted to during lineage commitment.

Our analysis shows multiple strategies for identifying and validating functionally relevant splicing changes in vitro. Although only a minority of AS events may impact self-renewal or pluripotency, analysis with AltAnalyze should highlight changes




Fig. 8. Distinct patterns of lineage marker expression for all Tcf3 knockdown lines. qPCR was performed on RNA extracted from multiple days of EB differentiation (days 1–9) for wild-type and Tcf3 isoform KD lines. Relative gene expression changes for all cell lines and time points are compared with gene expression of wild-type ESCs at day 9 of differentiation, where gene expression was typically highest. The lineages for which each marker is associated are listed in gray under the gene name for each. Values are mean \pm SEM of technical triplicates of pooled EB plates ($n = 96$ EBs or greater).

in isoform expression most likely to affect protein composition and expression. As whole-genome exon, junction, and RNA sequencing data become the standard method for expression analysis, we anticipate that these methods will become increasingly important to score for AS and predict the potential impact on protein composition and expression.

Materials and Methods

Tissue Isolation and Sample Preparation. Mouse E14 ES cells were grown in monolayer on gelatin-coated culture plates, maintained in medium supplemented with 10% FBS, sodium pyruvate, nonessential amino acids, β -mercaptoethanol, and LIF, and passaged with trypsin. EBs were derived by the hanging-drop method as described (39) in 20% FBS to enrich for the mesoderm lineage. For the microarray analyses, EBs were allowed to differentiate for 14 d. Isolated mouse (FVB/N) tissues consisted of age-matched adult cardiac atria and ventricle, myometrium from adult virgin, 14.5 d (quiescent), 18.5 d (term) gestation, and 6 h postpartum (46), and mouse adult cardiac ventricles from a model of dilated cardiomyopathy (47) and single-transgenic α -myosin heavy chain promoter–tetracycline transactivator controls. Affymetrix CEL files from a published tissue compendium (14) and Nova2 knockout (18) dataset were also included. Total RNA was isolated from cell cultures or snap-frozen tissues, in biological triplicates or greater, using TRIzol extraction and purified with the Qiagen RNA Purification Kit. The AltMouse A microarray design, sample preparation, and hybridization protocol have been described (14). Microarray data were deposited at the NCBI Gene Expression Omnibus (GEO) database (accession no. GSE17021). Additional details on alternative exon analyses are provided in *SI Materials and Methods*.

miRNA Targeting and Expression of Serca2 Isoforms. To assess miRNA repression of SERCA2b, we prepared a DNA construct that transcribes SERCA2b and GFP in a single transcript. This construct was derived by substituting the CMV promoter of the Clontech pDsRed-N1 vector with the PGK promoter driving the expression of eGFP and replacing the DsRed sequence with human SERCA2b full-length cDNA (BC035588). The SERCA2b construct was transfected into HEK-293 cells (300 ng per well) in the presence of siRNAs for phospholamban (negative control), SERCA2 (positive control), or mimics for each miRNA of the Pre-miR miRNA Precursor Library–Human V2 (Ambion) at a concentration of 10 nM. Transfections were performed in duplicates, and fluorescence was quantified using an automated fluorescent microscope (InCell Analyzer 1000; Amersham Biosciences) and a custom algorithm (48). miRNA binding sites identified in the screen were bioinformatically predicted using annotations from AltAnalyze, sequence alignment, and RNA22 (http://cbcsrv.watson.ibm.com/rna22_download_content.html). To assess miRNA binding-site targeting of miRNA-200b, an oligonucleotide duplex for the predicted Serca2b 3'UTR miRNA-200b binding site was obtained from Integrated DNA Technologies and cloned into the pMIR-reporter vector

(Ambion) immediately downstream of the luciferase protein. The miRNA-200b pMIR-reporter was transfected into HL-1 cells (0.4 μ g) with Lipofectamine 2000 (Invitrogen) with or without miR-200b, -429, or -1 mimics (50 pmoles) from Dharmacon, and luciferase activity was measured after a 24-h incubation. Repression of the pMIR-reporters was measured by normalizing luciferase to Renilla activity.

Isoform-Specific Expression/shRNA in Mouse ES Cells. Stable isoform-specific KD of Tcf3 alternatively spliced isoforms were obtained in E14 ES cells with sequence-specific shRNAs, delivered by lentiviral infection. Three 19-mer shRNAs were designed to target Tcf3(I) (GGATGGTGCCTCCACATT), Tcf3(s) (CCAGCACTTGTCCAAACA), and a constitutive region of Tcf3 (GCACCTACACAGATGAA) by selecting overlapping predictions from the program PSICOLIGOMAKER 1.5 (<http://web.mit.edu/jacks-lab/protocols/pSico.html>) and the Broad Institute mouse hairpin library (<http://www.broadinstitute.org/rnai/trc/lib>). The pSicoR-Ef1 α -mCh-Puro lentiviral construct was created by replacing the CMV promoter of pSicoR-mCherry with Ef1 α and adding a T2A-puromycin-resistance gene cassette following mCherry (49). Isoform-specific shRNAs were ligated into the pSicoR-Ef1 α -mCh-puro construct and cotransfected with the viral packaging plasmids pMDLgprRE, pRSV_Rev, and pVSV-G (Clontech) into HEK293 cells with FuGENE6

(Roche) as described (49). Harvested supernatant from viral-producing HEK293 cells was filtered through a 0.45- μ m filter, and 100 μ L incubated with 200,000 ES cells on rotator for 3 h. Cells were plated onto gelatinized tissue-culture plates, grown under feeder-free conditions in the presence of LIF, and selected for puromycin-resistant colonies for at least 5 d. Clonal populations of mCherry-expressing ES cells were screened with isoform-specific qPCR primers to select for clones with optimal isoform-specific KD. cDNAs for both the two Tcf3 isoforms were also expressed in Tcf3^{-/-} ES cells on 129/Sv background (G51) by electroporation or transfection of a linearized pCDNA3-Tcf3(s) (pBM58) (4) or pCDNA3-Tcf3(I) constructs. The pCDNA3-Tcf3(I) construct was obtained by removal of a 670-bp fragment by Kpn1-Pml1 (NEB) digestion of pBM58 and insertion of the corresponding 712-bp ES cell RT-PCR fragment from Tcf3(I).

ACKNOWLEDGMENTS. We thank Gary Howard and Stephen Ordway, Drs. Chuck Sugnet, Jernej Ule, and Kathryn Ivey for data analysis advice, and Jo Dee Fish and Chris Barker from the Gladstone Histology and Genomic Cores. This work was supported by National Institutes of Health Grants GM080223, HG003053, HL66621 (to B.R.C. and N.S.), HL057181, HL092851 (to E.S.), R01CA128571 (to B.J.M.), R37HL059502, R33HL088266 and California Institute for Regenerative Medicine RC1-000132-1 (to M.M.).

- Yamanaka S (2008) Pluripotency and nuclear reprogramming. *Philos Trans R Soc Lond B Biol Sci* 363:2079–2087.
- Vallier L, Pedersen RA (2005) Human embryonic stem cells: An in vitro model to study mechanisms controlling pluripotency in early mammalian development. *Stem Cell Rev* 1:119–130.
- Boyer LA, et al. (2005) Core transcriptional regulatory circuitry in human embryonic stem cells. *Cell* 122:947–956.
- Pereira L, Yi F, Merrill BJ (2006) Repression of Nanog gene transcription by Tcf3 limits embryonic stem cell self-renewal. *Mol Cell Biol* 26:7479–7491.
- Tam WL, et al. (2008) Tcf3 regulates embryonic stem cell pluripotency and self-renewal by the transcriptional control of multiple lineage pathways. *Stem Cells* 26:2019–2031.
- Cole MF, Johnstone SE, Newman JJ, Kagey MH, Young RA (2008) Tcf3 is an integral component of the core regulatory circuitry of embryonic stem cells. *Genes Dev* 22:746–755.
- Loh YH, et al. (2006) The Oct4 and Nanog transcription network regulates pluripotency in mouse embryonic stem cells. *Nat Genet* 38:431–440.
- Cooper TA (2005) Alternative splicing regulation impacts heart development. *Cell* 120:1–2.
- Duursma AM, Kedde M, Schrier M, le Sage C, Agami R (2008) miR-148 targets human DNMT3b protein coding region. *RNA* 14:872–877.
- Pritsker M, Doniger TT, Kramer LC, Westcot SE, Lemischka IR (2005) Diversification of stem cell molecular repertoire by alternative splicing. *Proc Natl Acad Sci USA* 102:14290–14295.
- Yeo GW, et al. (2007) Alternative splicing events identified in human embryonic stem cells and neural progenitors. *PLOS Comput Biol* 3:1951–1967.
- Kunars G, Wong KY, Stanton LW, Lipovich L (2008) Detailed characterization of the mouse embryonic stem cell transcriptome reveals novel genes and intergenic splicing associated with pluripotency. *BMC Genomics* 9:155.
- Salomonis N, et al. (2009) Alternative splicing in the differentiation of human embryonic stem cells into cardiac precursors. *PLOS Comput Biol* 5:e1000553.
- Sugnet CW, et al. (2006) Unusual intron conservation near tissue-regulated exons found by splicing microarrays. *PLOS Comput Biol* 2:e4.
- Krishnan K, Salomonis N, Guo S (2008) Identification of Spt5 target genes in zebrafish development reveals its dual activity in vivo. *PLoS ONE* 3:e3621.
- Chu PJ, Larsen JK, Chen CC, Best PM (2004) Distribution and relative expression levels of calcium channel beta subunits within the chambers of the rat heart. *J Mol Cell Cardiol* 36:423–434.
- Sato N, et al. (2003) A novel variant of cardiac myosin-binding protein-C that is unable to assemble into sarcomeres is expressed in the aged mouse atrium. *Mol Biol Cell* 14:3180–3191.
- Ule J, et al. (2005) Nova regulates brain-specific splicing to shape the synapse. *Nat Genet* 37:844–852.
- Tyson-Capper AJ (2007) Alternative splicing: An important mechanism for myometrial gene regulation that can be manipulated to target specific genes associated with preterm labour. *BMC Pregnancy Childbirth* 7 (Suppl 1):S13.
- Curley M, Morrison JJ, Smith TJ (2004) Analysis of Maxi-K alpha subunit splice variants in human myometrium. *Reprod Biol Endocrinol* 2:67.
- Dabertrand F, Fritz N, Mironneau J, Macrez N, Morel JL (2007) Role of RYR3 splice variants in calcium signaling in mouse nonpregnant and pregnant myometrium. *Am J Physiol Cell Physiol* 293:C848–C854.
- Biesiadecki BJ, Elder BD, Yu ZB, Jin JP (2002) Cardiac troponin T variants produced by aberrant splicing of multiple exons in animals with high instances of dilated cardiomyopathy. *J Biol Chem* 277:50275–50285.
- Spring CM, et al. (2005) The catenin p120ctn inhibits Kaiso-mediated transcriptional repression of the beta-catenin/TCF target gene matrilysin. *Exp Cell Res* 305:253–265.
- Imbalzano AN, Jones SN (2005) Snf5 tumor suppressor couples chromatin remodeling, checkpoint control, and chromosomal stability. *Cancer Cell* 7:294–295.
- Bachmann M, Hennemann H, Xing PX, Hoffmann I, Mörry T (2004) The oncogenic serine/threonine kinase Pim-1 phosphorylates and inhibits the activity of Cdc25C-associated kinase 1 (C-TAK1): A novel role for Pim-1 at the G2/M cell cycle checkpoint. *J Biol Chem* 279:48319–48328.
- Dugast-Darzacq C, Grange T, Schreiber-Agus NB (2007) Differential effects of Mxi1-SRalpha and Mxi1-SRbeta in Myc antagonism. *FEBS J* 274:4643–4653.
- Huang SC, et al. (2005) Mitotic regulation of protein 4.1R involves phosphorylation by cdc2 kinase. *Mol Biol Cell* 16:117–127.
- Ishtiani T, et al. (1999) The TAK1-NLK-MAPK-related pathway antagonizes signalling between beta-catenin and transcription factor TCF. *Nature* 399:798–802.
- Katoh Y, Katoh M (2007) Comparative genomics on PROM1 gene encoding stem cell marker CD133. *Int J Mol Med* 19:967–970.
- Wang H, Mo P, Ren S, Yan C (2010) Activating transcription factor 3 activates p53 by preventing E6-associated protein from binding to E6. *J Biol Chem* 285:13201–13210.
- Rocha S, Martin AM, Meek DW, Perkins ND (2003) p53 represses cyclin D1 transcription through down regulation of Bcl-3 and inducing increased association of the p52 NF-kappaB subunit with histone deacetylase 1. *Mol Cell Biol* 23:4713–4727.
- Wisdom R, Johnson RS, Moore C (1999) c-Jun regulates cell cycle progression and apoptosis by distinct mechanisms. *EMBO J* 18:188–197.
- Vallier L, Alexander M, Pedersen RA (2005) Activin/Nodal and FGF pathways cooperate to maintain pluripotency of human embryonic stem cells. *J Cell Sci* 118:4495–4509.
- Dugast-Darzacq C, Pirity M, Blanck JK, Scherl A, Schreiber-Agus N (2004) Mxi1-SRalpha: A novel Mxi1 isoform with enhanced transcriptional repression potential. *Oncogene* 23:8887–8899.
- Gee SL, et al. (2000) Alternative splicing of protein 4.1R exon 16: Ordered excision of flanking introns ensures proper splice site choice. *Blood* 95:692–699.
- Greene AL, et al. (2000) Overexpression of SERCA2b in the heart leads to an increase in sarcoplasmic reticulum calcium transport function and increased cardiac contractility. *J Biol Chem* 275:24722–24727.
- Misquitta CM, Mwanjewe J, Nie L, Grover AK (2002) Sarcoplasmic reticulum Ca(2+) pump mRNA stability in cardiac and smooth muscle: Role of the 3'-untranslated region. *Am J Physiol Cell Physiol* 283:C560–C568.
- Domian IJ, et al. (2009) Generation of functional ventricular heart muscle from mouse ventricular progenitor cells. *Science* 326:426–429.
- Ivey KN, et al. (2008) MicroRNA regulation of cell lineages in mouse and human embryonic stem cells. *Cell Stem Cell* 2:219–229.
- Yi F, Pereira L, Merrill BJ (2008) Tcf3 functions as a steady state limiter of transcriptional programs of mouse embryonic stem cell self renewal. *Stem Cells* 26:1951–1960.
- Cavallo RA, et al. (1998) Drosophila Tcf and Groucho interact to repress Wingless signalling activity. *Nature* 395:604–608.
- Roose J, et al. (1998) The Xenopus Wnt effector XTcf-3 interacts with Groucho-related transcriptional repressors. *Nature* 395:608–612.
- Veeman MT, Slusarski DC, Kaykas A, Louie SH, Moon RT (2003) Zebrafish prickle, a modulator of noncanonical Wnt/Fz signaling, regulates gastrulation movements. *Curr Biol* 13:680–685.
- Nishiyama A, et al. (2009) Uncovering early response of gene regulatory networks in ESCs by systematic induction of transcription factors. *Cell Stem Cell* 5:420–433.
- Chellini C, et al. (2008) Stathmin-like 2, a developmentally-associated neuronal marker, is expressed and modulated during osteogenesis of human mesenchymal stem cells. *Biochem Biophys Res Commun* 374:64–68.
- Salomonis N, et al. (2005) Identifying genetic networks underlying myometrial transition to labor. *Genome Biol* 6:R12.
- Redfern CH, et al. (2000) Conditional expression of a Gi-coupled receptor causes ventricular conduction delay and a lethal cardiomyopathy. *Proc Natl Acad Sci USA* 97:4826–4831.
- Bushway PJ, Mercola M (2006) High-throughput screening for modulators of stem cell differentiation. *Methods Enzymol* 414:300–316.
- Grskovic M, Chaivorapol C, Gaspar-Maia A, Li H, Ramalho-Santos M (2007) Systematic identification of cis-regulatory sequences active in mouse and human embryonic stem cells. *PLoS Genet* 3:e145.