

Ejercicios Certamen 2 Mecánica Clásica Avanzada

Simon Fonseca

June 9, 2026

1 Problema 1

A point-like nonrelativistic particle of mass m scatters in the time-dependent spherically symmetric harmonic potential

$$U(r, t) = \frac{k(t)}{2} r^2, \quad k(t) = \frac{\kappa}{t^2}, \quad \kappa > 0 \quad (1.1)$$

- (a) Identify all continuous symmetries of the action. Analyze if there are symmetries of type

$$\mathbf{r} \rightarrow \alpha \mathbf{r}, \quad t \rightarrow \beta t, \quad \alpha, \beta = \text{const.} \quad (1.2)$$

which do not change the equations of motion.

Respuesta: El lagrangiano será

$$L = \frac{1}{2} m \left(\dot{r}^2 + r^2 \dot{\theta}^2 + r^2 \dot{\phi}^2 \sin^2 \theta \right) - \frac{\kappa}{2t^2} r^2 \quad (1.3)$$

La variable ϕ es cíclica, por lo que M_z será conservado. Aun más, el sistema es esféricamente simétrico, por lo que todo el vector de momento angular se conservará:

$$\mathbf{M} = \text{const.} \quad (1.4)$$

Reemplazando $\mathbf{r} \rightarrow \alpha \mathbf{r}, t \rightarrow \beta t$ en las derivadas,

$$\frac{\partial r}{\partial t} \rightarrow \frac{\partial \alpha r}{\partial \beta t} = \frac{\alpha}{\beta} \frac{\partial r}{\partial t} \quad (1.5)$$

Luego en el lagrangiano

$$\begin{aligned} L' &= \frac{1}{2} m \left(\frac{\alpha^2}{\beta^2} \dot{r}^2 + \frac{\alpha^2}{\beta^2} r^2 \dot{\theta}^2 + \frac{\alpha^2}{\beta^2} r^2 \dot{\phi}^2 \sin^2 \theta \right) - \frac{\kappa}{2\beta^2 t^2} \alpha^2 r^2 \\ L' &= \frac{\alpha^2}{\beta^2} \left[\frac{1}{2} m \left(\dot{r}^2 + r^2 \dot{\theta}^2 + r^2 \dot{\phi}^2 \sin^2 \theta \right) - \frac{\kappa}{2t^2} r^2 \right] \\ L' &= \frac{\alpha^2}{\beta^2} L \end{aligned}$$

Ya que multiplicar un lagrangiano por una cantidad constante no cambia las ecuaciones de movimiento, tendremos una simetría sí y solo sí $\alpha, \beta \in \mathbb{R}, \beta \neq 0$.

Sin embargo, al considerar el efecto sobre la acción,

$$S = \int L dt \rightarrow S' = \int L' dt' = \int \frac{\alpha^2}{\beta^2} L \beta dt = \frac{\alpha^2}{\beta} S$$

Para que podamos usar el teorema de Noether la acción deberá ser invariante, por lo que

$$\begin{aligned} \frac{\alpha^2}{\beta} = 1 &\Rightarrow \alpha = \sqrt{\beta} \\ \Rightarrow \mathbf{r} \rightarrow \alpha \mathbf{r}, \quad t \rightarrow \alpha^2 t, \quad \alpha = \text{const.} \end{aligned} \tag{1.6}$$

■

- (b) Construct the integrals of motion which might exist according to Noether's theorem. Demonstrate that these quantities are conserved using the equations of motion.

Respuesta: El momento angular vendrá dado por

$$\mathbf{M} = m \mathbf{r} \times \dot{\mathbf{r}} = \text{const.} \tag{1.7}$$

Para la simetría de escala, escribiendo las transformaciones en notación con ε

$$\alpha = e^{\varepsilon(t)} \approx 1 + \varepsilon(t)$$

tal que α es generado por cambios infinitesimales ε , los cambios de coordenadas y tiempo será,, entonces

$$\begin{aligned} \mathbf{r}' &\approx \mathbf{r} + \delta \mathbf{r} \approx \mathbf{r} + \varepsilon \mathbf{r}, & t' &\approx t + \delta t \approx t + (1 + \varepsilon)^2 t \approx t + 2\varepsilon t \\ \delta \mathbf{r} &\approx \varepsilon \mathbf{r}, & \delta t &\approx 2\varepsilon t \end{aligned}$$

$$\begin{aligned} \Rightarrow d\mathbf{r}' &= d\left(e^{\varepsilon(t)} \mathbf{r}\right) \\ &= e^\varepsilon d\mathbf{r} + \mathbf{r} e^\varepsilon d\varepsilon \\ &= e^\varepsilon \dot{\mathbf{r}} dt + \mathbf{r} e^\varepsilon \dot{\varepsilon} dt \\ &\approx (1 + \varepsilon) \dot{\mathbf{r}} dt + \mathbf{r} (1 + \varepsilon) \dot{\varepsilon} dt \\ &\approx \underbrace{\dot{\mathbf{r}} dt}_{d\mathbf{r}} + \underbrace{\varepsilon \dot{\mathbf{r}} dt}_{\mathcal{O}(\varepsilon)} + \underbrace{\mathbf{r} \dot{\varepsilon} dt}_{\mathcal{O}(\dot{\varepsilon})} + \underbrace{\mathbf{r} \varepsilon \dot{\varepsilon} dt}_{\mathcal{O}(\varepsilon \dot{\varepsilon})} \\ \\ \Rightarrow dt' &= d\left(e^{2\varepsilon(t)} t\right) \\ &= e^{2\varepsilon} dt + t e^{2\varepsilon} \cdot 2d\varepsilon \\ &= e^{2\varepsilon} dt + 2t e^{2\varepsilon} \dot{\varepsilon} dt \\ &\approx (1 + \varepsilon)^2 dt + 2t(1 + \varepsilon)^2 \dot{\varepsilon} dt \\ &\approx (1 + 2\varepsilon + \varepsilon^2) dt + 2t(1 + 2\varepsilon + \varepsilon^2) \dot{\varepsilon} dt \\ &\approx dt + \underbrace{2\varepsilon dt}_{\mathcal{O}(\varepsilon)} + \underbrace{\varepsilon^2 dt}_{\mathcal{O}(\varepsilon^2)} + \underbrace{2t \dot{\varepsilon} dt}_{\mathcal{O}(\dot{\varepsilon})} + \underbrace{4t \varepsilon \dot{\varepsilon} dt}_{\mathcal{O}(\varepsilon \dot{\varepsilon})} + \underbrace{2t \varepsilon^2 \dot{\varepsilon} dt}_{\mathcal{O}(\varepsilon^2 \dot{\varepsilon})} \end{aligned}$$

Todos los términos $\varepsilon\dot{\varepsilon}$, ε^2 , $\varepsilon^2\dot{\varepsilon}$ son insignificantes, y después de esto deberemos diferenciar por $\dot{\varepsilon}$, por lo que solo nos interesarán los términos $\mathcal{O}(\dot{\varepsilon})$

$$d\mathbf{r}' \approx (\dot{\mathbf{r}} + \dot{\varepsilon}\mathbf{r}) dt, \quad dt' \approx (1 + 2\dot{\varepsilon}t) dt$$

Luego, la velocidad

$$\dot{\mathbf{r}}' \equiv \frac{d\mathbf{r}'}{dt'} \approx \frac{(\dot{\mathbf{r}} + \dot{\varepsilon}\mathbf{r}) dt}{(1 + 2\dot{\varepsilon}t) dt}$$

Aproximando la división con **Mathematica**,

$$\dot{\mathbf{r}}' \approx \dot{\mathbf{r}} + \delta\dot{\mathbf{r}} \approx \dot{\mathbf{r}} + \dot{\varepsilon}(\mathbf{r} - 2\dot{\mathbf{r}}t)$$

$$\delta\dot{\mathbf{r}} \approx \dot{\varepsilon}(\mathbf{r} - 2\dot{\mathbf{r}}t)$$

Luego, aproximando el lagrangiano a primer orden,

$$\begin{aligned} L'(\mathbf{r}', \dot{\mathbf{r}}', t') &= L'(\mathbf{r} + \delta\mathbf{r}, \dot{\mathbf{r}} + \delta\dot{\mathbf{r}}, t + \delta t) \\ &\approx L(\mathbf{r}, \dot{\mathbf{r}}, t) + \frac{\partial L}{\partial \mathbf{r}} \cdot \delta\mathbf{r} + \frac{\partial L}{\partial \dot{\mathbf{r}}} \cdot \delta\dot{\mathbf{r}} + \frac{\partial L}{\partial t} \delta t \\ &\approx L(\mathbf{r}, \dot{\mathbf{r}}, t) + \frac{\partial L}{\partial \mathbf{r}} \cdot \varepsilon\mathbf{r} + \frac{\partial L}{\partial \dot{\mathbf{r}}} \cdot \dot{\varepsilon}(\mathbf{r} - 2\dot{\mathbf{r}}t) + \frac{\partial L}{\partial t} 2\varepsilon t \end{aligned}$$

De nuevo, solo nos interesarán los términos $\mathcal{O}(\dot{\varepsilon})$, por lo que, considerando que $\partial L / \partial \dot{\mathbf{r}} = m\dot{\mathbf{r}}$

$$L' \approx L + m\dot{\mathbf{r}} \cdot \dot{\varepsilon}(\mathbf{r} - 2\dot{\mathbf{r}}t)$$

Por lo que la acción

$$\begin{aligned} S' &= \int L' dt' \\ &\approx \int [L + m\dot{\mathbf{r}} \cdot \dot{\varepsilon}(\mathbf{r} - 2\dot{\mathbf{r}}t)] [1 + 2\dot{\varepsilon}t] dt \\ &\approx \int L dt + \int m\dot{\mathbf{r}} \cdot \dot{\varepsilon}(\mathbf{r} - 2\dot{\mathbf{r}}t) dt + \int 2\dot{\varepsilon}tL dt + \int 2m\dot{\mathbf{r}} \cdot \dot{\varepsilon}^2 t(\mathbf{r} - 2\dot{\mathbf{r}}t) dt \end{aligned}$$

El último término es $\mathcal{O}(\dot{\varepsilon}^2)$, por lo que es insignificante. Tendremos entonces

$$S' \approx \int [L + m\dot{\mathbf{r}} \cdot \dot{\varepsilon}(\mathbf{r} - 2\dot{\mathbf{r}}t) + 2\dot{\varepsilon}tL] dt$$

Y la variación de L será

$$\begin{aligned} S' &\approx \int [L + \delta L] dt \approx \int [L + \dot{\varepsilon}(m\dot{\mathbf{r}} \cdot (\mathbf{r} - 2\dot{\mathbf{r}}t) + 2tL)] dt \\ \delta L &\approx \dot{\varepsilon}[m\dot{\mathbf{r}} \cdot (\mathbf{r} - 2\dot{\mathbf{r}}t) + 2tL] \end{aligned}$$

Finalmente, la cantidad conservada será

$$\begin{aligned}
Q &= \frac{\partial \delta L}{\partial \dot{\epsilon}} \\
&= m\dot{\mathbf{r}} \cdot (\mathbf{r} - 2\dot{\mathbf{r}}t) + 2tL \\
&= \mathbf{p} \cdot \mathbf{r} - 2m\dot{\mathbf{r}}^2 t + 2t \left(\frac{1}{2}m\dot{\mathbf{r}}^2 - U \right) \\
&= \mathbf{p} \cdot \mathbf{r} - 2t \left(\frac{1}{2}m\dot{\mathbf{r}}^2 + U \right) \\
&= \mathbf{p} \cdot \mathbf{r} - 2Ht
\end{aligned}$$

■

- (c) We have seen in lectures that in static (time-independent) Coulomb potential the particle's trajectory always remains in the plane (so-called scattering plane). Analyze if this property remains valid in the time-dependent potential given in Eq. (1). Analyze if the Kepler's laws remain valid in this potential field.

Respuesta: El potencial es central y el momento angular $\mathbf{M}$ es conservado componente por componente, por lo que el plano de rotación será constante.

La segunda ley de Kepler se conservará porque el campo es central; la primera y tercera no lo harán, ya que dependen de una trayectoria específica que en este caso será alterada.

■

2 Problema 2

A relativistic particle of mass m moves in the spherically symmetric potential $U(r) = \alpha/r + \beta/r^2$.

- (a) Write out the lagrangian of the system, and the corresponding equations of motion.

Respuesta: En el marco del laboratorio,

$$\begin{aligned}
L &= -mc^2 \sqrt{1 - \frac{\dot{\mathbf{r}}^2}{c^2}} - \left(\frac{\alpha}{r} + \frac{\beta}{r^2} \right) \\
&= -mc \sqrt{c^2 - \dot{r}^2 - r^2 \dot{\theta}^2 - r^2 \dot{\phi}^2 \sin^2 \theta} - \frac{\alpha}{r} - \frac{\beta}{r^2}
\end{aligned} \tag{2.1}$$

Usando **Mathematica**, los momentos generalizados:

$$p_r = \frac{\partial L}{\partial \dot{r}} = \gamma m \dot{r}, \quad p_\theta = \frac{\partial L}{\partial \dot{\theta}} = \gamma m r^2 \dot{\theta}, \quad p_\phi = \frac{\partial L}{\partial \dot{\phi}} = \gamma m r^2 \dot{\phi} \sin^2 \theta$$

Tenemos las ecuaciones de Euler-Lagrange:

$$\frac{dp_i}{dt} = \frac{\partial L}{\partial q_i}$$

$$\frac{d}{dt} [\gamma m \dot{r}] = \frac{\partial}{\partial r} \left[-mc \sqrt{c^2 - \dot{r}^2 - r^2 \dot{\theta}^2 - r^2 \dot{\phi}^2 \sin^2 \theta} - \frac{\alpha}{r} - \frac{\beta}{r^2} \right]$$

- (b) Identify all symmetries of the action and construct all integrals of motion which might exist in this problem.

Respuesta:

- (c) Try to integrate the equations of motion and find trajectories of bound orbits.

Respuesta:

- (d) Analyze the behavior of your solution in the limit $c \rightarrow \infty$. Assuming that c is large yet finite, find explicitly the first relativistic correction to the trajectory.

Respuesta:

3 Problema 3

In the lectures 12-13 we discussed that the covariant equations of motion for the massive particle in the electromagnetic field A^μ may be obtained from the action

$$S = \int d\tau \Lambda, \quad \Lambda = -mc^2 \sqrt{\dot{x}_\mu \dot{x}^\mu} + e \dot{x}_\mu A^\mu(x) \quad (3.1)$$

where τ is the proper time, $x^\mu(\tau)$ is the trajectory of the particle, and $\dot{x}^\mu = dx^\mu/d\tau$ is the 4-velocity of the particle. Assume that we replaced the covariant lagrangian 3.1 with

$$\tilde{\Lambda} = \frac{1}{2} mc^2 \dot{x}_\mu \dot{x}^\mu + e \dot{x}_\mu A^\mu(x) \quad (3.2)$$

- (a) Demonstrate that the action S is covariant under transformations of the Lorentz group when we use the lagrangian 3.2.

Respuesta: S es covariante porque sus componentes ($\tilde{\Lambda}$ y τ) están escritas usando objetos covariantes. ■

- (b) Write out the generalized momentum $\mathcal{P}_\mu = d\tilde{\Lambda}/d\dot{x}^\mu$ and the equations of motion which correspond to the lagrangian 3.2.

Respuesta: El momento

$$\begin{aligned}
\mathcal{P}_\mu &= \frac{d\tilde{\Lambda}}{d\dot{x}^\mu} \\
&= \frac{d}{d\dot{x}^\mu} \left[\frac{1}{2} mc^2 \dot{x}_\alpha \dot{x}^\alpha + e \dot{x}_\gamma A^\gamma(x) \right] \\
&= \frac{1}{2} mc^2 \frac{d}{d\dot{x}^\mu} [\eta_{\alpha\beta} \dot{x}^\beta \dot{x}^\alpha] + e A^\gamma(x) \frac{d}{d\dot{x}^\mu} [\eta_{\lambda\gamma} \dot{x}^\lambda] \\
&= \frac{1}{2} mc^2 \eta_{\alpha\beta} [\delta_\mu^\beta \dot{x}^\alpha + \dot{x}^\beta \delta_\mu^\alpha] + e A^\gamma(x) \eta_{\lambda\gamma} \delta_\mu^\lambda \\
&= \frac{1}{2} mc^2 [\eta_{\alpha\mu} \dot{x}^\alpha + \eta_{\mu\beta} \dot{x}^\beta] + e A^\gamma(x) \eta_{\mu\gamma} \\
&= \frac{1}{2} mc^2 [\dot{x}_\mu + \dot{x}_\mu] + e A_\mu(x) \\
&= mc^2 \dot{x}_\mu + e A_\mu(x)
\end{aligned}$$

Las ecuaciones de Euler-Lagrange:

$$\begin{aligned}
\frac{\partial \tilde{\Lambda}}{\partial x^\mu} &= \frac{d\mathcal{P}_\mu}{d\theta} \\
e \dot{x}_\alpha \partial_\mu A^\alpha(x) &= mc^2 \ddot{x}_\mu + e \frac{dA_\mu(x)}{d\tau} \\
\Rightarrow mc^2 \ddot{x}_\mu &= e \left(\dot{x}_\alpha \partial_\mu A^\alpha(x) - \frac{\partial A_\mu(x)}{\partial x^\beta} \frac{dx^\beta}{d\tau} \right) \\
mc^2 \ddot{x}_\mu &= e (\eta_{\alpha\lambda} \dot{x}^\lambda \partial_\mu A^\alpha - \dot{x}^\beta \partial_\beta A_\mu) \\
mc^2 \ddot{x}_\mu &= e (\dot{x}^\lambda \partial_\mu A_\lambda - \dot{x}^\beta \partial_\beta A_\mu) \\
mc^2 \ddot{x}_\mu &= e \dot{x}^\nu (\partial_\mu A_\nu - \partial_\nu A_\mu)
\end{aligned}$$

■

- (c) Demonstrate that your equations of motion are equivalent to the equations of motion found in lectures 12,13 using the lagrangian 3.1. Demonstrate that the same equations can be obtained with any lagrangian $\bar{\Lambda}$ given by

$$\bar{\Lambda} = mc^2 f(\dot{x}_\mu \dot{x}^\mu) + e \dot{x}_\mu A^\mu(x) \quad (3.3)$$

where $f(y)$ is an arbitrary function which satisfies $\partial f(y)/\partial y = 1/2$ at $y = 1$ (clearly, the lagrangians (3.1, 3.2) correspond to $f(y) = \sqrt{y}$ and $f(y) = y/2$, respectively).

Respuesta: Los momentos generalizados del primer lagrangiano serán

$$\mathcal{P}_\mu = \frac{d\Lambda}{d\dot{x}^\mu} \quad (3.4)$$

$$= \frac{d}{d\dot{x}^\mu} \left[-mc^2 \sqrt{\dot{x}_\mu \dot{x}^\mu} + e\dot{x}_\mu A^\mu(x) \right] \quad (3.5)$$

$$= -mc^2 \frac{d}{d\dot{x}^\mu} \sqrt{\eta_{\alpha\beta} \dot{x}^\alpha \dot{x}^\beta} + eA^\lambda(x) \frac{d}{d\dot{x}^\mu} (\eta_{\lambda\gamma} \dot{x}^\gamma) \quad (3.6)$$

$$= -mc^2 \frac{d}{d\dot{x}^\mu} \sqrt{\eta_{\alpha\beta} \dot{x}^\alpha \dot{x}^\beta} + eA^\lambda(x) \eta_{\lambda\gamma} \delta_\mu^\gamma \quad (3.7)$$

- (d) Find the nonrelativistic lagrangian L which corresponds to the lagrangian 3.2. Pay attention that the nonrelativistic lagrangian L contributes to the action as $S = \int L dt$.