

Theory & Manual

for

AlkCalc

Simon Euchner*

20.08.2025

* Electronic Mail: euchner.se@gmail.com

Contents

I. Introduction	1
II. Theory	1
i. Single-atom/ion theory	1
ii. Numerical method	5
iii. Radial matrix elements	8
iv. Oscillator strengths	9
v. State lifetimes	10
III. Manual	12
References	20

I. Introduction

Bound single-electron eigenfunctions of alkali-metal atoms and alkaline-earth metal ions are of significant relevance in quantum computing, quantum information processing, quantum simulation, Rydberg physics, quantum optics, However, except for the special case of the Hydrogen atom, it is not possible to solve the time-independent Schrödinger equation analytically. Because of this, there is the need to obtain eigenenergies and eigenstates numerically. The library `AlkCalc` is designed for exactly this purpose: it allows the user to compute the relevant part of the bound-state energy spectrum with the associated eigenstates.

The software `AlkCalc` is split into two parts: a first part used to compute the eigenenergies and eigenstates *once* and store the resulting data. And a second part, referred to as the *library functions*, which implement useful functionality for reading and processing the precomputed atomic data. For instance, there is a function to compute radial transition matrix elements.

In the various README files included in the file tree of `AlkCalc`, technical details for computing the eigenenergies and eigenstates are provided. However, these files do not include a discussion on the theory and physics behind the code. Further, there is no reference manual for the library functions.

This document's purpose is to add the missing pieces: in Sec. II the theory and the physics associated with the code are discussed. Section III is a comprehensive reference manual for the library functions of `AlkCalc`.

II. Theory

This section is organised as follows: in Subsec. i the theory and the physics of single-atom/ion systems are discussed. In Subsec. ii the numerical method based on B-splines is presented. In Subsec. iii it is shown how radial transitions matrix elements are computed by `AlkCalc`. In Subsec. iv it is shown how oscillator strengths for electronic transitions are computed. Finally, in Subsec. v it is shown how lifetimes of electronic states are computed by `AlkCalc`.

i. Single-atom/ion theory

The starting point is an alkali-metal atom or alkaline-earth-metal ion. We are interested in systems where all electrons are in closed shells except for one valence electron. Examples of such systems are ^1H , ^{38}Rb , $^{88}\text{Sr}^+$. This property allows a Kepler-problem description for the ‘effective nucleus’ (the nucleus ‘screened’ by the closed-shell electrons) and the valence electron. The dynamics of the valence electron and the effective nucleus can be described with the following Hamiltonian:

$$H = \frac{\mathbf{p}^2}{2m} + V(r; L^2, S^2, J^2). \quad (\text{II.1})$$

Here, $m = Mm_e/(M + m_e)$ is the reduced mass of the system, composed of the electron’s mass m_e and the atom’s or ion’s mass M . The centre-of-mass motion is decoupled and $\mathbf{r}$, $\mathbf{p}$ are the relative canonical co-ordinates, i.e., $\mathbf{r}$ is the distance vector between the centre-of-mass position of the effective nucleus and the position of the valence electron. Further, $\mathbf{L}$, $\mathbf{S}$, and $\mathbf{J} = \mathbf{L} + \mathbf{S}$ are the

orbital angular momentum operator, the electron's spin operator, and the total angular momentum operator, respectively. Finally, the radial potential V , depending on the distance $r = \|\mathbf{r}\|$, describes the interaction between the valence electron and the effective nucleus. It accounts for several effects: far from the nucleus the valence electron experiences a Coulomb potential with effective charge $Z_{\text{eff}} = 1$ for alkali-metal atom and $Z_{\text{eff}} = 2$ for singly ionised alkaline-earth metal atoms. This is because the nucleus of charge Z is 'screened' by the $Z - 1$ (or $Z - 2$, for alkaline-earth metal ions) ground-state electrons in closed-shells. However, valence electrons with orbits sufficiently close to the nucleus experience the non-trivial charge distribution posed by the nucleus in conjugation with the tightly bound electron cloud. This leads to short-ranged corrections to the effective Coulomb potential with charge Z_{eff} at long distances. Following Refs. [1–4] these corrections are described by the following potential:

$$V(r; \mathbf{L}^2, \mathbf{S}^2, J^2) = V_C(r) + V_P(r) + V_R(r; \mathbf{L}^2, \mathbf{S}^2, J^2). \quad (\text{II.2})$$

The first contribution is

$$V_C(r) = -\frac{e^2}{4\pi\epsilon_0} \frac{Z_n(r)}{r}, \quad Z_n(r) = Z_c + (Z - Z_c)e^{-k_1 r} + k_2 r e^{-k_3 r} - k_4 r^2 e^{-k_3 r}. \quad (\text{II.3})$$

This is a modified Coulomb potential which has the tail of a Coulomb potential with charge Z_{eff} and short-ranged corrections for small radii accounting for the charge-distribution of the effective nucleus. Here, the parameters k_j ($j = 1, 2, 3, 4$) are empirically determined from spectroscopic data [1, 2]. The second contribution to the potential (II.2) is

$$V_P(r) = -\frac{e^2}{(4\pi\epsilon_0)^2} \frac{\alpha_d}{2r^4} \left[1 - \exp\left(-\frac{r^6}{r_c^6}\right) \right]. \quad (\text{II.4})$$

It accounts for the polarisability of the screened nucleus, which gets polarised by the presence of the valence electron. Here, r_c is the effective nucleus' size and α_d its static electric polarisability. Finally,

$$V_R(r; \mathbf{L}^2, \mathbf{S}^2, J^2) = \frac{1}{2m^2 c^2} \frac{V'_{\text{NR}}(r)}{rN(r)} \mathbf{L} \cdot \mathbf{S}, \quad N(r) = \left[1 - \frac{V_{\text{NR}}(r)}{2mc^2} \right]^2, \quad V_{\text{NR}}(r) = V_C(r) + V_P(r). \quad (\text{II.5})$$

It describes (relativistic) coupling between the orbital angular momentum and the valence electrons' spin. This expression results from relativistic perturbation theory [3]. Here, V_{NR} is the non-relativistic model potential and c the speed of light. In Ref. [4] the potential (II.5) is employed for the treatment of alkaline-earth metal ions. In Ref. [5] the simplified version

$$V_R(r) = \frac{\alpha}{2r^3} \mathbf{L} \cdot \mathbf{S} \quad (\text{II.6})$$

is employed for alkali-metal atoms. Here, α is the fine-structure constant. Throughout `AlkCalc` we employ the relativistically correct version in Eq. (II.5), not the simplified version in Eq. II.6. Finally, for expressing the spin-operator $\mathbf{S}$ in Eq. (II.5) and Eq. (II.6) the representation in Pauli's (spin) matrices is chosen:

$$\mathbf{S} = \frac{\hbar}{2} \begin{bmatrix} \sigma_x \\ \sigma_y \\ \sigma_z \end{bmatrix}. \quad (\text{II.7})$$

The Pauli spin matrices are

$$\sigma_x = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}, \quad \sigma_y = \begin{bmatrix} 0 & -i \\ i & 0 \end{bmatrix}, \quad \sigma_z = \begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix}. \quad (\text{II.8})$$

This simple choice for representing the spin is possible, because throughout this work we are interested only in single electron systems with total spin $s = 1/2$.

At this point the Hamiltonian which determines the eigenenergies and the eigenstates is defined. In the following, the goal is to cast eigenvalue problem posed by Hamiltonian (II.1) into a more practical form for the numerical procedure. The first step is to rewrite the product $\mathbf{L} \cdot \mathbf{S}$ in terms of the total angular momentum operator $\mathbf{J} = \mathbf{L} + \mathbf{S}$:

$$\mathbf{L} \cdot \mathbf{S} = \frac{1}{2} [\mathbf{J}^2 - \mathbf{L}^2 - \mathbf{S}^2]. \quad (\text{II.9})$$

Next, we express the square of the momentum operator in spherical co-ordinates,

$$\mathbf{p}^2 = (-i\hbar\nabla)^2 = -\hbar^2\nabla^2 = -\frac{\hbar^2}{r^2} \frac{\partial}{\partial r} \left(r^2 \frac{\partial}{\partial r} \right) + \frac{\mathbf{L}^2}{r^2}. \quad (\text{II.10})$$

The second contribution is the centrifugal barrier. In a classical Kepler problem this is what prevents the electron crashing into the nucleus. Expressed in the spherical coordinates, Hamiltonian (II.1) takes the form

$$H = -\frac{\hbar^2}{2mr^2} \frac{\partial}{\partial r} \left(r^2 \frac{\partial}{\partial r} \right) + V_{\text{eff}}(r; \mathbf{L}^2, \mathbf{S}^2, \mathbf{J}^2), \quad V_{\text{eff}}(r; \mathbf{L}^2, \mathbf{S}^2, \mathbf{J}^2) = V(r; \mathbf{L}^2, \mathbf{S}^2, \mathbf{J}^2) + \frac{\mathbf{L}^2}{2mr^2}. \quad (\text{II.11})$$

The full Hilbert space this Hamiltonian acts on can be split into an angular momentum factor and a radial one. A basis of the angular momentum Hilbert space is given by the simultaneous eigenstates Φ_{l,s,j,m_j} of the operators (Cartan subalgebra) $\mathbf{L}^2$, $\mathbf{S}^2$, $\mathbf{J}^2$, and J_z . Note that a simultaneous eigenbasis of $\mathbf{L}^2$, $\mathbf{S}^2$, $\mathbf{J}^2$, and J_z exists, because these operators commute pairwise. The so-called *spinors* Φ_{l,s,j,m_j} constitute the so-called *coupled basis*, also known as the *fine-structure* basis in the literature. These spinors are given by linear combinations of the uncoupled basis states

$$Y_{l,m_l} \otimes \chi_{s,m_s} \quad (\text{II.12})$$

via SU(2) Clebsch-Gordan coefficients [6]. Here, $Y_{l,m_l} : [0, \pi] \times [0, 2\pi] \rightarrow \mathbb{C}$ are spherical harmonics defined as

$$Y_{l,m_l}(\vartheta, \varphi) = \sqrt{\frac{2l+1}{4\pi} \frac{(l-m_l)!}{(l+m_l)!}} P_l^{m_l}(\cos(\vartheta)) e^{im_l\varphi}. \quad (\text{II.13})$$

In this expression $P_l^{m_l} : [-1, 1] \rightarrow \mathbb{R}$ are the associated Legendre polynomials defined for $l = 0, 1, 2, \dots$ and $m_l = 0, 1, \dots, l-1, l$. These polynomials can be computed via the recursion

$$(l-m_l)P_l^{m_l}(x) = x(2l-1)P_{l-1}^{m_l}(x) - (l+m_l-1)P_{l-2}^{m_l}(x), \quad (\text{II.14})$$

with starting point

$$P_{m_l}^{m_l}(x) = (-1)^{m_l} \frac{(2m_l)!}{2^{m_l} m_l!} (1-x^2)^{m_l/2}, \quad (\text{II.15})$$

in combination with $P_l^{m_l}(x) = 0$ for $l < m_l$. Negative values of m_l are treated with the relation

$$P_l^{-m_l} = (-1)^{m_l} \frac{(l - m_l)!}{(l + m_l)!} P_l^{m_l}. \quad (\text{II.16})$$

In this definition of the associated Legendre polynomial the so-called *Condon-Shortley phase* is included. This follows the definition in Ref. [6].¹ Finally, χ_{s,m_s} in Eq. (II.12) are two-spinors defined as

$$\chi_{\frac{1}{2},\frac{1}{2}} = \begin{bmatrix} 1 \\ 0 \end{bmatrix}, \quad \chi_{\frac{1}{2},-\frac{1}{2}} = \begin{bmatrix} 0 \\ 1 \end{bmatrix}. \quad (\text{II.17})$$

Explicitly (for the case of interest $s = 1/2$), we find the following expressions for the coupled basis states Φ_{l,s,j,m_j} :

$$\Phi_{l,s,j,m_j} = \langle Y_{l,m_j-s} \otimes \chi_{s,s}, \Phi_{l,s,j,m_j} \rangle Y_{l,m_j-s} \otimes \chi_{s,s} + \langle Y_{l,m_j+s} \otimes \chi_{s,-s}, \Phi_{l,s,j,m_j} \rangle Y_{l,m_j+s} \otimes \chi_{s,-s}, \quad (\text{II.18})$$

where the scalar products are (possibly vanishing) Clebsch-Gordan coefficients. These spinor transform according to irreducible representations of $\text{SU}(2)$. This is seen in the following way: consider the unitary and faithful representation $\rho(\alpha) = \exp(i\alpha \cdot J)$, where $\alpha \in \mathbb{R}^3$ is a vector of length $\|\alpha\| \in [0, 2\pi)$. Its action on the spinors is

$$\rho(\alpha)\Phi_{l,s,j,m_j}(\vartheta, \varphi) = \sum_{q=1}^{d_j} D^{(\Gamma_j)}(\alpha)_{m_j,q} \Phi_{l,s,j,q}(\vartheta, \varphi). \quad (\text{II.19})$$

In this expression, d_j is the dimension of the irreducible representation Γ_j of $\text{SU}(2)$ and $D^{(\Gamma_j)}$ is a unitary matrix representation of the irreducible representation Γ_j .

To rewrite the Hamiltonian the crucial step is to use the fact that it has $\text{SU}(2)$ symmetry, i.e., $[\rho(\alpha), H] = 0$ for all α .² Due to this symmetry the Hamiltonian is automatically block diagonal with block dimensions given by integer multiplicities μ_j of the dimensions $d_j = 2j + 1$ of the irreducible representations Γ_j labelled by $j = 0, 1/2, 1, 3/2, 2, \dots$. The multiplicities μ_j are $\mu_j = 2$ for $j \neq 0$ and $\mu_0 = 0$. This is because for each $l > 0$ there are the two possible orbital angular momentum quantum numbers $l = j - 1/2$ and $l = j + 1/2$ and for $s = 1/2$ a total angular momentum of $j = 0$ is not possible. This entails that the Hamiltonian in the basis of the spinors Φ_{l,s,j,m_j} can be decomposed into the following direct sum:

$$\bigoplus_{l=0}^{\infty} \bigoplus_{j \in \{|l-s|, l+s\}} \bigoplus_{m_j=-j}^j H_{l,s,j,m_j}, \quad H_{l,s,j,m_j} = \langle \Phi_{l,s,j,m_j}, H \Phi_{l,s,j,m_j} \rangle. \quad (\text{II.20})$$

Note that the off-diagonal matrix elements within the blocks associated with each irreducible representation Γ_j vanish, i.e., $\langle \Phi_{l,s,j,m'_j}, H \Phi_{l,s,j,m_j} \rangle = 0$ for $m'_j \neq m_j$. This is a result of the $\text{U}(1) \subset$

1 This definition entails that the factor $(-1)^{m_l}$ does not explicitly appear in the definition of the spherical harmonics. This remark is placed here to avoid confusion of whether the spherical harmonics are with or without the Condon-Shortley phase. In the definition of Y_{l,m_l} used in this work, the Condon-Shortley phase is included.

2 The symmetry group is even bigger because parity is also conserved. So the full symmetry group is $\text{SU}(2) \times \mathbb{Z}_2$.

SU(2) symmetry of the full Hamiltonian in Eq. (II.1), which is associated to the generator L_z . This symmetry further entails that the diagonal matrix elements $\langle \Phi_{l,s,j,m_j}, H \Phi_{l,s,j,m_j} \rangle$ do not depend on m_j . Therefore, the full Hamiltonian (II.1) in the basis of the spinors $\Phi_{l,s,j}$ can be expressed as the more simple direct sum

$$\bigoplus_{l=0}^{\infty} \bigoplus_{j \in \{|l-s|, l+s\}} 1_{2j+1} \otimes H_{l,s,j}. \quad (\text{II.21})$$

Here, 1_{2j+1} is the identity matrix of dimension $2j + 1$ and

$$H_{l,s,j} = -\frac{\hbar^2}{2mr^2} \frac{\partial}{\partial r} \left(r^2 \frac{\partial}{\partial r} \right) + V_{\text{eff}}(r; \hbar^2 l[l+1], \hbar^2 s[s+1], \hbar^2 j[j+1]). \quad (\text{II.22})$$

For completeness ρ can be expressed in the basis of $\Phi_{l,s,j}$ as well. It is given by the following direct sum of the irreducible representations Γ_j of SU(2):

$$\rho = \bigoplus_{j>0} 2\Gamma_j. \quad (\text{II.23})$$

The block structure of the Hamiltonian reduces the eigenvalue problem defined by the time-independent Schrödinger equation

$$H\psi = E\Psi \quad (\text{II.24})$$

to the blocks $H_{l,s,j}$ on the Hilbert space associated with the radial degrees of freedom. In particular, solving for the full eigenenergies and eigenstates is reduced to solving the radial eigenvalue problems

$$H_{l,s,j} R_{n,l,s,j} = E_{n,l,s,j} R_{n,l,s,j}, \quad (\text{II.25})$$

where $R_{n,l,s,j}$ is the n -th radial eigenstate of $H_{l,s,j}$. The total eigenstate Ψ is then given by the product of the radial eigenstates and the spinors:

$$\Psi_{n,l,s,j,m_j}(r, \vartheta, \varphi) = R_{n,l,s,j}(r) \Phi_{l,s,j,m_j}(\vartheta, \varphi). \quad (\text{II.26})$$

This shows that solving for the full eigenstates Ψ_{n,l,s,j,m_j} is equivalent to solving for the radial eigenstates $R_{n,l,s,j}$. In general, the eigenenergies $E = E_{n,l,s,j}$ and the radial eigenstates $R_{n,l,s,j}$ must be calculated numerically. This task is exactly what `AlkCalc` is designed to do.

ii. Numerical method

The radial eigenstates, generally, cannot be obtained analytically. `AlkCalc` provides a platform to solve for these states numerically. To make the problem addressable by a computer, we express the eigenproblem in terms of dimensionless quantities. To this end we express the Hamiltonians $H_{l,s,j}$ in units of Hartree, $E_H = 1/(4\pi\epsilon_0 a_B)$, where $a_B = \hbar/(m_e c \alpha) \approx 52.9$ pm is Bohr's radius. The radial eigenfunctions we parametrise as

$$R_{n,l,s,j}(r) = \frac{1}{\sqrt{a_B^3}} \frac{f_{n,l,s,j}(r/a_B)}{r/a_B}. \quad (\text{II.27})$$

where $f_{n,l,s,j}$ is dimensionless. The eigenproblem in Eq. (II.25) in the new eigenfunctions takes the form

$$\left[-\frac{1}{2} \frac{d^2}{dt^2} + \frac{l(l+1)}{2t^2} + C \frac{V(a_B t; \hbar^2 l[l+1], \hbar^2 s[s+1], \hbar^2 j[j+1])}{E_H} \right] f_{n,l,s,j}(t) = \bar{E}_{n,l,s,j} f_{n,l,s,j}(t), \quad (\text{II.28})$$

where we defined

$$C = \frac{1}{1 + m_e/M} \quad (\text{II.29})$$

and the eigenenergies may be extracted from the relation

$$\frac{E_{l,s,j,n}}{E_H} = \frac{\bar{E}_{n,l,s,j}}{C}. \quad (\text{II.30})$$

Rewriting the normalisation condition in terms of the eigenfunctions $f_{n,l,s,j}$ leads to

$$1 = \int_0^\infty dr r^2 R_{n,l,s,j}^2(r) = \int_0^\infty (a_B dt) (a_B t)^2 \frac{1}{a_B^3} \frac{f_{n,l,s,j}^2(t)}{t^2} = \int_0^\infty dt f_{n,l,s,j}^2(t). \quad (\text{II.31})$$

Here we assumed that the (radial) eigenfunctions are real-valued, which is justified as the Hamiltonian is not only an hermitian but also a symmetric operator. In conclusion, computing the normalised eigenfunctions $f_{n,l,s,j}$ yields the (automatically normalised) radial eigenstates by employing Eq. (II.27) for the conversion.

To solve the eigenproblem in Eq. (II.28) numerically, we employ a finite element method (FEM) instead of the conceptually (and programmatically) more simple finite difference method (FDM) for, mainly³, one reason: With a FEM we are completely free in choosing the mesh (the position of the discretisation points) for the radial domain, while keeping the associated finite-dimensional matrix problem real and symmetric. This is important because the problem we face presents itself with drastically different energy scales, introduced by the diverging potential: the eigenfunctions $f_{n,l,s,j}$ oscillate quickly for small arguments and slowly for large ones, where the valence electron is almost free. Thus, a mesh tailored to this behaviour leads to a significant reduction of computational resources (the fast oscillation can be efficiently resolved without introducing ‘too many’ points in the slowly oscillating regime).

The first step in setting-up the FEM is to agree on a finite closed interval $I = [0, t_{\max}]$, where $t_{\max} > 0$ sets the maximal value of the domain we consider. The value $t_{\max}$ must be chosen large enough to ensure that the boundary condition $f_{n,l,s,j}(t \geq t_{\max}) = 0$ is fulfilled. This equal sign is actually never true, because the radial eigenstates asymptotically approach zero. However, if we reach zero up to machine precision, it is numerically true. Of course, one must ensure that the finite set of solutions that shall be computed has support within this interval. The mesh we define as a family $(t_k)_{k=1,\dots,N-1}$ in I with the properties $t_0 = 0$, $t_{N-1} = t_{\max}$, and $t_0 < t_1 < \dots < t_{N-2} < t_{N-1}$, where we say $N \geq 3$. Of course, typical values are $N \sim 10^5$ - 10^7 . Next, we define the derived family

3 It also helps to save some disk space, because one must store much less data to represent the eigenfunctions $f_{n,l,s,j}$. Typically, disk space is not a relevant factor these days, but we see no need to be unnecessarily wasteful with it either.

of step sizes $(h_k)_{k=1,\dots,N-1}$, $h_k = t_k - t_{k-1}$. The elements $(\phi_k)_{k=0,\dots,N-1}$, $\phi_k : I \rightarrow [0, 1]$, we define as

$$\phi_k(t) = \begin{cases} (t - t_{k-1})/(t_k - t_{k-1}), & t \in (t_{k-1}, t_k) \\ (t_{k+1} - t)/(t_{k+1} - t_k), & t \in (t_k, t_{k+1}) \\ 0, & \text{else} \end{cases} \quad (\text{II.32})$$

for $k = 1, \dots, N - 2$, and for $k = 0$ and $k = N - 1$ the Dirichlet boundary conditions $f_{n,l,s,j}(0) = f_{n,l,s,j}(t_{\max}) = 0$ lead to the canonical choice $\phi_0 = \phi_{N-1} \equiv 0$. To obtain a matrix problem which approximates Eq. (II.28) we compute projections of it onto the elements with the real scalar product ('dummy' functions $\alpha, \beta : I \rightarrow \mathbb{R}$)

$$B(\alpha, \beta) = \int_I dt \alpha(t) \beta(t). \quad (\text{II.33})$$

To simplify the notation we fix the quantum numbers n, l, s , and j and say $f \equiv f_{n,l,s,j}$. Moreover, we collect all potential contributions into the new potential v , defined as

$$v(t) = \frac{l(l+1)}{2t^2} + C \frac{V(a_B t; \hbar^2 l[l+1], \hbar^2 s[s+1], \hbar^2 j[j+1])}{E_H}. \quad (\text{II.34})$$

Finally, we say $\lambda = \bar{E}_{n,l,s,j}$, such that Eq. (II.28) simply becomes

$$-\frac{1}{2}f''(t) + v(t)f(t) = \lambda f(t). \quad (\text{II.35})$$

Before we project this equation onto the elements, we approximately expand f in them:

$$f = \sum_{k=1}^{N-2} f_k \phi_k. \quad (\text{II.36})$$

This expansion automatically fulfils the Dirichlet boundary conditions $f(0) = f(t_{\max}) = 0$. Projection, i.e., applying the map $B(\phi_k, \cdot)$ for all $k = 1, \dots, N - 2$, yields a matrix eigenproblem whose solutions converge to solutions of Eq. (II.28) as N tends to infinity:

$$(K + V)f = \lambda Mf. \quad (\text{II.37})$$

In this expression $M_{kk'} = B(\phi_k, \phi_{k'})$ is the so-called mass matrix, $K_{kk'} = B(\phi'_k, \phi'_{k'})/2$ the so-called stiffness matrix⁴, and $V_{kk'} = B(\phi_k, v\phi_{k'})$ the potential energy contribution. Each matrix is $(N - 2)$ -dimensional, since $k, k' = 1, \dots, N - 2$, and M and K can be computed analytically; in particular

$$M = \frac{1}{6} \begin{bmatrix} 2(h_1 + h_2) & h_2 & & & & \\ h_2 & 2(h_2 + h_3) & h_3 & & & \\ & h_3 & 2(h_3 + h_4) & h_4 & & \\ & & \ddots & \ddots & h_{N-2} & \\ & & & h_{N-2} & 2(h_{N-2} + h_{N-1}) & \end{bmatrix} \quad (\text{II.38})$$

⁴ Note that because the elements obey the Dirichlet boundary conditions $\phi_k(0) = \phi_k(t_{\max}) = 0$, $k = 1, \dots, N - 2$, integrating by parts allows to write $B(\phi''_k, \phi_{k'}) = -B(\phi'_k, \phi'_{k'})$.

and

$$K = \frac{1}{2} \begin{bmatrix} \frac{1}{h_1} + \frac{1}{h_2} & -\frac{1}{h_2} & & & \\ -\frac{1}{h_2} & \frac{1}{h_2} + \frac{1}{h_3} & -\frac{1}{h_3} & & \\ & -\frac{1}{h_3} & \frac{1}{h_3} + \frac{1}{h_4} & -\frac{1}{h_4} & \\ & & \ddots & \ddots & \\ & & & -\frac{1}{h_{N-2}} & \frac{1}{h_{N-2}} + \frac{1}{h_{N-1}} \end{bmatrix}. \quad (\text{II.39})$$

Note that both of these matrices are symmetric. The potential matrix, generally, must be computed numerically. However, here we choose the simple approximation

$$B(\phi_k, v\phi_{k'}) \approx \frac{v(t_k) + v(t_{k'})}{2} M_{kk'}, \quad (\text{II.40})$$

which is justified if the step size is small enough. Note that this particular choice of approximations keeps the matrix symmetric, such that $K + V$ is symmetric. Furthermore, M represents a scalar product and is thus positive definite. This means that the matrix eigenproblem in Eq. (II.37) can be efficiently solved with the Lanczos algorithm.

Next, we express the normalisation condition in Eq. (II.31) in terms of our elements and the vector $\mathbf{f}$. To this end we calculate

$$\int_0^\infty dt f^2(t) \approx \int_I dt f^2(t) = \sum_{k=1}^{N-2} \sum_{k'=1}^{N-2} f_k f_{k'} B(\phi_k, \phi_{k'}) = \mathbf{f}^T M \mathbf{f} = \|\mathbf{f}\|_M^2 \stackrel{!}{=} 1, \quad (\text{II.41})$$

i.e., the length of $\mathbf{f}$ must be unity in the by the scalar product M induced norm $\|\cdot\|_M$.

iii. Radial matrix elements

To compute radial matrix elements, we first calculate, for $p \in \mathbb{R}$ such that the integral exists,

$$\begin{aligned} \frac{r_p^{n,l,s,j;n',l',s',j'}}{a_B^p} &\equiv \frac{1}{a_B^p} \langle R_{n,l,s,j}, \text{id}^p R_{n',l',s',j'} \rangle = \int_0^\infty dr r^2 R_{n,l,s,j}(r) \frac{r^p}{a_B^p} R_{n',l',s',j'}(r) \\ &= \int_0^\infty (dt a_B) (a_B t)^2 f_{n,l,s,j}(t) \frac{(a_B t)^p}{a_B^p} \frac{1}{a_B^3} \frac{f_{n',l',s',j'}(t)}{t^2} = B(f_{n,l,s,j}, \text{id}^p f_{n',l',s',j'}). \end{aligned} \quad (\text{II.42})$$

Here, ‘id’ denotes the identity on the non-negative real numbers. Next, we approximate this result with solutions to the matrix eigenproblem in Eq. (II.37), i.e., we represent $f_{n,l,s,j}$ and $f_{n',l',s',j'}$ approximately with the vectors $\mathbf{f}_{n,l,s,j}$ and $\mathbf{f}_{n',l',s',j'}$, respectively [see Eq. (II.36)]. This allows to write

$$B(f_{n,l,s,j}, \text{id}^p f_{n',l',s',j'}) \approx \mathbf{f}_{n,l,s,j}^T R_p \mathbf{f}_{n',l',s',j'}, \quad (\text{II.43})$$

where the components of the real symmetric matrix R_p are defined as

$$(R_p)_{k,k'} = B(\phi_k, \text{id}^p \phi_{k'}) = \begin{cases} B(\phi_k, \text{id}^p \phi_{k+1}), & k' = k + 1 \text{ or } k = k' + 1 \ (0 < k, k' < N - 2) \\ B(\phi_k, \text{id}^p \phi_k), & k' = k \ (0 < k, k' \leq N - 2) \\ 0, & \text{else} \end{cases}. \quad (\text{II.44})$$

To numerically compute the remaining integrals we use a two-point Gaussian quadrature. This method approximates the integral as [7]:

$$\int_a^b dt \alpha(t) \approx \frac{b-a}{2} \left[\alpha \left(\frac{a+b}{2} - \frac{b-a}{2\sqrt{3}} \right) + \alpha \left(\frac{a+b}{2} + \frac{b-a}{2\sqrt{3}} \right) \right], \quad (\text{II.45})$$

for any sufficiently smooth function α and $a < b$. Note that this result is exact if α is a polynomial of degree three, which is the case for radial dipole matrix elements. This method is straightforward to implement in a computer program, such that the problem of calculating matrix elements is effectively reduced to computing matrix products. However, note that for too large p there can be numerical overflow or underflow when computing t^p . But, for 64-bit floating point arithmetic, $p = \pm 10$ is easily possible (if the integral exists), such that, typically, all practical cases are included. Explicitly, we could verify that for Hydrogen the range $p = -2, \dots, 10$ yields reasonably good results. Finally, we remark that one can improve on the precision by employing higher-order Gaussian quadrature algorithms as described in [7].

iv. Oscillator strengths

The so-called ‘oscillator strength’, for a transition from an initial state $\Psi_i \equiv \Psi_{n',l',s,j',m'_j}$ to a final state $\Psi_f \equiv \Psi_{n,l,s,j,m_j}$ is [6]

$$f_{i \rightarrow f} = \sum_{m'_j = -j'}^{j'} \frac{2}{3} \bar{E}_{f,i} |\langle \Psi_f, \mathbf{r} / a_B \Psi_i \rangle|^2, \quad (\text{II.46})$$

with the energy difference $E_H \bar{E}_{f,i} = E_f - E_i$. Note that we sum over all final m'_j , because we are interested in transitions between the fine-structure states without resolving the Zeeman sub-levels. The oscillator strength is a measure for how strongly the induced Hertzian (electric) dipole oscillates: the radiation power is large for spatially large dipoles and fast oscillation frequencies, captured by the matrix element and energy difference, respectively. In principle, we already have all the machinery to numerically compute this quantity, however, one can reduce the expression to a much simpler form by using angular momentum algebra, in particular, relations from [8]. To simplify the expression, we first use (Condon-Shortley phase)

$$r_x = \sqrt{\frac{2\pi}{3}} r (Y_{1,-1} - Y_{1,1}), \quad r_y = i \sqrt{\frac{2\pi}{3}} r (Y_{1,-1} + Y_{1,1}), \quad r_z = \sqrt{\frac{4\pi}{3}} r Y_{1,0}. \quad (\text{II.47})$$

With this, the oscillator strength becomes

$$f_{i \rightarrow f} = \frac{4\pi}{3} \frac{2}{3} \bar{E}_{f,i} \langle r R_f, r^2 / a_B r R_i \rangle^2 \sum_{m'_j = -j'}^{j'} \sum_{q=-1}^1 |\langle \Phi_f, Y_{1,q} \Phi_i \rangle|^2 \quad (\text{II.48})$$

Employing relations from [8, p. 236, 244, 260, 445, 496] one can show that the remaining matrix element is

$$\langle \Phi_f, Y_{1,q} \Phi_i \rangle = (-1)^{j'+j+1+l'+s-m'_j} \sqrt{\frac{3}{4\pi}} \begin{pmatrix} j' & 1 & j \\ -m'_j & q & m_j \end{pmatrix} \sqrt{(2j+1)(2j'+1)(2l+1)} \begin{Bmatrix} l & s & j \\ j' & 1 & l' \end{Bmatrix} C_{l',0}^{l,0,1,0}, \quad (\text{II.49})$$

(l, j)	$\mapsto$	(l', j')	a_l
$(l, l + s)$	$\mapsto$	$(l + 1, l + s)$	$\frac{1}{(2l+3)(2l+1)}$
$(l, l - s)$	$\mapsto$	$(l + 1, l + s)$	$\frac{l+1}{2l+1}$
$(l, l + s)$	$\mapsto$	$(l + 1, l + 3s)$	$\frac{l+2}{2l+3}$
$(l, l - s)$	$\mapsto$	$(l - 1, l - s)$	$\frac{1}{(2l+1)(2l-1)}$
$(l, l + s)$	$\mapsto$	$(l - 1, l - s)$	$\frac{l}{2l+1}$
$(l, l - s)$	$\mapsto$	$(l - 1, l - 3s)$	$\frac{l-1}{2l-1}$

Table 1: Angular factors. Table containing the angular factors a_l [see Eq. (II.50)] for computing oscillator strengths. Throughout this table, $s = 1/2$.

expressed in terms of a Wigner $3jm$ -symbol, a Wigner $6j$ -symbol, and a Clebsch-Gordan coefficient. With symmetry relations and a sum rule from [8, p. 236, 259] we can compute the sum over q and m'_j in the expression for the oscillator strength, which yields

$$\sum_{m'_j} f_{i \rightarrow f} = \frac{2}{3} \bar{E}_{f,i} |\langle r R_{n',l',s,j'}, r^2 / a_B R_{n,l,s,j} \rangle|^2 a_l, \quad a_l = l_{\max}(2j' + 1) \left\{ \begin{matrix} l & s & j \\ j' & 1 & l' \end{matrix} \right\}^2, \quad (\text{II.50})$$

where $l_{\max} = \max(l, l')$. There are only six transitions for which a_l is non-zero. Using relations from [8, p. 298, 300] we arrive at the results shown in Tab. 1. With this it is now straight-forward to compute the oscillator strengths numerically with a computer program. As a final remark, note that the oscillator strength does not depend on m_j any more, i.e., by summing over final states, we find that the oscillators for different m_j all have the same strength.

v. State lifetimes

To compute the lifetime of a fine-structure state, we need to calculate the so-called ‘Einstein-A’ coefficient [6, 9]. It is a rate that describes the number of spontaneous emissions per unit time at zero temperature. The total decay rate, at zero temperature, is given by the sum over all Einstein-A coefficients. In the dipole approximation this sum is given by the expression [6]:

$$A_i = \sum_f A_{i \rightarrow f}, \quad A_{i \rightarrow f} = \frac{E_H}{\hbar} \frac{4}{3} \alpha^2 \bar{E}_{i,f}^3 \sum_{u=x,y,z} |\langle \Psi_f, r_u / a_B \Psi_i \rangle|^2. \quad (\text{II.51})$$

In terms of the oscillator strength we obtain

$$A_i = \frac{2\alpha^3 E_H}{\hbar} \sum_{n',l',j'} \bar{E}_{i,f}^2 (-f_{i \rightarrow f}). \quad (\text{II.52})$$

The rate in Eq. (II.52) is the spontaneous emission rate, which only appears for transitions where $E_i > E_f$. Note that, implicitly, this restricts the quantum numbers the sum is running over. Further, this is the correct decay rate only for zero temperature case. At finite temperature, spontaneous transitions increasing the energy may be driven as well, if thermal fluctuations are sufficiently

strong. The background spectral density is given by Planck's law, which yields for the average photon number at frequency ω and temperature T :

$$n(\omega, T) = \frac{1}{\exp(\hbar\omega/(k_B T)) - 1}, \quad (\text{II.53})$$

where k_B is Boltzmann's constant. From Einstein's rate equations [9] it follows that the total rate for decay out of the initial state is

$$\frac{\hbar}{E_H} \Gamma = 2\alpha^3 \sum_{n', l', j'} \bar{E}_{i,f}^2 (-f_{i \rightarrow f}) (1 + n(E_{i,f}/\hbar, T)) + 2\alpha^3 \sum_{n', l', j'} \bar{E}_{f,i}^2 f_{i \rightarrow f} n(E_{f,i}/\hbar, T), \quad (\text{II.54})$$

where each sum only runs over quantum numbers such that the associated term is positive. That is, the first sum is only over final states with lower, and the second one only over final states with larger, energy than the initial state. With this formula, the lifetime of a fine-structure state is $\tau = \Gamma^{-1}$.

III. Manual

Reference manual for the `AlkCalc` library functions. Prototypes for the following functions are defined in the header ‘`interface/alkcalc.h`’.

There is always a decision to be made when dealing with half-integer quantum numbers numerically. Here we agree on the following: say q is a quantum number that may take half-integer values, e.g., j , m_j . To ‘clean up’ the numerical value we do $2q \mapsto [2.0 \times q + 0.5]$. The ‘floor’ function is defined as $[x] = k \in \mathbb{Z} (x \in \mathbb{R})$, where k is the unique integer such that $x \in [k, k + 1)$. In practice, this means that for any $q \in [-0.75, -0.25)$ we use $q = -1/2$, for $q \in [-0.25, 0.25)$ we use $q = 0$, for $q \in [0.25, 0.75)$ we use $q = 1/2$, and so on, in both the negative and positive direction. In the following manual, this convention is applied to all quantum numbers that can take half-integer values, unless stated otherwise.

FUNCTION

`alkcalc_Enlsj`

DESCRIPTION

Eigenenergies in units of Hartree. The requested eigenenergy is read from the corresponding data file in the directory ‘`data`’ and returned.

ARGUMENTS

`char *species`

String to specify atom/ion species, e.g., `1H` for Hydrogen, or `88SR+` for the $^{88}\text{Sr}^+$ ion.

`int32_t n`

Principal quantum number.

`int32_t l`

Orbital angular momentum quantum number $l = 0, 1, \dots, n - 1$.

`double s`

Electron spin quantum number. Not an argument, as $s = 1/2$ is always assumed.

`double j`

Total angular momentum quantum number $j = |l - 1/2|, l + 1/2$.

RETURN

`double E`

E is the eigenenergy specified by the quantum numbers n , l , $s = 1/2$, and j .

FUNCTION

alkcalc_fnlsj

DESCRIPTION

Numerical equivalent to $f_{n,l,s,j}$ in Eq. (II.27). The data is read from the file containing the requested state. This data is stored at a user specified location. The location where AlkCalc will search for the data can be set in the file 'interface/alkcalc.h'. The result is owned by the caller and should be freed with `alkcalc_state_free`.

ARGUMENTS

char result

Character for deciding if discretisation data is returned; see RETURN down below. If result is f (full), the full result is returned, and if result is p (partial), the discretisation data is not returned.

char *species

String to specify atom/ion species, e.g., 1H for Hydrogen, or 88SR+ for the $^{88}\text{Sr}^+$ ion.

int32_t n

Principal quantum number.

int32_t l

Orbital angular momentum quantum number $l = 0, 1, \dots, n - 1$.

double s

Electron spin quantum number. Not an argument, as $s = 1/2$ is always assumed.

double j

Total angular momentum quantum number $j = |l - 1/2|, l + 1/2$.

RETURN

alkcalc_state *f

f contains eight members: N, dim, t, h, and fnlsj.

int32_t N: Number of points, N , to discretise the interval $[0, t_{\max}]$ (see Sec. II).

int32_t n: Principal quantum number.

int32_t l: Orbital angular momentum quantum number.

double j: Total angular momentum quantum number.

int32_t dim: Dimension of eigenproblem, i.e., $\text{dim} = N - 2$ [see Eq. (II.36)].

double *t: Array of length N , containing the mesh $(t_k)_{k=0,\dots,N-1}$ (see Sec. II).

double *h: Array of length $N - 1$, containing the steps $(h_k)_{k=1,\dots,N-1}$ (see Sec. II).

double *fnlsj: dim-dimensional eigenvector f [see Eq. (II.36)].

FUNCTION

alkcalc_rp

DESCRIPTION

Radial matrix element of r^p , $p \in \mathbb{R}$ [see Eq. (II.42)]. For $|p| \gtrsim 10$ there is the risk of numerical over or underflow of the 64-bit floating-point arithmetic. It is the user's responsibility to ensure that the integral associated to the desired matrix element exists. In case it does not, the returned value is wrong and no error is thrown. ^1H (Hydrogen atom) and $^2\text{He}^+$ (Helium ion) can be used to benchmark atoms and ions, respectively.

ARGUMENTS

char *species

String to specify atom/ion species, e.g., 1H for Hydrogen, or 88SR+ for the $^{88}\text{Sr}^+$ ion.

int32_t nb

Principal quantum number of bra.

int32_t lb

Orbital angular momentum quantum number $l = 0, 1, \dots, n - 1$ of bra.

double sb

Electron spin quantum number. Not an argument, as $s = 1/2$ is always assumed.

double jb

Total angular momentum quantum number of bra, j .

double p

Power of radius operator; cf. Eq. (II.42). For instance, $p = 0$ yields the scalar product of the states. For $p = 1$ and $p = 2$, respectively, dipole and quadrupole matrix elements are computed. Generally, $p \in \mathbb{R}$.

int32_t nk

Principal quantum number of ket.

int32_t lk

Orbital angular momentum quantum number $l = 0, 1, \dots, n - 1$ of ket.

double sk

Electron spin quantum number. Not an argument, as $s = 1/2$ is always assumed.

double jk

Total angular momentum quantum number of ket, j .

RETURN

double rp

$r_p^{n,l,s,j;n',l',s',j'}$ in Eq. (II.42) in units of a_B^p , where a_B is Bohr's radius. Quantum numbers with a prime correspond to the bra, and the ones without one, to the ket.

FUNCTION

alkcalc_cj1m1j2m2jmj

DESCRIPTION

Clebsch-Gordan coefficients, $C_{j,m_j}^{j_1,m_1,j_2,m_2} = \langle j_1, m_1; j_2, m_2 | j_1, j_2, j, m_j \rangle$, where j is the total angular momentum composed of the two individual ones j_1 and j_2 . To fix the phase uniquely, we adopt the Condon-Shortley sign convention, that is, the Clebsch-Gordan coefficients with $m_j = j_1 + j_2$ are positive. In particular, for the case $j_1 = l$ and $j_2 = s$,

$$C_{j,m_j}^{l,m_l,s,m_s} = \langle Y_{l,m_l} \otimes \chi_{s,m_s}, \Phi_{l,s,j,m_j} \rangle, \quad (\text{III.1})$$

with the states $Y_{l,m_l} \otimes \chi_{s,m_s}$ of the uncoupled basis and the coupled basis states $\Phi_{l,s,j,m}$; see Eqs. (II.13, II.17), and Eq. (II.18), respectively. The return type (see RETURN down below) allows to extract the requested coefficient symbolically as well as numerically. The function returns zero in case the arguments do not make sense, e.g., if $j_1 < m_1$; or if j_1 is half-integer and m_1 is integer; or if j does not lay between (including bounds) $|j_1 - j_2|$ and $|j_1 + j_2|$.

ARGUMENTS

double j1

Angular momentum quantum number of first factor.

double m1

Magnetic quantum number of first factor.

double j2

Angular momentum quantum number of second factor.

double m2

Magnetic quantum number of second factor.

double j

Total angular momentum quantum number.

double mj

Magnetic quantum number of total angular momentum.

RETURN

alkcalc_cg c

c contains three members: sign, numerator, and denominator.

int8_t sign: Sign, σ , of the Clebsch-Gordan coefficient.

int64_t numerator: Numerator, ν , of the Clebsch-Gordan coefficient.

int64_t denominator: Denominator, μ , of the Clebsch-Gordan coefficient.

The requested Clebsch-Gordan coefficient is given by $\sigma \sqrt{\nu/\mu}$. Euklid's algorithm is employed to reduce ν and μ . This representation allows to obtain Clebsch-Gordan coefficients symbolically as well as numerically.

FUNCTION

alkcalc_YlmlXsms

DESCRIPTION

Angular factor of the full state in the uncoupled basis, i.e., the two-component spinor $Y_{l,m_l}(\vartheta, \varphi)\chi_{s,m_s}$. The input quantum numbers are checked for inconsistencies and an error is thrown in case one is detected. The angles ϑ and φ have the ranges $[0, \pi]$ and $[0, 2\pi]$, respectively. This condition is checked up to floating point machine precision (type `double`) and an error will be thrown in case either condition is not obeyed.

ARGUMENTS

`int32_t l`

Orbital angular momentum quantum number $l = 0, 1, \dots, n - 1$.

`int32_t ml`

Magnetic quantum number of orbital angular momentum.

`double s`

Electron spin quantum number. Not an argument, as $s = 1/2$ is always assumed.

`double ms`

Magnetic quantum number of electron spin, m_s . For any $m_s < 0$ the value $m_s = -1/2$ is used internally, and for $m_s \geq 0$ the value $m_s = 1/2$.

`double theta`

Polar angle (Zenitwinkel), ϑ , in range $[0, \pi]$.

`double phi`

Azimuthal angle (Azimut), φ , in range $[0, 2\pi]$.

RETURN

`alkcalc_spinor s`

`s` has two components: `u` and `v`.

`u` is the spin-up ($m_s = 1/2$) component.

`v` is the spin-down ($m_s = -1/2$) component.

FUNCTION

alkcalc_Philsjmj

DESCRIPTION

Angular factor of the full eigenstate in the coupled (fine-structure) basis, i.e., the two-component spinor $\Phi_{l,s,j,m_j}(\vartheta, \varphi)$ [see Eq. (II.18)]. The input quantum numbers are checked for inconsistencies and an error is thrown in case one is detected. The angles ϑ and φ have the ranges $[0, \pi]$ and $[0, 2\pi]$, respectively. This condition is checked up to floating point machine precision (type `double`) and an error will be thrown in case either condition is not obeyed. To compute the resulting two-component spinor, Clebsch-Gordon coefficients are needed, for which the sign convention detailed in Eq. (III.1) is employed.

ARGUMENTS

`int32_t l`

Orbital angular momentum quantum number $l = 0, 1, \dots, n - 1$.

`double s`

Electron spin quantum number. Not an argument, as $s = 1/2$ is always assumed.

`double j`

Total angular momentum quantum number $j = |l - 1/2|, l + 1/2$.

`double mj`

Magnetic quantum number of total angular momentum, m_j .

`double theta`

Polar angle (Zenitwinkel), ϑ , in range $[0, \pi]$.

`double phi`

Azimuthal angle (Azimut), φ , in range $[0, 2\pi]$.

RETURN

`alkcalc_spinor s`

`s` has two components: `u` and `v`.

`u` is the spin-up ($m_s = 1/2$) component.

`d` is the spin-down ($m_s = -1/2$) component.

FUNCTION

alkcalc_fitof

DESCRIPTION

Oscillator strength for the transition between an initial state Ψ_i and a final state Ψ_f , as defined in Subsec. iv, that is, it was already summed over the magnetic quantum number of the final state.

ARGUMENTS

char *species

String to specify atom/ion species, e.g., 1H for Hydrogen, or 88SR+ for the $^{88}\text{Sr}^+$ ion.

int32_t ni

Principal quantum number of the initial state Ψ_i .

int32_t li

Orbital angular momentum quantum number $l = 0, 1, \dots, n - 1$ of the initial state Ψ_i .

double si

Electron spin quantum number. Not an argument, as $s = 1/2$ is always assumed.

double ji

Total angular momentum quantum number of the initial state Ψ_i , j .

int32_t nf

Principal quantum number of the final state Ψ_f .

int32_t lf

Orbital angular momentum quantum number $l = 0, 1, \dots, n - 1$ of the final state Ψ_f .

double sf

Electron spin quantum number of the final state Ψ_f . Not an argument here, since $s = 1/2$ is always assumed.

double jf

Total angular momentum quantum number of the final state Ψ_f , j .

RETURN

double fitof

$f_{i \rightarrow f}$ is the oscillator strength of the selected electronic transition, summed over m'_j of the final state. Note that because of this sum, it is the same for all m_j of the initial state.

FUNCTION

alkcalc_tau

DESCRIPTION

Lifetime of the fine-structure state Ψ_{n,l,s,j,m_j} in nanoseconds. Note that, since the oscillator-strengths [see Eq. (II.50)] do not depend on m_j , also the lifetimes do not depend on it [see Eq. (II.54)].

ARGUMENTS

double T

Temperature of black-body excitation spectrum in Kelvin.

char *species

String to specify atom/ion species, e.g., 1H for Hydrogen, or 88SR+ for the $^{88}\text{Sr}^+$ ion.

int32_t n

Principal quantum number.

int32_t dn

Range for principal quantum number. At zero temperature, this argument is not used. For finite temperature, the absorption processes in Eq. (II.54) allow ‘decay’ to states with higher energy. In principle, one needs to take into account all transitions to these states, of which there are infinitely many. Here, we only take into account dn levels above Ψ_{n,l,s,j,m_j} to truncate the series, for both $l' = l \pm 1$. In other words, the sum in Eq. (II.54), taking into account absorption, contains $2 \times \text{dn}$ terms for $l > 0$ and dn for $l = 0$.

int32_t l

Orbital angular momentum quantum number $l = 0, 1, \dots, n - 1$.

double s

Electron spin quantum number. Not an argument, as $s = 1/2$ is always assumed.

double j

Total angular momentum quantum number $j = |l - 1/2|, l + 1/2$.

RETURN

double tau

τ is the lifetime of the fine-structure state Ψ_{n,l,s,j,m_j} (independent of m_j) in nanoseconds.

References

- [1] M. Aymar, C. H. Greene, and E. Luc-Koenig, ‘Multichannel Rydberg spectroscopy of complex atoms’, *Rev. Mod. Phys.* **68**, 1015 (1996).
- [2] M. Marinescu, H. R. Sadeghpour, and A. Dalgarno, ‘Dispersion coefficients for alkali-metal dimers’, *Phys. Rev. A* **49**, 982 (1994).
- [3] E. U. Condon and S. G. H., *The Theory of Atomic Spectra* (Cambridge University Press, Cambridge, 1951) 2nd ed.
- [4] J. W. P. Wilkinson, K. Bolsmann, T. L. M. Guedes, M. Müller, and I. Lesanovsky, ‘Two-qubit gate protocols with microwave-dressed Rydberg ions in a linear Paul trap’, *New. J. Phys.* **27**, 064502 (2025).
- [5] N. Šibalić, J. D. Pritchard, C. S. Adams, and K. J. Weatherill, ‘ARC: An open-source library for calculating properties of alkali Rydberg atoms’, *Comput. Phys. Commun.* **220**, 319–331 (2017).
- [6] H. Friedrich, *Theoretical Atomic Physics*, K. H. Becker, J.-M. Di Meglio, S. Hassani, B. Munro, R. Needs, W. T. Rhodes, S. Scott, H. E. Stanley, M. Stutzmann, and A. Wipf (Springer International Publishing AG, Heidelberg, 2017) 4th ed. ISBN: 978-3-319-47769-5.
- [7] R. L. Burden and J. D. Faires, *Numerical Analysis* (Cengage Learning, Boston, MA, 2011) 9th ed. ISBN: 978-0-538-73351-9 pp. 1–888.
- [8] D. A. Varshalovich, A. N. Moskalev, and V. K. Khersonskii, *Quantum Theory of Angular Momentum* (WORLD SCIENTIFIC, Leningrad, 1988) ISBN: 9789971509965.
- [9] A. Einstein, ‘Zur Quantentheorie der Strahlung’, *Physik. Zeitschr.* **XVIII**, 121–128 (1917).