

The dynamics of adaptive neuronal networks: influence of topology on synchronisation

Simon Aertssen, s181603

Supervisors

Erik Martens and Poul Hjorth

February 1st 2021

DTU Compute

Department of Applied Mathematics and Computer Science



Contents

1. Introduction
2. The Theta Neuron Model
3. Network Topologies
4. Mean Field Reductions
5. *Investigation*: Mean Field Reductions for undirected graphs
6. Hebbian Learning and Synaptic Plasticity
7. *Investigation*: Emerging Network Topologies
8. Conclusion and Discussion

Neuronal activity


- Neurons receive neurotransmitters
- Stimuli control the neurons' membrane potential
- Action potential = explosion of electrical activity
- Synapse releases the neurons' neurotransmitter

The Theta Neuron Model


Model formulation

$$\dot{\theta} = (1 - \cos \theta) + (1 + \cos \theta) \cdot I \quad \theta \in \mathbb{T} \quad (1)$$


- SNIC bifurcation



Excitable regime: $I < 0$






Bifurcation: $I = 0$



Periodic regime: $I > 0$

The Theta Neuron Model



Network Topologies

Network Topologies

Mean Field Reductions

Mean Field Reductions

Investigation: Mean Field Reductions for undirected graphs

***Investigation:* Mean Field Reductions for undirected graphs**

Hebbian Learning and Synaptic Plasticity

Hebbian Learning and Synaptic Plasticity

Investigation: Emerging Network Topologies

Investigation: Emerging Network Topologies

Conclusion and Discussion

Conclusion and Discussion