

Maximum Principle for Absolute Minimisers

Simone Carano and Nikos Katzourakis

March 23, 2026

Abstract

2020 MSC:

Key words and phrases: Vectorial Calculus of Variations in L^∞ ; higher order problems; Absolute Minimisers, Maximum Principle.

1 Introduction

For $n, N, k \in \mathbb{N}$, let $\Omega \subset \mathbb{R}^n$ be an open bounded set and $H : \Omega \times \mathbb{R}^N \times \mathbb{R}^{Nn} \times \dots \times \mathbb{R}^{Nn^k} \rightarrow \mathbb{R}$ be $\mathcal{L}(\Omega) \otimes \mathcal{B}(\mathbb{R}^\alpha)$ -measurable, where we have identified $\mathbb{R}^\alpha \cong \mathbb{R}^N \times \mathbb{R}^{Nn} \times \dots \times \mathbb{R}^{Nn^k}$ for $\alpha = \alpha(n, N, k) = N \frac{n^{k+1}-1}{n-1}$. Here $\mathcal{L}(\Omega)$ is the Lebesgue σ -algebra of Ω and $\mathcal{B}(\mathbb{R}^\alpha)$ is the Borel σ -algebra of $\mathbb{R}^\alpha$. Let $U \subseteq \Omega$ be an open subset and consider the supremal functional

$$E_\infty(u, U) := \operatorname{ess\,sup}_{x \in U} H(x, D^{[k]}u(x)), \quad (1.1)$$

where $u \in W^{k,\infty}(\Omega, \mathbb{R}^N)$ and $D^{[k]}u := (u, Du, \dots, D^k u)$ is the jet of order k of u .

The goal herein is to prove a maximum principle property for the absolute minimisers of the functional E_∞ . We recall that $u \in W^{k,\infty}(\Omega, \mathbb{R}^N)$ is an absolute minimiser of E_∞ if

$$E_\infty(u, U) \leq E_\infty(u + \Phi, U) \quad \forall \Phi \in W_0^{k,\infty}(U, \mathbb{R}^N) \quad \forall U \subseteq \Omega \text{ open}.$$

Under very mild assumptions on the supremand H , we prove that for an absolute minimiser the essential supremum in (1.1) is “attained at the boundary of U ”. In order to give a precise meaning to this expression, we refer to the supremum on ∂U of the essential limsup of $H(\cdot, D^{[k]}u)$, which is defined as

$$H(x, D^{[k]}u(x))^* := \lim_{\varepsilon \rightarrow 0} \operatorname{ess\,sup}_{B_\varepsilon(x) \cap U} H(\cdot, D^{[k]}u) \quad \forall x \in \partial U. \quad (1.2)$$

Therefore, we give the following definition.

Definition 1.1 (Maximum principle). *We say that $u \in W^{k,\infty}(\Omega, \mathbb{R}^N)$ satisfies the maximum principle for E_∞ in Ω if*

$$\operatorname{ess\,sup}_U H(\cdot, D^{[k]}u) = \sup_{\partial U} H(\cdot, D^{[k]}u)^* \quad \forall U \subseteq \Omega \text{ open}. \quad (1.3)$$

We prove that this property is necessarily satisfied by absolute minimisers. Although it is not sufficient: the cone function $u(x) = |x|$ belongs to $W^{1,\infty}(\mathbb{R}^n)$ and clearly satisfies the maximum principle for $E_\infty(u, \Omega) := \operatorname{ess\,sup}_\Omega |Du|$, but it is not an absolute minimiser. However, we will see

that the maximum principle is a property that characterises a class of maps which is larger than the one of absolute minimisers, and are the ones that minimise E_∞ with respect to compactly supported variations.

Before stating our main results, we underline that E_∞ in (1.1) is the general object of study in the L^∞ -Calculus of Variations, a field initiated by Aronsson in [3]. His pioneering work on the scalar first order case (namely when $N = k = 1$) has been well developed by now, and most challenges have been thoroughly analysed and understood (see [19] for a survey reference). More recently, in [17], the second author approached the vectorial case ($N > 1$), which turned out to be more involved and intriguing than the scalar one. For this reason, a complete theory is still far from reach. The situation is very similar in the higher order case ($k > 1$), recently started in [22, 27]. Without any pretension of being exhaustive, we refer also to [17, 20, 21, 23] for a glimpse of the literature on vectorial first order problems and to [9, 13, 24–26] for higher order ones. Furthermore, the fractional order case ($k \notin \mathbb{N}$) has been recently explored in [10].

It is worth to mention that, given boundary data $u_0 \in W^{k,\infty}(\Omega, \mathbb{R}^N)$, the existence (and/or uniqueness) of absolute minimisers to the Dirichlet problem in $W_{u_0}^{k,\infty}(\Omega, \mathbb{R}^N)$ associated to E_∞ in (1.1) is an open problem already when $k = 1$ and $n, N > 1$ or when $N = 1$ and $n, k > 1$. There are some exceptions in the second order case, whenever H has special dependence in the second derivatives of u . We refer to [22, 25, 27] for the scalar second order case and to [9] for the vectorial one (see also [10] for the fractional case).

Now we proceed to specify the assumptions on the function H . By denoting $X = (X_0, X_1, \dots, X_k) \in \mathbb{R}^N \times \mathbb{R}^{Nn} \times \dots \times \mathbb{R}^{Nn^k} = \mathbb{R}^\alpha$, we assume that $H : \Omega \times \mathbb{R}^\alpha \rightarrow \mathbb{R}$ is Carathéodory and radially non-decreasing w.r.t. X , i.e. for almost every $x \in \Omega$ and every $X \in \mathbb{R}^\alpha$ the function

$$t \mapsto H(x, tX) \text{ is non-decreasing in } \mathbb{R}^+.$$

Notice that radial monotonicity implies that, for almost every $x \in \Omega$, $H(x, \cdot)$ has a global minimum at $X = 0$. Without loss of generality we can assume that $H(x, 0) = 0$ a.e. $x \in \Omega$, so that H takes values in $\mathbb{R}^+$ and we can write $E_\infty(u, U) = \|H(\cdot, D^{[k]}u)\|_{L^\infty(U)}$ for every $U \subseteq \Omega$ open.

We point out that any level convex function with global minimum at the origin is necessarily radially non-decreasing, so that our result applies to the class of level convex functions, which is a common assumption for H in L^∞ variational problems (see e.g. [4]). Moreover, one can even apply the result to non-continuous supremands: indeed, for a function $g : \mathbb{R}^+ \rightarrow \mathbb{R}^+$ lower semicontinuous and non-decreasing, set $G = g \circ H$ and $E'_\infty(u, U) := \text{ess sup}_U G(\cdot, D^{[k]}u)$. It is not difficult to see that E_∞ and E'_∞ share the same minimisers. In addition, if $u \in W^{k,\infty}(\Omega, \mathbb{R}^N)$ satisfies the maximum principle for E_∞ in Ω , then for every $U \subseteq \Omega$ open

$$E'_\infty(u, U) = g \left(\text{ess sup}_U H(\cdot, D^{[k]}u) \right) = g \left(\text{ess sup}_{\partial U} H(\cdot, D^{[k]}u)^* \right) = \left(\text{ess sup}_{\partial U} G(\cdot, D^{[k]}u)^* \right),$$

i.e. u satisfies the maximum principle for E'_∞ as well.

The second assumption on H concerns the uniformity w.r.t. x of its modulus continuity: since $H(x, \cdot)$ is continuous in $\mathbb{R}^\alpha$, it is uniformly continuous on every closed ball $\overline{B}_R(0) \subset \mathbb{R}^\alpha$. Thus, we require that the modulus of continuity on $B_R(0)$ is independent of x . More precisely, we assume

$$\begin{aligned} \forall R > 0 \exists \omega_R \in C(\mathbb{R}^+) \text{ non-decreasing, with } \omega_R(0) = 0 : \\ |H(x, X) - H(x, X')| \leq \omega_R(|X - X'|), \quad \text{a.e. } x \in \Omega, \quad \forall X \in \overline{B}_R(0). \end{aligned} \quad (1.4)$$

Now we can state our main result.

Theorem 1.2 (Maximum Principle for Absolute Minimisers). *Let $\Omega \subset \mathbb{R}^n$ be a bounded open set and let $H : \Omega \times \mathbb{R}^\alpha \rightarrow \mathbb{R}^+$ be a Carathéodory function satisfying (1.4), with $H(x, 0) = 0$ and $H(x, \cdot)$*

radially non-decreasing for a.e. $x \in \Omega$. Assume that $u \in W^{k,\infty}(\Omega, \mathbb{R}^N)$ is an absolute minimiser for the functional E_∞ , defined in (1.1). Then u satisfies the maximum principle for E_∞ in Ω , namely

$$\operatorname{ess\,sup}_U H(\cdot, D^{[k]}u) = \sup_{\partial U} H(\cdot, D^{[k]}u)^* \quad \forall U \subseteq \Omega \text{ open.} \quad (1.5)$$

where $H(\cdot, D^{[k]}u)^*$ is the essential limsup of $H(\cdot, D^{[k]}u)$, defined in (1.2).

Before stating our second theorem, we give the definition of absolute minimiser w.r.t. compactly supported variations.

Definition 1.3 (The class $\operatorname{AM}_c(\Omega, \mathbb{R}^N)$). *We say that $u \in W^{k,\infty}(\Omega, \mathbb{R}^N)$ is an absolute minimiser w.r.t. compactly supported variations, and we write $u \in \operatorname{AM}_c(\Omega, \mathbb{R}^N)$, if*

$$E_\infty(u, U) \leq E_\infty(u + \Phi, U) \quad \forall \Phi \in W_c^{k,\infty}(\Omega, \mathbb{R}^N) \quad \forall U \subseteq \Omega \text{ open.}$$

Our second result is the following.

Theorem 1.4. *Let $\Omega \subset \mathbb{R}^n$ be a bounded open set and let $H : \Omega \times \mathbb{R}^\alpha \rightarrow \mathbb{R}^+$ be a Carathéodory function with $H(x, 0) = 0$ and $H(x, \cdot)$ radially non-decreasing for a.e. $x \in \Omega$. Consider the functional E_∞ , defined in (1.1). Then, the following are equivalent for a map $u \in W^{k,\infty}(\Omega, \mathbb{R}^N)$:*

- i) u satisfies the maximum principle for E_∞ in Ω ;
- ii) $u \in \operatorname{AM}_c(\Omega, \mathbb{R}^N)$.

The proof of Theorems 1.2 and 1.4 are in Sections, after a short preliminary section.

2 Preparatory tools

2.1 On radial monotonicity

In this first preliminary section, we recall some notions of radial monotonicity for function on $\mathbb{R}^\alpha$, together with some relevant examples.

Definition 2.1. *A function $H : \mathbb{R}^\alpha \rightarrow \mathbb{R}$ is said to be*

- i) *radially non-decreasing if for every $X \in \mathbb{R}^\alpha$ the function $t \mapsto H(tX)$ is non-decreasing in $\mathbb{R}^+$;*
- ii) *strictly radially increasing if for every $X \in \mathbb{R}^\alpha$ the function $t \mapsto H(tX)$ is strictly increasing in $\mathbb{R}^+$;*
- iii) *strongly radially increasing if there exists a continuous function $c : [0, 1] \times \mathbb{R}^+ \rightarrow \mathbb{R}^+$, with $c(1, \cdot) = c(\cdot, 0) = 0$ and $c > 0$ in $(0, 1) \times (0, \infty)$, such that*

$$H(tX) \leq H(X) - c(t, |X|) \quad \forall t \in [0, 1] \quad \forall X \in \mathbb{R}^\alpha.$$

Notice that $i) \Rightarrow ii) \Rightarrow iii)$, and implications do not invert in general. The above definition is in complete analogy with the notion of level convexity and its stronger versions. In particular, any level convex (resp. strictly level convex) function with global minimum at the origin is radially non-decreasing (resp. strictly radially increasing). Of course, a radially non-decreasing function need not to be level convex, since its level sets only need to be star shaped w.r.t. the origin.

Strongly radially increasing functions can be seen as strict radially increasing functions with a modulus of monotonicity. This class of functions will be particularly important for us, since in the proof of Theorem 1.2 we will approximate radially non-decreasing functions with strongly radially increasing ones in the uniform convergence.

Example 2.2. Relevant examples of strongly radially increasing functions are:

- Any strongly convex function with global minimum at the origin;
- The cone function associated to a (smooth) open set $S \subset \mathbb{R}^n$ star shaped w.r.t. the origin with $\nu_S(X) \cdot X > 0$ for every $X \in \partial S$, where ν_S is the outward normal vector to ∂S .
- Functions of the form $H_{\lambda,\beta}(X) = H(X) + \lambda|X|^\beta$, with $\lambda, \beta > 0$ and H radially non-decreasing.

2.2 Properties of the essential limsup

In this second preliminary section, we give the definition of essential limsup, together with some properties.

Definition 2.3 (Essential limsup). *Let $U \subseteq \mathbb{R}^n$ be a Borel set and $f \in L^\infty(U)$. We say that the function $f^* \in L^\infty(U)$, defined as*

$$f^*(x) := \lim_{\varepsilon \rightarrow 0} \operatorname{ess\,sup}_{B_\varepsilon(x) \cap U} f \quad \forall x \in U,$$

is the essential limsup of f .

From [18, Proposition 9], we have that $f \leq f^*$ almost everywhere and f^* is upper semicontinuous on U . In general, $f \neq f^*$ on a set of positive measure: for instance, consider $f = \mathbb{1}_{\mathbb{R}^n \setminus K}$ for a nowhere dense compact set K , with $\mathcal{L}^n(K) > 0$. Moreover, again by [18, Proposition 9], the definition of f^* allows us to give a pointwise meaning to the essential supremum, indeed for any Borel set $U \subseteq \mathbb{R}^n$ we have

$$\sup_U f^* = \operatorname{ess\,sup}_U f. \quad (2.1)$$

For a set $E \subset \mathbb{R}^n$ and $\rho > 0$, we denote by E^ρ the open ρ -neighbourhood of E , namely

$$E^\rho := \{x \in \mathbb{R}^n : \operatorname{dist}(x, \partial E) < \rho\}.$$

The following lemma is in order.

Lemma 2.4. *Let $U \subset \mathbb{R}^n$ be an open bounded set. Then for any $K \subseteq \partial U$ compact subset, we have*

$$\sup_K f^* = \lim_{\rho \rightarrow 0} \operatorname{ess\,sup}_{K^\rho \cap U} f.$$

Proof. We start with noticing that for every $x \in K$ there exists $\rho > 0$ such that $B_\rho(x) \cap U \subset K^\rho \cap U$, so that

$$\operatorname{ess\,sup}_{B_\rho(x) \cap U} f \leq \operatorname{ess\,sup}_{K^\rho \cap U} f.$$

By passing to the limit as $\rho \rightarrow 0$ in both sides and taking the supremum on K in the left hand side, we obtain

$$\sup_K f^* \leq \lim_{\rho \rightarrow 0} \operatorname{ess\,sup}_{K^\rho \cap U} f.$$

It remains to prove the opposite inequality. By definition (2.3), for any fixed $x \in K$ and $\sigma > 0$, there exists $\rho_{x,\sigma} > 0$ such that

$$f^*(x) \geq \operatorname{ess\,sup}_{B_{\rho_{x,\sigma}}(x) \cap U} f - \sigma. \quad (2.2)$$

The family of balls $\{B_{\rho_{x,\sigma}}(x); x \in K\}$ is an open covering of K . Thus, we can extract a finite subcovering $\{B_{\rho_{x_i,\sigma}}(x_i); x_i \in K\}_{i=1,\dots,m}$. For seek of brevity, let us denote $B_{i,\sigma} = B_{\rho_{x_i,\sigma}}(x_i)$.

We claim that there exists $\rho_\sigma > 0$ such that $K^{\rho_\sigma} \subset \bigcup_{i=1}^m B_{i,\sigma}$.

By contradiction, suppose that for every $j \in \mathbb{N}$ there exists $y_j \in K^{1/j} \setminus \bigcup_{i=1}^m B_{i,\sigma}$. This is equivalent to say that

$$\forall j \in \mathbb{N} \quad \exists y_j \in \mathbb{R}^n : \begin{cases} y_j \notin B_{i,\sigma} & \forall i = 1, \dots, m, \\ \text{dist}(y_j, K) \leq 1/j. \end{cases} \quad (2.3)$$

Let $\bar{y}_j \in K$ be a point realising $d(y_j, K)$. Since K is compact, there exists a subsequence $(\bar{y}_{j_k}) \subset (\bar{y}_j)$ such that $\bar{y}_{j_k} \rightarrow \bar{x}$ as $k \rightarrow \infty$, for some $\bar{x} \in K$. Moreover, since $\text{dist}(y_{j_k}, \bar{y}_{j_k}) \leq 1/j_k$, we have also that $y_{j_k} \rightarrow \bar{x}$ as $k \rightarrow \infty$. But being $\{B_{i,\sigma}\}_{i=1,\dots,m}$ a covering of K , we must have that eventually $y_{j_k} \in B_{i,\sigma}$ for some $i \in \{1, \dots, m\}$, which contradicts (2.3).

Now, define $U_\sigma = \bigcup_{i=1}^m (B_{i,\sigma} \cap U)$ and notice that $K^\rho \cap U \subset U_\sigma$. Thus, using also (2.2), we have

$$\begin{aligned} \lim_{\rho \rightarrow 0} \text{ess sup}_{K^\rho \cap U} f &\leq \text{ess sup}_{K^{\rho_\sigma} \cap U} f \\ &\leq \text{ess sup}_{U_\sigma} f \\ &\leq \max_{i=1,\dots,m} \text{ess sup}_{B_{i,\sigma} \cap U} f \\ &\leq \max_{i=1,\dots,m} f^*(x_i) + \sigma \\ &\leq \sup_K f^* + \sigma. \end{aligned}$$

By letting $\sigma \rightarrow 0$, we conclude the proof. $\square$

3 Proof of the main results

This section is devoted to the proof of Theorems 1.2 and 1.4. The proof of the latter is basically a consequence of the argument in the proof of the former. Indeed, the key idea in the proof of Theorem 1.2 is to find, for every $U \subseteq \Omega$, a suitable competitor in $u_\lambda \in W_u^{k,\infty}(U, \mathbb{R}^N)$ to compare with the absolute minimiser u . Actually, it turns out that $u_\lambda \in u + W_c^{k,\infty}(U, \mathbb{R}^N)$, and this will be the crucial observation for the proof of Theorem 1.4.

We start with proving Theorem 1.2: the key argument is based on the case H is strongly radially increasing in X uniformly w.r.t. x . The result in the general case, i.e. when H is only radially non-decreasing in X , is recovered by an approximation argument.

Proof of Theorem 1.2:

Fix $U \subseteq \Omega$ an open subset. First, notice that for every $x \in \partial U$ and every $\rho > 0$, we clearly have

$$\text{ess sup}_{B_\rho(x) \cap U} H(\cdot, D^{[k]}u) \leq \text{ess sup}_U H(\cdot, D^{[k]}u).$$

From definition 2.3, by passing to the limit as $\rho \rightarrow 0$ and then to the supremum over all $x \in \partial U$ in the left hand side, we obtain

$$\sup_{\partial U} H(\cdot, D^{[k]}u)^* \leq \text{ess sup}_U H(\cdot, D^{[k]}u). \quad (3.1)$$

Let us prove the opposite inequality. We divide the proof in two steps: First, we treat the case when $H(x, \cdot)$ is strongly radially increasing, uniformly in x . Then, we will deal with the general case.

Step 1: $H(x, \cdot)$ strongly radially increasing on $\mathbb{R}^\alpha$, uniformly w.r.t $x \in \Omega$.

Assume that there exists a continuous function $c : [0, 1] \times \mathbb{R}^+ \rightarrow \mathbb{R}^+$, with $c(1, \cdot) = c(\cdot, 0) = 0$ and $c > 0$ in $(0, 1) \times (0, \infty)$, such that

$$H(x, tX) \leq H(x, X) - c(t, |X|) \quad \forall t \in [0, 1] \quad \forall X \in \mathbb{R}^\alpha \quad \text{a.e. } x \in \Omega. \quad (3.2)$$

We start by setting

$$M := \operatorname{ess\,sup}_U H(\cdot, D^{[k]}u). \quad (3.3)$$

If $M = 0$, there is nothing to prove, so we can assume that $M > 0$. For seek of contradiction, assume that (3.1) holds strict. Then, by applying Lemma 2.4 to $K = \partial U$, one can find $\delta > 0$ such that

$$\lim_{\rho \rightarrow 0} \operatorname{ess\,sup}_{U^\rho \cap U} H(\cdot, D^{[k]}u) + 2\delta \leq M.$$

In particular, there exists $\varepsilon > 0$ such that

$$\operatorname{ess\,sup}_{U^\varepsilon \cap U} H(\cdot, D^{[k]}u) + \delta \leq M. \quad (3.4)$$

By possibly reducing ε in size, we can choose a function $\varphi_\varepsilon \in C^\infty(\mathbb{R}^n)$, with $0 \leq \varphi_\varepsilon \leq 1$, such that $\varphi_\varepsilon = 0$ in $U \setminus U^\varepsilon$ and $\varphi_\varepsilon = 1$ in $U^{\varepsilon/2}$. Now, for $\lambda \in (0, 1)$, set

$$u_\lambda := \lambda u + (1 - \lambda)\varphi_\varepsilon u.$$

Notice that $u_\lambda \in W^{k, \infty}(U, \mathbb{R}^N)$. Moreover, $u_\lambda = u$ on $U^{\varepsilon/2} \cap U$, so that

$$u_\lambda \in u + W_c^{k, \infty}(U, \mathbb{R}^N) \subset W_u^{1, \infty}(U, \mathbb{R}^N). \quad (3.5)$$

Let us compute $D^{[k]}u_\lambda$. For every $h = 1, \dots, k$, we have

$$D^h u_\lambda = \lambda D^h u + (1 - \lambda) D^h(\varphi_\varepsilon u). \quad (3.6)$$

By the Leibniz formula,

$$\partial_{i_1, \dots, i_h}^h (\varphi_\varepsilon u) = \sum_{j=0}^h \binom{h}{j} (\partial_{i_{j+1}, \dots, i_h}^{h-j} \varphi_\varepsilon) (\partial_{i_1, \dots, i_j}^j u).$$

In particular, for every $h = 0, \dots, k$, we can estimate

$$|D^h(\varphi_\varepsilon u)| \leq 2^h \|\varphi_\varepsilon\|_{W^{h, \infty}(U)} \|u\|_{W^{h, \infty}(U, \mathbb{R}^N)} \quad \text{in } U,$$

which gives

$$|D^{[k]}(\varphi_\varepsilon u)| \leq 2^{k+1} \|\varphi_\varepsilon\|_{W^{k, \infty}(U)} \|u\|_{W^{k, \infty}(U, \mathbb{R}^N)} \quad \text{in } U. \quad (3.7)$$

Putting together (3.6) and (3.7), we obtain

$$|D^{[k]}u_\lambda - \lambda D^{[k]}u| \leq (1 - \lambda) 2^{k+1} \|\varphi_\varepsilon\|_{W^{k, \infty}(U)} \|u\|_{W^{k, \infty}(U, \mathbb{R}^N)} \quad \text{in } U. \quad (3.8)$$

In particular, if we set

$$C_1 = C_1(\varepsilon, k, \lambda) = (1 - \lambda) 2^{k+1} \|\varphi_\varepsilon\|_{W^{k, \infty}(U)} + \lambda,$$

then

$$|D^{[k]}u_\lambda| \leq C_1 \|u\|_{W^{k,\infty}(U, \mathbb{R}^N)} \quad \text{in } U.$$

Now, we recall assumption (1.4) and apply it to

$$R := C_1 \|u\|_{W^{k,\infty}(U, \mathbb{R}^N)}, \quad D^{[k]}u_\lambda(x), \lambda D^{[k]}u(x) \in \overline{\mathbb{B}}_R(0) \quad \text{for a.e. } x \in U,$$

obtaining the existence of a continuous function $\omega = \omega_R : \mathbb{R}^+ \rightarrow \mathbb{R}^+$, non-decreasing, with $\omega(0) = 0$, such that

$$|H(x, D^{[k]}u_\lambda(x)) - H(x, \lambda D^{[k]}u(x))| \leq \omega(|D^{[k]}u_\lambda(x) - \lambda D^{[k]}u(x)|) \quad \text{a.e. } x \in U.$$

From (3.8) and the monotonicity of ω , setting

$$C_2 = C_2(\varepsilon, k) = 2^{k+1} \|\varphi_\varepsilon\|_{W^{k,\infty}(U)} \|u\|_{W^{k,\infty}(U, \mathbb{R}^N)},$$

we get

$$|H(\cdot, D^{[k]}u_\lambda) - H(\cdot, \lambda D^{[k]}u)| \leq \omega((1-\lambda)C_2) \quad \text{a.e. in } U.$$

By restricting the previous estimate to $U^\varepsilon \cap U$ and passing to the essential supremum on this set, we have

$$\begin{aligned} \operatorname{ess\,sup}_{U^\varepsilon \cap U} H(\cdot, D^{[k]}u_\lambda) &\leq \operatorname{ess\,sup}_{U^\varepsilon \cap U} H(\cdot, \lambda D^{[k]}u) + \omega((1-\lambda)C_2) \\ &\leq M - \delta + \omega((1-\lambda)C_2), \end{aligned}$$

where in the last inequality we have applied (3.4). Now, for λ sufficiently close to 1, by continuity of ω , we have $\omega((1-\lambda)C_2) \leq \delta/2$, giving

$$\operatorname{ess\,sup}_{U^\varepsilon \cap U} H(\cdot, D^{[k]}u_\lambda) \leq M - \frac{\delta}{2} < M. \quad (3.9)$$

On the other hand, let us prove that

$$\operatorname{ess\,sup}_{U \setminus U^\varepsilon} H(\cdot, D^{[k]}u_\lambda) < M. \quad (3.10)$$

To this purpose, notice that, since $\varphi_\varepsilon = 0$ in $U \setminus U^\varepsilon$, we have $u_\lambda = \lambda u$ in $U \setminus U^\varepsilon$. Thus, set

$$\overline{M} := \operatorname{ess\,sup}_U H(\cdot, \lambda D^{[k]}u)$$

and aim at proving $\overline{M} < M$. Without loss of generality, we can assume $\overline{M} > 0$. So, by definition of essential supremum, for every $\delta \in (\overline{M}/2, \overline{M})$, there exists $x_\delta \in U$ such that

$$H(x_\delta, \lambda D^{[k]}u(x_\delta)) \geq \overline{M} - \delta. \quad (3.11)$$

On the other hand, by recalling assumption (3.2) and (3.3), we have

$$\begin{aligned} H(x_\delta, \lambda D^{[k]}u(x_\delta)) &\leq H(x_\delta, D^{[k]}u(x_\delta)) - c(\lambda, |D^{[k]}u(x_\delta)|) \\ &\leq M - c(\lambda, |D^{[k]}u(x_\delta)|). \end{aligned} \quad (3.12)$$

Now we claim that there exists $\sigma > 0$ independent of δ such that

$$c(\lambda, |D^{[k]}u(x_\delta)|) \geq \sigma. \quad (3.13)$$

Indeed, from (3.11) and the continuity of H at $X = 0$, we infer that $|D^{[k]}u(x_\delta)|$ is bounded away from 0. Moreover, since $u \in W^{k,\infty}(\Omega, \mathbb{R}^N)$, we have that $|D^{[k]}u(x_\delta)|$ is also bounded from above. In other words, there exist two constants $R \geq r > 0$, independent of δ , such that

$$r \leq |D^{[k]}u(x_\delta)| \leq R.$$

Hence, since $c > 0$ in $(0, 1) \times (0, \infty)$, by continuity we have

$$c(\lambda, |D^{[k]}u(x_\delta)|) \geq \min_{\rho \in [r, R]} c(\lambda, \rho) =: \sigma > 0.$$

Putting together (3.11), (3.12), and (3.13), we obtain

$$\overline{M} - \delta \leq M - c(\lambda, |D^{[k]}u(x_\delta)|) \leq M - \sigma,$$

for $\sigma > 0$ independent of δ . We conclude that $\overline{M} \leq M - \sigma < M$, by letting $\delta \rightarrow 0$. As a consequence, we deduce (3.10), indeed

$$\begin{aligned} \operatorname{ess\,sup}_{U \setminus U^\varepsilon} H(\cdot, D^{[k]}u_\lambda) &= \operatorname{ess\,sup}_{U \setminus U^\varepsilon} H(\cdot, \lambda D^{[k]}u) \\ &\leq \operatorname{ess\,sup}_U H(\cdot, \lambda D^{[k]}u) \\ &= \overline{M} < M \end{aligned}$$

Finally, by recalling (3.3), and the inequalities (3.9) and (3.10), we obtain

$$\operatorname{ess\,sup}_U H(\cdot, D^{[k]}u_\lambda) \leq \max \left\{ \operatorname{ess\,sup}_{U^\varepsilon \cap U} H(\cdot, D^{[k]}u_\lambda), \operatorname{ess\,sup}_{U \setminus U^\varepsilon} H(\cdot, D^{[k]}u_\lambda) \right\} < M.$$

This inequality and (3.5) contradict the absolute minimality of u .

Step 2: General case.

Assume that $H(x, \cdot)$ is radially non-decreasing for almost every $x \in \Omega$. For $\varepsilon > 0$, consider the function $H_\varepsilon : \Omega \times \mathbb{R}^\alpha \rightarrow \mathbb{R}^+$ defined as

$$H_\varepsilon(x, X) = H(x, X) + \varepsilon |X|^2 \quad \forall X \in \mathbb{R}^\alpha \quad \text{a.e. } x \in \Omega.$$

Then, H_ε is strongly radially increasing in X , uniformly w.r.t. x (see also Example 2.2): indeed, it is enough to take $c(t, \rho) := \varepsilon(1 - t^2)\rho^2$. Hence, by (3.1) and *Step 1*, for every open subset $U \subseteq \Omega$, we have

$$\operatorname{ess\,sup}_U H_\varepsilon(\cdot, D^{[k]}u) = \sup_{\partial U} H_\varepsilon(\cdot, D^{[k]}u)^*. \quad (3.14)$$

Moreover, for every compact subset $K \subset \mathbb{R}^\alpha$, $H_\varepsilon \rightarrow H$ uniformly in $\Omega \times K$ as $\varepsilon \rightarrow 0$. Therefore, recalling Lemma (2.4), we have

$$\begin{aligned} \sup_{\partial U} H_\varepsilon(\cdot, D^{[k]}u)^* &= \lim_{\rho \rightarrow 0} \operatorname{ess\,sup}_{(\partial U)^\rho \cap U} H_\varepsilon(\cdot, D^{[k]}u) \\ &\leq \inf_{\rho > 0} \operatorname{ess\,sup}_{(\partial U)^\rho \cap U} H_\varepsilon(\cdot, D^{[k]}u). \end{aligned}$$

In particular, for every $\rho > 0$,

$$\sup_{\partial U} H_\varepsilon(\cdot, D^{[k]}u)^* \leq \operatorname{ess\,sup}_{(\partial U)^\rho \cap U} H_\varepsilon(\cdot, D^{[k]}u).$$

Passing to the limit in both sides as $\varepsilon \rightarrow 0$, by uniform convergence of H_ε to H , we get

$$\limsup_{\varepsilon \rightarrow 0} \sup_{\partial U} H_\varepsilon(\cdot, D^{[k]}u)^* \leq \operatorname{ess\,sup}_{(\partial U)^\rho \cap U} H(\cdot, D^{[k]}u) \quad \forall \rho > 0.$$

Sending $\rho \rightarrow 0$ in the right hand side, using again Lemma (2.4), we obtain

$$\limsup_{\varepsilon \rightarrow 0} \sup_{\partial U} H_\varepsilon(\cdot, D^{[k]}u)^* \leq \sup_{\partial U} H(\cdot, D^{[k]}u)^*. \quad (3.15)$$

On the other hand, by recalling (3.1) and the uniform convergence of H_ε to H , we can write

$$\sup_{\partial U} H(\cdot, D^{[k]}u)^* \leq \operatorname{ess\,sup}_U H(\cdot, D^{[k]}u) = \lim_{\varepsilon \rightarrow 0} \operatorname{ess\,sup}_U H_\varepsilon(\cdot, D^{[k]}u). \quad (3.16)$$

Putting together (3.15) and (3.16), identity (3.14) allows to conclude

$$\operatorname{ess\,sup}_U H(\cdot, D^{[k]}u) = \sup_{\partial U} H(\cdot, D^{[k]}u)^*.$$

The proof is completed. □

4 On the gradient maximum principle for p -harmonic maps

METTERLA QUESTA SEZIONE??

ACKNOWLEDGEMENTS: S.C. and N.K. acknowledge partial financial support through the EP-SRC grant EP/X017109/1. S.C. acknowledges partial financial support through the EPSRC grant EP/X017206/1. S.C. is a member of Gruppo Nazionale per l'Analisi Matematica, la Probabilità e le loro Applicazioni (GNAMPA) of the Istituto Nazionale di Alta Matematica (INdAM) of Italy.

DATA AVAILABILITY: Our manuscript has no associated data.

CONFLICT OF INTEREST: The authors have no Conflict of interest to declare for this article.

References

- [1] L. Ambrosio, A. Carlotto, A. Massaccesi, *Lectures on Elliptic Partial Differential Equations*, Publications of the Scuola Normale Superiore (2019).
- [2] N. Ansini, F. Prinari, *On the lower semicontinuity of supremal functional under differential constraints*, ESAIM - Control, Opt. and Calc. Var., **21** (4), 1053–1075 (2015).
- [3] G. Aronsson, *Minimization problems for the functional $\sup_x F(x, f(x), f'(x))$ I, II, III*, Arkiv fur Mat. 33 - 53 (1965); 409 - 431 (1966); 509 - 512 (1969).
- [4] E.N. Barron, R.R. Jensen, C.Y. Wang, *The Euler equation and absolute minimizers of L^∞ functionals*, Arch. Rational Mech. Anal. **157**, 255-283 (2001).
- [5] V. I. Bogachev, N. V. Krylov, M. Röckner, *Elliptic and parabolic equations for measures*, Uspekhi Mat. Nauk **64**, 5–116 (2009), translation in Russian Math. Surveys **64**, 973–1078 (2009).
- [6] A. Braides, *A handbook of Γ -convergence*, Handbook of Differential Equations: Stationary Partial Differential Equations, North-Holland, **3**, 101-213 (2006).

- [7] C. Brizzi, L. De Pascale, *A property of Absolute Minimizers in L^∞ Calculus of Variations and of solutions of the Aronsson-Euler equation*, Advances in Differential Equations 28 (3/4), 287-310 (2023).
- [8] L. Bungert, Y. Korolev, *Eigenvalue problems in L^∞ : optimality conditions, duality, and relations with optimal transport*, Communications of the American Mathematical Society 2:345 (2022).
- [9] S. Carano, N. Katzourakis, R. Moser, *Existence, uniqueness and characterisation of vector-valued absolute minimisers for a second order L^∞ -variational problem*, Preprint arXiv <https://arxiv.org/abs/2504.04181> (2025).
- [10] S. Carano, R. Moser, *An L^∞ -variational problem involving the Fractional Laplacian*, Preprint arXiv <https://arxiv.org/abs/2510.14476> (2025).
- [11] B. Dacorogna, *Direct Methods in the Calculus of Variations*, 2nd ed., Appl. Math. Sci. 78, Springer, New York, 2008.
- [12] J. M. Danskin, *The theory of max-min with applications*, J. SIAM Appl. Math. **14**(4), 641-665 (1966).
- [13] B. Dutton, N. Katzourakis, *On Second-Order L^∞ Variational Problems with Lower-Order Terms*, DDGD..... (2026).
- [14] M. Giaquinta, L. Martinazzi, *An Introduction to the Regularity Theory for Elliptic Systems, Harmonic Maps and Minimal Graphs*, Publications of the Scuola Normale Superiore (2018).
- [15] D. G. de Figueiredo, J. P. Gossez, *Strict Monotonicity of Eigenvalues and Unique Continuation*, Communications in Partial Differential Equations **17**, 339–346 (1992).
- [16] R. Hardt, L. Simon, *Nodal sets for solutions of elliptic equations*, J. Differential Geom. **30** 505 - 522 (1989).
- [17] N. Katzourakis, *L^∞ Variational Problems for maps and the Aronsson PDE System*, J. Differential Equations **253**(7), 2123 - 2139 (2012).
- [18] N. Katzourakis, *An L^∞ regularisation strategy to the inverse source identification problem for elliptic equations*, SIAM.....
- [19] N. Katzourakis, *An Introduction to Viscosity Solutions for Fully Nonlinear PDE with Applications to Calculus of Variations in L^∞* , Springer Briefs in Mathematics, DOI 10.1007/978-3-319-12829-0 (2015).
- [20] N. Katzourakis, *∞ -minimal submanifolds*, Proc. Amer. Math. Soc. **142**, 2797-2811 (2014).
- [21] N. Katzourakis, *On the structure of ∞ -harmonic maps*, Comm. Partial Differential Equations **39**, 2091–2124 (2014).
- [22] N. Katzourakis, R. Moser, *Existence, uniqueness and structure of second order absolute minimisers*, ARMA **231**(3), 1615-1634 (2019).
- [23] N. Katzourakis, R. Moser, *Minimisers of supremal functionals and mass-minimising 1-currents*, Calc. Var. **64**(26) (2025).

- [24] N. Katzourakis, R. Moser, *Variational problems in L^∞ involving semilinear second order differential operators*, ESAIM: COCV, **29**(76) (2023).
- [25] N. Katzourakis, R. Moser, *Existence, uniqueness and characterisation of local minimisers in higher order Calculus of Variations in L^∞* , ArXiv Preprint, <https://arxiv.org/abs/2403.12625> (2024).
- [26] N. Katzourakis, T. Pryer, *On the numerical approximation of p -Biharmonic and ∞ -Biharmonic functions*, Numerical Methods for Partial Differential Equations, **35**(1), 155-180 (2018).
- [27] N. Katzourakis, T. Pryer, *2nd order L^∞ Variational Problems and the ∞ -Polylaplacian*, Advances in Calculus of Variations **13**, 115-140 (2020).
- [28] C. Kreisbeck, E. Zappale, *Lower semicontinuity and relaxation of nonlocal L^∞ -functionals*, Calculus of Variations and PDE, **59** (138), 1–36 (2020).
- [29] M. Manetti, *Topology*, Springer Cham, DOI 10.1007/978-3-319-16958-3 (2015).
- [30] Q. Miao, C. Wang, Y. Zhou, *Uniqueness of Absolute Minimizers for L^∞ -Functionals Involving Hamiltonians $H(x, p)$* , Archive for Rational Mechanics and Analysis **223** (1), 141–198 (2017).
- [31] R. Moser, *Structure and classification results for the ∞ -elastica problem*, American Journal of Mathematics **144**(5), 1299-1329 (2022).
- [32] R. Moser, H. Schwetlick, *Minimizers of a weighted maximum of the Gauss curvature*, Annals of Global Analysis and Geometry **41**, 199-207 (2012).
- [33] F. Peng, C. Wang, Y. Zhou, *Regularity of absolute minimizers for continuous convex Hamiltonians*, Journal of Differential Equations, Volume 274 (2021), 1115-1164.
- [34] F. Prinari, E. Zappale, *A Relaxation Result in the Vectorial Setting and Power Law Approximation for Supremal Functionals*, J Optim. Theory Appl., **186**, 412–452 (2020).
- [35] A. M. Ribeiro, E. Zappale, *Existence of minimisers for nonlevel convex functionals*, SIAM J. Control Opt., **52** (5), 3341–3370 (2014).
- [36] A. M. Ribeiro, E. Zappale, *Revisited convexity notions for L^∞ variational problems*, Revista Matemática Complutense, published online, <https://doi.org/10.1007/s13163-024-00499-0> (2024).
- [37] O. A. Oleĭnik, E. V. Radkevič, *On the analyticity of solutions of linear partial differential equations*, Math. USSR Sb. **19**(581) (1973).
- [38] H. Whitney, *Analytic Extensions of Differentiable Functions Defined in Closed Sets*, Transactions of the American Mathematical Society **36**(1), 63-89 (1934).

SIMONE CARANO
 Department of Mathematical Sciences, University of Bath
 Bath BA2 7AY, UK
 E-mail: sc3705@bath.ac.uk

NIKOS KATZOURAKIS
Department of Mathematics and Statistics, University of Reading
Whiteknights Campus, Pepper Lane, Reading RG6 6AX, UK
E-mail: n.katzourakis@reading.ac.uk

ROGER MOSER
Department of Mathematical Sciences, University of Bath
Bath BA2 7AY, UK
E-mail: r.moser@bath.ac.uk