

Energy Maximum Principle for Absolute Minimisers

Simone Carano, Nikos Katzourakis and Roger Moser

May 21, 2026

Abstract

2020 MSC:

Key words and phrases: Vectorial Calculus of Variations in L^∞ ; higher order problems; Absolute Minimisers, Maximum Principle.

1 Introduction

For $n, N, k \in \mathbb{N}$, let $\Omega \subset \mathbb{R}^n$ be an open bounded set and $H : \Omega \times \mathbb{R}^N \times \mathbb{R}^{Nn} \times \dots \times \mathbb{R}^{Nn^k} \rightarrow \mathbb{R}$ be $\mathcal{L}(\Omega) \otimes \mathcal{B}(\mathbb{R}^\alpha)$ -measurable, where we have identified $\mathbb{R}^\alpha \cong \mathbb{R}^N \times \mathbb{R}^{Nn} \times \dots \times \mathbb{R}^{Nn^k}$ for $\alpha = \alpha(n, N, k) = N \frac{n^{k+1}-1}{n-1}$. here $\mathcal{L}(\Omega)$ is the Lebesgue σ -algebra of Ω and $\mathcal{B}(\mathbb{R}^\alpha)$ is the Borel σ -algebra of $\mathbb{R}^\alpha$. Let $U \subseteq \Omega$ be an open subset and consider the supremal functional

$$E_\infty(u, U) := \operatorname{ess\,sup}_{x \in U} H(x, D^{[k]}u(x)), \quad (1.1)$$

where $u \in W^{k,\infty}(\Omega, \mathbb{R}^N)$ and $D^{[k]}u := (u, Du, \dots, D^k u)$ is the jet of order k of u .

The goal herein is to prove a maximum principle property for the absolute minimisers of the functional E_∞ . We recall that $u \in W^{k,\infty}(\Omega, \mathbb{R}^N)$ is an absolute minimiser of E_∞ if

$$E_\infty(u, U) \leq E_\infty(u + \Phi, U) \quad \forall \Phi \in W_0^{k,\infty}(U, \mathbb{R}^N) \quad \forall U \subseteq \Omega \text{ open.}$$

Under very mild assumptions on the supremand H , we prove that for an absolute minimiser the essential supremum in (1.1) is “attained at the boundary of U ”. In order to give a precise meaning to this expression, we refer to the supremum on ∂U of the *essential limsup* of $H(\cdot, D^{[k]}u)$, which is defined as

$$H(x, D^{[k]}u(x))^* := \lim_{\varepsilon \rightarrow 0} \operatorname{ess\,sup}_{B_\varepsilon(x) \cap U} H(\cdot, D^{[k]}u) \quad \forall x \in \partial U. \quad (1.2)$$

Therefore, we give the following definition.

Definition 1.1 (Energy maximum principle). *We say that $u \in W^{k,\infty}(\Omega, \mathbb{R}^N)$ satisfies the energy maximum principle for E_∞ in Ω if*

$$\operatorname{ess\,sup}_U H(\cdot, D^{[k]}u) = \sup_{\partial U} H(\cdot, D^{[k]}u)^* \quad \forall U \subseteq \Omega \text{ open.} \quad (1.3)$$

If $u \in C^k(\overline{\Omega}, \mathbb{R}^N)$ and $H \in C(\Omega \times \mathbb{R}^\alpha)$, this can be rephrased as

$$\max_{\overline{U}} H(\cdot, D^{[k]}u) = \max_{\partial U} H(\cdot, D^{[k]}u) \quad \forall U \subseteq \Omega \text{ open.}$$

We prove that this property is necessarily satisfied by absolute minimisers. Although it is not sufficient: the cone function $u(x) = |x|$ belongs to $W^{1,\infty}(\mathbb{R}^n)$ and clearly satisfies the energy (or, equivalently, gradient) maximum principle for $E_\infty(u, \Omega) := \text{ess sup}_\Omega |Du|$, but it is not an absolute minimiser. However, we will see that the energy maximum principle is a property that characterises a class of maps which is larger than the one of absolute minimisers, and are the ones that minimise $E_\infty(\cdot, U)$ with respect to compactly supported variations in U , for every $U \subseteq \Omega$.

Before stating our main results, we underline that E_∞ in (1.1) is the general object of study in the L^∞ -Calculus of Variations, a field initiated by Aronsson in [1]. his pioneering work on the scalar first order case (namely when $N = k = 1$) has been well developed by now, and most challenges have been thoroughly analysed and understood (see [13] for a survey reference). More recently, the vectorial case ($N > 1$) and the higher order case ($k > 1$) have been approached respectively in [11] and [16, 21]. In these contexts, a complete theory is still far from reach. Without any pretension of being exhaustive, we refer also to [14, 15, 17] for a glimpse of the literature on vectorial first order problems and to [6, 8, 18–20] for higher order ones. Furthermore, the fractional order case ($k \notin \mathbb{N}$) has been recently explored in [7].

It is worth mentioning that, given boundary data $u_0 \in W^{k,\infty}(\Omega, \mathbb{R}^N)$, the existence (and/or uniqueness) of absolute minimisers to the Dirichlet problem in $W_{u_0}^{k,\infty}(\Omega, \mathbb{R}^N)$ associated to E_∞ in (1.1) is an open problem already when $k = 1$ and $n, N > 1$ or when $N = 1$ and $n, k > 1$. There are some exceptions in the second order case, whenever H has special dependence in the second derivatives of u : relevant works are [16, 19, 21] for the scalar second order case and [6] for the vectorial one; surprisingly, similar results hold true also in the fractional case [7].

Now we proceed to specify the assumptions on the function H . By denoting $X = (X_0, X_1, \dots, X_k) \in \mathbb{R}^N \times \mathbb{R}^{Nn} \times \dots \times \mathbb{R}^{Nn^k} = \mathbb{R}^\alpha$, we assume that $H : \Omega \times \mathbb{R}^\alpha \rightarrow \mathbb{R}$ is Carathéodory and radially non-decreasing w.r.t. X , i.e. for almost every $x \in \Omega$ and every $X \in \mathbb{R}^\alpha$ the function

$$t \mapsto H(x, tX) \text{ is non-decreasing in } \mathbb{R}^+.$$

Notice that radial monotonicity implies that, for almost every $x \in \Omega$, $H(x, \cdot)$ has a global minimum at $X = 0$. Without loss of generality we can assume that $H(x, 0) = 0$ a.e. $x \in \Omega$, so that H takes values in $\mathbb{R}^+$. In this case, $E_\infty(u, U) = \|H(\cdot, D^{[k]}u)\|_{L^\infty(U)}$ for every $U \subseteq \Omega$ open.

We point out that any level convex function with global minimum at the origin is necessarily radially non-decreasing, so that our result applies to the class of level convex functions, which is a common assumption for H in L^∞ variational problems (see e.g. [3]). Moreover, one can even apply the result to non-continuous supremands: indeed, for a function $g : \mathbb{R}^+ \rightarrow \mathbb{R}^+$ lower semicontinuous and non-decreasing, set $G = g \circ h$ and $E'_\infty(u, U) := \text{ess sup}_U G(\cdot, D^{[k]}u)$. It is not difficult to see that E_∞ and E'_∞ share the same minimisers. In addition, if $u \in W^{k,\infty}(\Omega, \mathbb{R}^N)$ satisfies the energy maximum principle for E_∞ in Ω , then for every $U \subseteq \Omega$ open

$$E'_\infty(u, U) = g \left(\text{ess sup}_U H(\cdot, D^{[k]}u) \right) = g \left(\text{ess sup}_{\partial U} H(\cdot, D^{[k]}u)^* \right) = \left(\text{ess sup}_{\partial U} G(\cdot, D^{[k]}u)^* \right),$$

i.e. u satisfies the energy maximum principle for E'_∞ as well.

The second assumption on H concerns the uniformity w.r.t. x of its modulus of continuity: since $H(x, \cdot)$ is continuous in $\mathbb{R}^\alpha$, it is uniformly continuous on every closed ball $\overline{\mathbb{B}}_R(0) \subset \mathbb{R}^\alpha$. Thus, we require that the modulus of continuity on $\overline{\mathbb{B}}_R(0)$ is independent of x . More precisely, we assume

$$\begin{aligned} \forall R > 0 \exists \omega_R \in C(\mathbb{R}^+) \text{ non-decreasing, with } \omega_R(0) = 0 : \\ |H(x, X) - H(x, X')| \leq \omega_R(|X - X'|), \quad \text{a.e. } x \in \Omega, \quad \forall X \in \overline{\mathbb{B}}_R(0). \end{aligned} \tag{1.4}$$

Now we can state our main result.

Theorem 1.2 (Energy maximum principle for absolute minimisers). *Let $\Omega \subset \mathbb{R}^n$ be a bounded open set and let $H : \Omega \times \mathbb{R}^\alpha \rightarrow \mathbb{R}^+$ be a Carathéodory function satisfying (1.4), with $H(x, 0) = 0$ and $H(x, \cdot)$ radially non-decreasing for a.e. $x \in \Omega$. Assume that $u \in W^{k,\infty}(\Omega, \mathbb{R}^N)$ is an absolute minimiser for the functional E_∞ , defined in (1.1). Then u satisfies the energy maximum principle for E_∞ in Ω , namely*

$$\operatorname{ess\,sup}_U H(\cdot, D^{[k]}u) = \sup_{\partial U} H(\cdot, D^{[k]}u)^* \quad \forall U \subseteq \Omega \text{ open.} \quad (1.5)$$

where $H(\cdot, D^{[k]}u)^*$ is the essential limsup of $H(\cdot, D^{[k]}u)$, defined in (1.2).

Another way of interpreting the energy maximum principle property is by saying that, for every $U \subseteq \Omega$, the attainment set $U(u)$ must contain the boundary of U . The attainment set $U(u)$ is the collection of all points $x \in \bar{U}$ where $H(x, D^{[k]}u(x)) = \operatorname{ess\,sup}_U H(\cdot, D^{[k]}u)$, and it is well defined and non-empty in case $u \in C^k(\Omega, \mathbb{R}^N)$ and $H \in C(\Omega \times \mathbb{R}^\alpha)$. For a definition of $U(u)$ in the non-smooth setting we refer to [5], where the authors show also a minimality property of $U(u)$ for absolute minimisers in the scalar first order case.

Before stating our second result, we give the definition of absolute minimiser w.r.t. compactly supported variations.

Definition 1.3 (The class $\operatorname{AM}_c(\Omega, \mathbb{R}^N)$). *We say that $u \in W^{k,\infty}(\Omega, \mathbb{R}^N)$ is an absolute minimiser w.r.t. compactly supported variations, and we write $u \in \operatorname{AM}_c(\Omega, \mathbb{R}^N)$, if*

$$E_\infty(u, U) \leq E_\infty(u + \Phi, U) \quad \forall \Phi \in W_c^{k,\infty}(\Omega, \mathbb{R}^N) \quad \forall U \subseteq \Omega \text{ open.}$$

As a consequence of Theorem 1.2, we obtain the following characterisation.

Corollary 1.4. *Let $\Omega \subset \mathbb{R}^n$ be a bounded open set and let $H : \Omega \times \mathbb{R}^\alpha \rightarrow \mathbb{R}^+$ be a Carathéodory function satisfying (1.4), with $H(x, 0) = 0$ and $H(x, \cdot)$ radially non-decreasing for a.e. $x \in \Omega$. Consider the functional E_∞ , defined in (1.1). Then, the following are equivalent for a map $u \in W^{k,\infty}(\Omega, \mathbb{R}^N)$:*

- i) u satisfies the energy maximum principle for E_∞ in Ω ;
- ii) $u \in \operatorname{AM}_c(\Omega, \mathbb{R}^N)$.

The proof of Theorem 1.2 and Corollary 1.4 are in Section 3, preceded by a short preparatory section. Therein we discuss the different notions of radial monotonicity which are of interest for us and some properties of the essential limsup.

Finally, in Section 4, we give a proof of a gradient maximum principle for p -harmonic maps. The methods used therein are genuinely different from the ones in the previous sections, and are based on the theory of elliptic PDEs. For the scalar case, results of this kind are already present in the literature, e.g. in [9, Chapter 9]. We refer also to [22, Appendix B] for a proof of gradient maximum and minimum principles for the class of p -harmonic potentials on convex rings in the plane. However, to the best of our knowledge, the validity of a gradient maximum principle for vector-valued p -harmonic maps appears to be new.

2 Preparatory tools

2.1 On radial monotonicity

In this first preliminary section, we recall some notions of radial monotonicity for function on $\mathbb{R}^\alpha$, together with some relevant examples.

Definition 2.1. A function $h : \mathbb{R}^\alpha \rightarrow \mathbb{R}$ is said to be

- i) *radially non-decreasing* if for every $X \in \mathbb{R}^\alpha$ the function $t \mapsto h(tX)$ is non-decreasing in $\mathbb{R}^+$;
- ii) *strictly radially increasing* if for every $X \in \mathbb{R}^\alpha$ the function $t \mapsto h(tX)$ is strictly increasing in $\mathbb{R}^+$;
- iii) *strongly radially increasing* if there exists a continuous function $c : [0, 1] \times \mathbb{R}^+ \rightarrow \mathbb{R}^+$, with $c(1, \cdot) = c(\cdot, 0) = 0$ and $c > 0$ in $(0, 1) \times (0, \infty)$, such that

$$h(tX) \leq h(X) - c(t, |X|) \quad \forall t \in [0, 1] \quad \forall X \in \mathbb{R}^\alpha.$$

Notice that $i) \Rightarrow ii) \Rightarrow iii)$, and implications do not invert in general. The above definition is in complete analogy with the notion of (level) convexity and its stronger versions. In particular, any level convex (resp. strictly level convex) function with global minimum at the origin is radially non-decreasing (resp. strictly radially increasing). Of course, a radially non-decreasing function need not to be level convex, since its level subsets only need to be star shaped w.r.t. the origin.

Strongly radially increasing functions can be seen as strict radially increasing functions with a modulus of monotonicity. This class of functions will be particularly important for us, since in the proof of Theorem 1.2 we will approximate radially non-decreasing functions with strongly radially increasing ones in the uniform convergence.

Example 2.2. Relevant examples of strongly radially increasing functions are:

- Any strongly convex function with global minimum at the origin;
- The cone function associated to a (smooth) open set $S \subset \mathbb{R}^\alpha$ star shaper w.r.t. the origin with $\nu_S(X) \cdot X > 0$ for every $X \in \partial S$, where ν_S is the outward normal vector to ∂S .
- Functions of the form $h_{\lambda, \beta}(X) = h(X) + \lambda|X|^\beta$, with $\lambda, \beta > 0$ and h radially non-decreasing.

2.2 Properties of the essential limsup

In this second preliminary section, we give the definition of essential limsup, together with some properties.

Definition 2.3 (Essential limsup). Let $U \subseteq \mathbb{R}^n$ be a Borel set and $f \in L^\infty(U)$. We say that the function $f^* \in L^\infty(U)$, defined as

$$f^*(x) := \lim_{\varepsilon \rightarrow 0} \operatorname{ess\,sup}_{B_\varepsilon(x) \cap U} f \quad \forall x \in U,$$

is the *essential limsup* of f .

Of course, if U is open and f is continuous in U , then $f^* = f$. However, in the more general setting of Definition 2.3, we may have $f \neq f^*$ on a set of positive measure: for instance, consider $f = \mathbb{1}_{\mathbb{R}^n \setminus K}$ for a nowhere dense compact set $K \subset \mathbb{R}^n$, with $\mathcal{L}^n(K) > 0$. In general, from [12, Proposition 9], we have that $f \leq f^*$ almost everywhere and f^* is upper semicontinuous on U . Moreover, again by [12, Proposition 9], the definition of f^* allows us to give a pointwise meaning to the essential supremum, indeed for any Borel set $U \subseteq \mathbb{R}^n$ we have

$$\sup_U f^* = \operatorname{ess\,sup}_U f. \tag{2.1}$$

For a set $E \subset \mathbb{R}^n$ and $\rho > 0$, we denote by E^ρ the open ρ -neighbourhood of E , namely

$$E^\rho := \{x \in \mathbb{R}^n : \operatorname{dist}(x, E) < \rho\}.$$

The following lemma is in order.

Lemma 2.4. *Let $U \subset \mathbb{R}^n$ be an open bounded set. Then for any $K \subseteq \partial U$ compact subset, we have*

$$\sup_K f^* = \lim_{\rho \rightarrow 0} \operatorname{ess\,sup}_{K^\rho \cap U} f.$$

Proof. We start with noticing that for every $x \in K$ there exists $\rho > 0$ such that $B_\rho(x) \cap U \subset K^\rho \cap U$, so that

$$\operatorname{ess\,sup}_{B_\rho(x) \cap U} f \leq \operatorname{ess\,sup}_{K^\rho \cap U} f.$$

By passing to the limit as $\rho \rightarrow 0$ in both sides and taking the supremum on K in the left hand side, we obtain

$$\sup_K f^* \leq \lim_{\rho \rightarrow 0} \operatorname{ess\,sup}_{K^\rho \cap U} f.$$

It remains to prove the opposite inequality. By definition (2.3), for any fixed $x \in K$ and $\sigma > 0$, there exists $\rho_{x,\sigma} > 0$ such that

$$f^*(x) \geq \operatorname{ess\,sup}_{B_{\rho_{x,\sigma}}(x) \cap U} f - \sigma. \quad (2.2)$$

The family of balls $\{B_{\rho_{x,\sigma}}(x); x \in K\}$ is an open covering of K . Thus, we can extract a finite subcovering $\{B_{\rho_{x_i,\sigma}}(x_i); x_i \in K\}_{i=1,\dots,m}$. For seek of brevity, let us denote $B_{i,\sigma} = B_{\rho_{x_i,\sigma}}(x_i)$.

We claim that there exists $\rho_\sigma > 0$ such that $K^{\rho_\sigma} \subset \bigcup_{i=1}^m B_{i,\sigma}$.

By contradiction, suppose that for every $j \in \mathbb{N}$ there exists $y_j \in K^{1/j} \setminus \bigcup_{i=1}^m B_{i,\sigma}$. This is equivalent to say that

$$\forall j \in \mathbb{N} \quad \exists y_j \in \mathbb{R}^n : \begin{cases} y_j \notin B_{i,\sigma} & \forall i = 1, \dots, m, \\ \operatorname{dist}(y_j, K) \leq 1/j. \end{cases} \quad (2.3)$$

Let $\bar{y}_j \in K$ be a point realising $d(y_j, K)$. Since K is compact, there exists a subsequence $(\bar{y}_{j_k}) \subset (\bar{y}_j)$ such that $\bar{y}_{j_k} \rightarrow \bar{x}$ as $k \rightarrow \infty$, for some $\bar{x} \in K$. Moreover, since $\operatorname{dist}(y_{j_k}, \bar{y}_{j_k}) \leq 1/j_k$, we have also that $y_{j_k} \rightarrow \bar{x}$ as $k \rightarrow \infty$. But being $\{B_{i,\sigma}\}_{i=1,\dots,m}$ a covering of K , we must have that eventually $y_{j_k} \in B_{i,\sigma}$ for some $i \in \{1, \dots, m\}$, which contradicts (2.3).

Now, define $U_\sigma = \bigcup_{i=1}^m (B_{i,\sigma} \cap U)$ and notice that $K^{\rho_\sigma} \cap U \subset U_\sigma$. Thus, using also (2.2), we have

$$\begin{aligned} \lim_{\rho \rightarrow 0} \operatorname{ess\,sup}_{K^\rho \cap U} f &\leq \operatorname{ess\,sup}_{K^{\rho_\sigma} \cap U} f \\ &\leq \operatorname{ess\,sup}_{U_\sigma} f \\ &\leq \max_{i=1,\dots,m} \operatorname{ess\,sup}_{B_{i,\sigma} \cap U} f \\ &\leq \max_{i=1,\dots,m} f^*(x_i) + \sigma \\ &\leq \sup_K f^* + \sigma. \end{aligned}$$

By letting $\sigma \rightarrow 0$, we conclude the proof. $\square$

3 Proof of the main results

This section is devoted to the proof of Theorems 1.2 and Corollary 1.4. The proof of the latter is basically a consequence of the argument in the proof of the former. Indeed, the key idea in the proof of Theorem 1.2 is to find, for every $U \subseteq \Omega$, a suitable competitor in $u_\lambda \in W_u^{k,\infty}(U, \mathbb{R}^N)$ to compare with the absolute minimiser u . Actually, it turns out that $u_\lambda \in u + W_c^{k,\infty}(U, \mathbb{R}^N)$, and

this will be the crucial observation in the proof of Corollary 1.4.

We start with proving Theorem 1.2: the key argument is based on the case $H(x, \cdot)$ strongly radially increasing on $\mathbb{R}^\alpha$, uniformly w.r.t $x \in \Omega$. The result in the general case, i.e. when $H(x, \cdot)$ is radially non-decreasing, is recovered by an approximation argument.

Proof of Theorem 1.2:

Fix $U \subseteq \Omega$ an open subset. First, notice that for every $x \in \partial U$ and every $\rho > 0$, we clearly have

$$\operatorname{ess\,sup}_{B_\rho(x) \cap U} H(\cdot, D^{[k]}u) \leq \operatorname{ess\,sup}_U H(\cdot, D^{[k]}u).$$

From definition 2.3, by passing to the limit as $\rho \rightarrow 0$ and then to the supremum over all $x \in \partial U$ in the left hand side, we obtain

$$\sup_{\partial U} H(\cdot, D^{[k]}u)^* \leq \operatorname{ess\,sup}_U H(\cdot, D^{[k]}u). \quad (3.1)$$

Let us prove the opposite inequality. We divide the proof in two steps: First, we treat the case when $H(x, \cdot)$ is strongly radially increasing, uniformly in x . Then, we will deal with the general case.

Step 1: $H(x, \cdot)$ strongly radially increasing on $\mathbb{R}^\alpha$, uniformly w.r.t $x \in \Omega$.

Assume that there exists a continuous function $c : [0, 1] \times \mathbb{R}^+ \rightarrow \mathbb{R}^+$, with $c(1, \cdot) = c(\cdot, 0) = 0$ and $c > 0$ in $(0, 1) \times (0, \infty)$, such that

$$H(x, tX) \leq H(x, X) - c(t, |X|) \quad \forall t \in [0, 1] \quad \forall X \in \mathbb{R}^\alpha \quad \text{a.e. } x \in \Omega. \quad (3.2)$$

We start by setting

$$M := \operatorname{ess\,sup}_U H(\cdot, D^{[k]}u). \quad (3.3)$$

If $M = 0$, there is nothing to prove, so we can assume that $M > 0$. For seek of contradiction, assume that (3.1) holds strict. Then, by applying Lemma 2.4 to $K = \partial U$, one can find $\delta > 0$ such that

$$\lim_{\rho \rightarrow 0} \operatorname{ess\,sup}_{(\partial U)^\rho \cap U} H(\cdot, D^{[k]}u) + 2\delta \leq M.$$

In particular, there exists $\varepsilon > 0$ such that

$$\operatorname{ess\,sup}_{(\partial U)^\varepsilon \cap U} H(\cdot, D^{[k]}u) + \delta \leq M. \quad (3.4)$$

By possibly reducing ε in size, we can choose a function $\varphi_\varepsilon \in C^\infty(\mathbb{R}^n)$, with $0 \leq \varphi_\varepsilon \leq 1$, such that $\varphi_\varepsilon = 0$ in $U \setminus (\partial U)^\varepsilon$ and $\varphi_\varepsilon = 1$ in $(\partial U)^{\varepsilon/2}$. Now, for $\lambda \in (0, 1)$, set

$$u_\lambda := \lambda u + (1 - \lambda)\varphi_\varepsilon u.$$

Notice that $u_\lambda \in W^{k, \infty}(U, \mathbb{R}^N)$. Moreover, $u_\lambda = u$ on $(\partial U)^{\varepsilon/2} \cap U$, so that

$$u_\lambda \in u + W_c^{k, \infty}(U, \mathbb{R}^N) \subset W_u^{1, \infty}(U, \mathbb{R}^N). \quad (3.5)$$

Let us compute $D^{[k]}u_\lambda$. For every $h = 1, \dots, k$, we have

$$D^h u_\lambda = \lambda D^h u + (1 - \lambda) D^h(\varphi_\varepsilon u). \quad (3.6)$$

By the Leibniz formula,

$$\partial_{i_1, \dots, i_h}^h (\varphi_\varepsilon u) = \sum_{j=0}^h \binom{h}{j} (\partial_{i_{j+1}, \dots, i_h}^{h-j} \varphi_\varepsilon) (\partial_{i_1, \dots, i_j}^j u).$$

In particular, for every $h = 0, \dots, k$, we can estimate

$$|\mathbf{D}^h(\varphi_\varepsilon u)| \leq 2^h \|\varphi_\varepsilon\|_{W^{h,\infty}(U)} \|u\|_{W^{h,\infty}(U, \mathbb{R}^N)} \quad \text{in } U,$$

which gives

$$|\mathbf{D}^{[k]}(\varphi_\varepsilon u)| \leq 2^{k+1} \|\varphi_\varepsilon\|_{W^{k,\infty}(U)} \|u\|_{W^{k,\infty}(U, \mathbb{R}^N)} \quad \text{in } U. \quad (3.7)$$

Putting together (3.6) and (3.7), we obtain

$$|\mathbf{D}^{[k]}u_\lambda - \lambda \mathbf{D}^{[k]}u| \leq (1 - \lambda) 2^{k+1} \|\varphi_\varepsilon\|_{W^{k,\infty}(U)} \|u\|_{W^{k,\infty}(U, \mathbb{R}^N)} \quad \text{in } U. \quad (3.8)$$

In particular, if we set

$$C_1 = C_1(\varepsilon, k, \lambda) = (1 - \lambda) 2^{k+1} \|\varphi_\varepsilon\|_{W^{k,\infty}(U)} + \lambda,$$

then

$$|\mathbf{D}^{[k]}u_\lambda| \leq C_1 \|u\|_{W^{k,\infty}(U, \mathbb{R}^N)} \quad \text{in } U.$$

Now, we recall assumption (1.4) and apply it to

$$R := C_1 \|u\|_{W^{k,\infty}(U, \mathbb{R}^N)}, \quad \mathbf{D}^{[k]}u_\lambda(x), \lambda \mathbf{D}^{[k]}u(x) \in \overline{\mathbb{B}}_R(0) \quad \text{for a.e. } x \in U,$$

obtaining the existence of a continuous function $\omega = \omega_R : \mathbb{R}^+ \rightarrow \mathbb{R}^+$, non-decreasing, with $\omega(0) = 0$, such that

$$|\mathbf{H}(x, \mathbf{D}^{[k]}u_\lambda(x)) - \mathbf{H}(x, \lambda \mathbf{D}^{[k]}u(x))| \leq \omega(|\mathbf{D}^{[k]}u_\lambda(x) - \lambda \mathbf{D}^{[k]}u(x)|) \quad \text{a.e. } x \in U.$$

From (3.8) and the monotonicity of ω , setting

$$C_2 = C_2(\varepsilon, k) = 2^{k+1} \|\varphi_\varepsilon\|_{W^{k,\infty}(U)} \|u\|_{W^{k,\infty}(U, \mathbb{R}^N)},$$

we get

$$|\mathbf{H}(\cdot, \mathbf{D}^{[k]}u_\lambda) - \mathbf{H}(\cdot, \lambda \mathbf{D}^{[k]}u)| \leq \omega((1 - \lambda)C_2) \quad \text{a.e. in } U.$$

By restricting the previous estimate to $(\partial U)^\varepsilon \cap U$ and passing to the essential supremum on this set, we have

$$\begin{aligned} \operatorname{ess\,sup}_{(\partial U)^\varepsilon \cap U} \mathbf{H}(\cdot, \mathbf{D}^{[k]}u_\lambda) &\leq \operatorname{ess\,sup}_{(\partial U)^\varepsilon \cap U} \mathbf{H}(\cdot, \lambda \mathbf{D}^{[k]}u) + \omega((1 - \lambda)C_2) \\ &\leq M - \delta + \omega((1 - \lambda)C_2), \end{aligned}$$

where in the last inequality we have applied (3.4). Now, for λ sufficiently close to 1, by continuity of ω , we have $\omega((1 - \lambda)C_2) \leq \delta/2$, giving

$$\operatorname{ess\,sup}_{(\partial U)^\varepsilon \cap U} \mathbf{H}(\cdot, \mathbf{D}^{[k]}u_\lambda) \leq M - \frac{\delta}{2} < M. \quad (3.9)$$

On the other hand, let us prove that

$$\operatorname{ess\,sup}_{U \setminus (\partial U)^\varepsilon} H(\cdot, D^{[k]}u_\lambda) < M. \quad (3.10)$$

To this purpose, notice that, since $\varphi_\varepsilon = 0$ in $U \setminus (\partial U)^\varepsilon$, we have $u_\lambda = \lambda u$ in $U \setminus (\partial U)^\varepsilon$. Thus, set

$$\overline{M} := \operatorname{ess\,sup}_U H(\cdot, \lambda D^{[k]}u)$$

and aim at proving $\overline{M} < M$. Without loss of generality, we can assume $\overline{M} > 0$. So, by definition of essential supremum, for every $\delta \in (0, \overline{M}/2)$, there exists $x_\delta \in U$ such that

$$H(x_\delta, \lambda D^{[k]}u(x_\delta)) \geq \overline{M} - \delta. \quad (3.11)$$

On the other hand, by recalling assumption (3.2) and (3.3), we have

$$\begin{aligned} H(x_\delta, \lambda D^{[k]}u(x_\delta)) &\leq H(x_\delta, D^{[k]}u(x_\delta)) - c(\lambda, |D^{[k]}u(x_\delta)|) \\ &\leq M - c(\lambda, |D^{[k]}u(x_\delta)|). \end{aligned} \quad (3.12)$$

Now we claim that there exists $\sigma > 0$ independent of δ such that

$$c(\lambda, |D^{[k]}u(x_\delta)|) \geq \sigma. \quad (3.13)$$

Indeed, from (3.11) and the continuity of H at $X = 0$, we infer that $|D^{[k]}u(x_\delta)|$ is bounded away from 0. Moreover, since $u \in W^{k,\infty}(\Omega, \mathbb{R}^N)$, we have that $|D^{[k]}u(x_\delta)|$ is also bounded from above. In other words, there exist two constants $R \geq r > 0$, independent of δ , such that

$$r \leq |D^{[k]}u(x_\delta)| \leq R.$$

Hence, since c is continuous and strictly positive in $(0, 1) \times (0, \infty)$, we have

$$c(\lambda, |D^{[k]}u(x_\delta)|) \geq \min_{\rho \in [r, R]} c(\lambda, \rho) =: \sigma > 0.$$

Putting together (3.11), (3.12), and (3.13), we obtain

$$\overline{M} - \delta \leq M - c(\lambda, |D^{[k]}u(x_\delta)|) \leq M - \sigma,$$

for $\sigma > 0$ independent of δ . We conclude that $\overline{M} \leq M - \sigma < M$, by letting $\delta \rightarrow 0$.

As a consequence, we deduce (3.10), indeed

$$\begin{aligned} \operatorname{ess\,sup}_{U \setminus (\partial U)^\varepsilon} H(\cdot, D^{[k]}u_\lambda) &= \operatorname{ess\,sup}_{U \setminus (\partial U)^\varepsilon} H(\cdot, \lambda D^{[k]}u) \\ &\leq \operatorname{ess\,sup}_U H(\cdot, \lambda D^{[k]}u) \\ &= \overline{M} < M \end{aligned}$$

Finally, by recalling (3.3), and the inequalities (3.9) and (3.10), we obtain

$$\operatorname{ess\,sup}_U H(\cdot, D^{[k]}u_\lambda) \leq \max \left\{ \operatorname{ess\,sup}_{(\partial U)^\varepsilon \cap U} H(\cdot, D^{[k]}u_\lambda), \operatorname{ess\,sup}_{U \setminus (\partial U)^\varepsilon} H(\cdot, D^{[k]}u_\lambda) \right\} < M.$$

This inequality and (3.5) contradict the absolute minimality of u .

Step 2: General case.

Assume that $H(x, \cdot)$ is radially non-decreasing for almost every $x \in \Omega$. For $\varepsilon > 0$, consider the function $H_\varepsilon : \Omega \times \mathbb{R}^\alpha \rightarrow \mathbb{R}^+$ defined as

$$H_\varepsilon(x, X) = H(x, X) + \varepsilon|X|^2 \quad \forall X \in \mathbb{R}^\alpha \quad \text{a.e. } x \in \Omega.$$

Then, H_ε is strongly radially increasing in X , uniformly w.r.t. x (see also Example 2.2): indeed, it is enough to take $c(t, \rho) := \varepsilon(1 - t^2)\rho^2$ as a modulus of monotonicity. Hence, by (3.1) and *Step 1*, for every open subset $U \subseteq \Omega$, we have

$$\operatorname{ess\,sup}_U H_\varepsilon(\cdot, D^{[k]}u) = \sup_{\partial U} H_\varepsilon(\cdot, D^{[k]}u)^*. \quad (3.14)$$

Moreover, for every compact subset $K \subset \mathbb{R}^\alpha$, $H_\varepsilon \rightarrow H$ uniformly in $\Omega \times K$ as $\varepsilon \rightarrow 0$. Therefore, recalling Lemma (2.4), we have

$$\begin{aligned} \sup_{\partial U} H_\varepsilon(\cdot, D^{[k]}u)^* &= \lim_{\rho \rightarrow 0} \operatorname{ess\,sup}_{(\partial U)^\rho \cap U} H_\varepsilon(\cdot, D^{[k]}u) \\ &= \inf_{\rho > 0} \operatorname{ess\,sup}_{(\partial U)^\rho \cap U} H_\varepsilon(\cdot, D^{[k]}u). \end{aligned}$$

In particular, for every $\rho > 0$,

$$\sup_{\partial U} H_\varepsilon(\cdot, D^{[k]}u)^* \leq \operatorname{ess\,sup}_{(\partial U)^\rho \cap U} H_\varepsilon(\cdot, D^{[k]}u).$$

Passing to the limit in both sides as $\varepsilon \rightarrow 0$, by uniform convergence of H_ε to H , we get

$$\lim_{\varepsilon \rightarrow 0} \sup_{\partial U} H_\varepsilon(\cdot, D^{[k]}u)^* \leq \operatorname{ess\,sup}_{(\partial U)^\rho \cap U} H(\cdot, D^{[k]}u) \quad \forall \rho > 0.$$

Sending $\rho \rightarrow 0$ in the right hand side, using again Lemma (2.4), we obtain

$$\lim_{\varepsilon \rightarrow 0} \sup_{\partial U} H_\varepsilon(\cdot, D^{[k]}u)^* \leq \sup_{\partial U} H(\cdot, D^{[k]}u)^*. \quad (3.15)$$

On the other hand, by recalling (3.1) and the uniform convergence of H_ε to H , we can write

$$\sup_{\partial U} H(\cdot, D^{[k]}u)^* \leq \operatorname{ess\,sup}_U H(\cdot, D^{[k]}u) = \lim_{\varepsilon \rightarrow 0} \operatorname{ess\,sup}_U H_\varepsilon(\cdot, D^{[k]}u). \quad (3.16)$$

Putting together (3.15) and (3.16), identity (3.14) allows to conclude

$$\operatorname{ess\,sup}_U H(\cdot, D^{[k]}u) = \sup_{\partial U} H(\cdot, D^{[k]}u)^*.$$

The proof is completed. □

A straightforward consequence is Corollary 1.4, whose proof is detailed below.

Proof of Corollary 1.4:

Let us show that satisfying the energy maximum principle implies the membership to the class $\operatorname{AM}_c(\Omega, \mathbb{R}^N)$. Fix $u \in W^{1,\infty}(\Omega, \mathbb{R}^N)$, an open set $U \subseteq \Omega$ and a map $\Phi \in W_c^{1,\infty}(U, \mathbb{R}^N)$. Let $U' \Subset U$ be an open subset with $\operatorname{supp}(\Phi) \Subset U'$. Then we have

$$\begin{aligned} E_\infty(u + \Phi, U) &= \operatorname{ess\,sup}_U H(\cdot, D^{[k]}(u + \Phi)) \\ &\geq \operatorname{ess\,sup}_{U \setminus U'} H(\cdot, D^{[k]}(u + \Phi)) \\ &= \operatorname{ess\,sup}_{U \setminus U'} H(\cdot, D^{[k]}u) \end{aligned} \quad (3.17)$$

Notice that $(\partial U)^\rho \cap U \subset U \setminus U'$ for a sufficiently small $\rho > 0$. hence

$$\operatorname{ess\,sup}_{U \setminus U'} H(\cdot, D^{[k]}u) \geq \operatorname{ess\,sup}_{(\partial U)^\rho \cap U} H(\cdot, D^{[k]}u).$$

By passing to the limit as $\rho \rightarrow 0$ in the right hand side and invoking Lemma 2.4, we get

$$\operatorname{ess\,sup}_{U \setminus U'} H(\cdot, D^{[k]}u) \geq \lim_{\rho \rightarrow 0} \operatorname{ess\,sup}_{(\partial U)^\rho \cap U} H(\cdot, D^{[k]}u) = \sup_{\partial U} H(\cdot, D^{[k]}u)^*. \quad (3.18)$$

Thus, if u satisfies the energy maximum principle in (1.5), by putting together (3.17) and (3.18), we get

$$\begin{aligned} E_\infty(u + \Phi, U) &\geq \sup_{\partial U} H(\cdot, D^{[k]}u)^* \\ &= \operatorname{ess\,sup}_U H(\cdot, D^{[k]}u) \\ &= E_\infty(u, U), \end{aligned}$$

so that $u \in \operatorname{AM}_c(\Omega, \mathbb{R}^N)$.

For the reverse implication, one can just replicate the proof of Theorem 1.2. Indeed, inequality (3.1) holds for every $u \in W^{1,\infty}(\Omega, \mathbb{R}^N)$. The opposite inequality is obtained exactly with the same reasoning, since the competitor u_λ belongs to the class $u + W_c^{1,\infty}(U, \mathbb{R}^N)$ (recall (3.5)). $\square$

4 On the gradient maximum principle for p -harmonic maps

In this section, we prove a gradient maximum principle for p -harmonic maps. To our best knowledge, a result of this kind is not present in the literature, at least in the vectorial case. For the scalar one, a gradient maximum principle for quasilinear elliptic equations can be found in [9, Chapter 15]. We refer also to [22, Appendix B] regarding gradient maximum and minimum principles for p -harmonic potentials on convex rings in the plane. In the case of general p -harmonic maps, however, one can only expect a gradient maximum principle to hold, since a minimum principle is not true in general, as existing examples show (see [2]).

We show the following result.

Theorem 4.1. *Let $\Omega \subset \mathbb{R}^n$ an open set and $p \in [2, \infty)$. Assume that $u : \Omega \rightarrow \mathbb{R}^N$ is a p -harmonic map, i.e. $u \in W_{\operatorname{loc}}^{1,p}(\Omega, \mathbb{R}^N)$ satisfying*

$$\operatorname{div}(|Du|^{p-2}Du) = 0 \quad \text{in } \Omega, \quad (4.1)$$

in the sense of distributions. Then, for every open set $U \Subset \Omega$, we have

$$\max_{\overline{U}} |Du| = \max_{\partial U} |Du|. \quad (4.2)$$

Remark 4.2. We observe that the above maxima are well defined since $u \in C_{\operatorname{loc}}^{1,\alpha}(\Omega, \mathbb{R}^N)$, from regularity theory for p -harmonic maps [23].

Moreover, we underline that Theorem 1.2, in the particular case when $E_\infty(u, U) = \operatorname{ess\,sup}_U |Du|$, cannot be deduced from Theorem 4.1 by passing to the limit as $p \rightarrow \infty$. Indeed, it is not known if vector valued absolute minimisers of E_∞ can be approximated locally uniformly by p -harmonic maps. More in general, the lack of good approximation results for absolute minimisers via p -harmonic maps is one of the reasons why existence of absolute minimisers is still open in the vectorial case.

The content of the following lemma is well known, but we report its proof for seek of completeness.

Lemma 4.3. *Let $\Omega \subset \mathbb{R}^n$ be an open set and $u : \Omega \rightarrow \mathbb{R}^N$ be a p -harmonic map, $p \in [2, \infty)$. Set $\mathcal{C} := \{x \in \Omega : Du(x) = 0\}$. Then $u \in C^\infty(\Omega \setminus \mathcal{C})$.*

Proof. Define $F : \mathbb{R}^{N \times n} \rightarrow \mathbb{R}^{N \times n}$ as $F(X) := |X|^{p-2}X$. Then, we have

$$DF(X) = (p-2)|X|^{p-4}X \otimes X + |X|^{p-2}I,$$

where I is the identity map in $(\mathbb{R}^{N \times n})^{\otimes 2}$. Using the following identity

$$(X \otimes X)Y : Z = (X : Y)(X : Z) \quad \forall X, Y, Z \in \mathbb{R}^{N \times n},$$

we have that

$$DF(X)Y : Z = Y : DF(X)Z \quad \forall X, Y, Z \in \mathbb{R}^{N \times n},$$

i.e. $DF(X)$ is a symmetric 4-tensor. Moreover, for every $X, Y \in \mathbb{R}^{N \times n}$, we have

$$|X|^{p-2}|Y|^2 \leq DF(X)Y : Y = (p-2)|X|^{p-4}(X : Y)^2 + |X|^{p-2}|Y|^2 \leq (p-1)|X|^{p-2}|Y|^2.$$

In particular, if we assume $0 < \lambda \leq |X| \leq \Lambda$, then

$$\lambda^{p-2}|Y|^2 \leq DF(X)Y : Y \leq (p-1)\Lambda^{p-2}|Y|^2, \quad (4.3)$$

so that, $DF(X)$ is elliptic in the Legendre sense (see [10]).

Now, fix $U \Subset \Omega \setminus \mathcal{C}$ and observe that u solves

$$\operatorname{div}(F(Du)) = 0 \quad \text{in } U.$$

For every fixed $U' \Subset U$ and $i \in \mathbb{N}$, the difference quotient of a function $f : U \rightarrow \mathbb{R}$

$$D_h^i f(x) := \frac{f(x + he_i) - f(x)}{h}$$

is well defined on U' , for every $h \in \mathbb{R}$ with $0 < |h| < \operatorname{dist}(U', \partial U)$.

By applying the difference quotient to the differential system solved by u , we have

$$\operatorname{div}(D_h^i(F(Du))) = 0 \quad \text{in } U'.$$

For $x \in U'$, we write

$$\begin{aligned} D_h^i(F(Du)) &= \frac{1}{h} \int_0^1 \frac{d}{dt} [F(Du(x) + thD_h^i Du(x))] dt \\ &= \left(\int_0^1 DF(Du(x) + thD_h^i Du(x)) dt \right) D_h^i Du(x). \end{aligned}$$

Observe that

$$\mathbb{A}(x) := \int_0^1 DF(Du(x) + thD_h^i Du(x)) dt$$

is a symmetric 4-tensor field on U' . Let us prove that it is Legendre elliptic: first notice that there exists $\lambda > 0$ such that $|Du| \geq \lambda$ in U' . In addition, since $u \in C^{1,\alpha}(U)$, for $|h|$ sufficiently small we have

$$|Du(x + he_i) - Du(x)| \leq \frac{\lambda}{2} \quad \forall x \in U'.$$

Thus, for $t \in [0, 1]$, we estimate

$$|Du(x) + thD_h^i Du(x)| = |Du(x) + t(Du(x + he_i) - Du(x))| \geq \lambda - t\frac{\lambda}{2} \geq \frac{\lambda}{2} \quad \forall x \in U'.$$

On the other hand,

$$|Du(x) + thD_h^i Du(x)| \leq 3\|Du\|_{L^\infty(U)} \quad \forall x \in U'.$$

From (4.3), we deduce the Legendre ellipticity of $\mathbb{A}$ in U' (with ellipticity constants independent of h).

Therefore, we have proved that $w := D_h^i Du$ solves the linear elliptic system

$$\operatorname{div}(\mathbb{A}w) = 0 \quad \text{in } U'.$$

By Schauder estimates [10, Thm. 5.19], since $\mathbb{A}$ has α -Hölder coefficients, we have $w \in C^{1,\alpha}(U')$. From the properties of the difference quotient, we infer $Du \in C^{2,\alpha}(U')$. By a bootstrap argument and arbitrariness of U' , we conclude $u \in C^\infty(\Omega \setminus \mathcal{C})$. $\square$

Now, we are ready to prove Theorem 4.1.

Proof of Theorem 4.1:

First, we observe that if $p = 2$, the result is straightforward: every harmonic map $u : \Omega \rightarrow \mathbb{R}^N$ is smooth and satisfies $\Delta|Du|^2 \geq 0$ on Ω . So, we can assume $p > 2$ in the following.

We divide the proof in two steps: firstly, we prove the gradient maximum principle outside the closed set $\mathcal{C} = \{x \in \Omega : Du(x) = 0\}$; second, we show the argument for a generic open set $U \Subset \Omega$. Of course, we can assume that $\Omega \setminus \mathcal{C} \neq \emptyset$, otherwise u is constant on Ω .

Step 1: assume $U \Subset \Omega \setminus \mathcal{C}$.

In the following, the symbols $\cdot$ and $\langle \cdot, \cdot \rangle$ stand for the inner products in $\mathbb{R}^n$ and $\mathbb{R}^N$ respectively, while $:$ stands for the Hilbert-Schmidt inner product in $\mathbb{R}^{N \times n}$.

By Lemma 4.3, we have $u \in C^\infty(\overline{U}, \mathbb{R}^N)$. So, we can differentiate (4.1) with respect to x_i , obtaining

$$0 = \partial_i [\operatorname{div}(|Du|^{p-2} Du)] = \operatorname{div}((p-2)|Du|^{p-4}(Du : D\partial_i u)Du + |Du|^{p-2} D\partial_i u) \quad \text{in } U.$$

By taking the inner product with $\partial_i u$, we have

$$\begin{aligned} 0 &= \langle \operatorname{div}((p-2)|Du|^{p-4}(Du : D\partial_i u)Du + |Du|^{p-2} D\partial_i u), \partial_i u \rangle \\ &= \operatorname{div}((p-2)|Du|^{p-4}(Du : D\partial_i u)\langle Du, \partial_i u \rangle + |Du|^{p-2}\langle D\partial_i u, \partial_i u \rangle) - \\ &\quad - (p-2)|Du|^{p-4}(Du : D\partial_i u)^2 - |Du|^{p-2}|D\partial_i u|^2, \quad \text{in } U. \end{aligned} \tag{4.4}$$

From (4.4), we deduce

$$\operatorname{div}((p-2)|Du|^{p-4}(Du : D\partial_i u)\langle Du, \partial_i u \rangle + |Du|^{p-2}\langle D\partial_i u, \partial_i u \rangle) \geq 0, \quad \text{in } U. \tag{4.5}$$

By taking the sum over $i = 1, \dots, n$ in (4.5) and observing that

$$\sum_{i=1}^n (Du : D\partial_i u)\langle Du, \partial_i u \rangle = (Du^T Du) \cdot \langle D^2 u, Du \rangle,$$

we obtain

$$\operatorname{div}((p-2)|Du|^{p-4}(Du^T Du) \cdot \langle D^2 u, Du \rangle + |Du|^{p-2}\langle D^2 u, Du \rangle) \geq 0, \quad \text{in } U. \tag{4.6}$$

Now, we define

$$f := \frac{1}{p}|Du|^p, \quad A := I + (p-2)\frac{Du^T Du}{|Du|^2},$$

where I is the identity matrix of $\mathbb{R}^{n \times n}$. Thus,

$$Df = |Du|^{p-2} \langle D^2 u, Du \rangle,$$

so that we can rewrite (4.6) as

$$\operatorname{div}(ADf) \geq 0, \quad \text{in } U. \quad (4.7)$$

Since $u \in C^\infty(\overline{U}, \mathbb{R}^N)$, the function f is a classical subsolution to a linear elliptic partial differential inequality in divergence form, with smooth and bounded coefficients. By the maximum principle for elliptic PDEs, we have

$$\max_{\overline{U}} f = \max_{\partial U} f,$$

yielding (4.2).

Step 2: general case.

Let $U \Subset \Omega$ be an open subset. Then, without loss of generality, we can assume that $U \not\subset \mathcal{C}$. Indeed, if $U \subset \mathcal{C}$, then also $\overline{U} \subset \mathcal{C}$, so that (4.2) trivially holds. Therefore, since $U \setminus \mathcal{C}$ is a non-empty open subset of Ω , there exists $\varepsilon > 0$ such that $U_\varepsilon := U \setminus \overline{\mathcal{C}^\varepsilon}$ is a non-empty open subset of Ω , where $\mathcal{C}^\varepsilon$ is the open ε -neighbourhood of $\mathcal{C}$. Thanks to *Step 1*, since $U_\varepsilon \Subset \mathbb{R}^n \setminus \mathcal{C}$, we have

$$\max_{\overline{U}_\varepsilon} |Du| = \max_{\partial U_\varepsilon} |Du|. \quad (4.8)$$

Now, set $M := \max_{\overline{U}} |Du| > 0$. By possibly reducing ε in size, the continuity of Du ensures

$$\max_{\overline{\mathcal{C}^\varepsilon}} |Du| < M. \quad (4.9)$$

Hence

$$M = \sup_U |Du| = \max \left\{ \sup_{U_\varepsilon} |Du|, \max_{\overline{\mathcal{C}^\varepsilon}} |Du| \right\} = \sup_{U_\varepsilon} |Du| = \max_{\overline{U}_\varepsilon} |Du|.$$

In particular, from (4.8) we have

$$\max_{\partial U_\varepsilon} |Du| = M.$$

On the other hand, we can write

$$\partial U_\varepsilon = (\partial U \setminus \overline{\mathcal{C}^\varepsilon}) \cup (U \cap \partial \mathcal{C}^\varepsilon),$$

so that, using (4.9),

$$M = \sup_{\partial U_\varepsilon} |Du| = \max \left\{ \sup_{\partial U \setminus \overline{\mathcal{C}^\varepsilon}} |Du|, \sup_{U \cap \partial \mathcal{C}^\varepsilon} |Du| \right\} \leq \max \left\{ \max_{\partial U} |Du|, \max_{\overline{\mathcal{C}^\varepsilon}} |Du| \right\} = \max_{\partial U} |Du|,$$

which ensures the validity of (4.2). $\square$

ACKNOWLEDGEMENTS: S.C and R.M. acknowledge partial financial support through the EP-SRC grant EP/X017206/1. S.C and N.K. acknowledge partial financial support through the EPSRC grant EP/X017109/1. S.C. is a member of Gruppo Nazionale per l'Analisi Matematica, la Probabilità e le loro Applicazioni (GNAMPA) of the Istituto Nazionale di Alta Matematica (INdAM) of Italy.

DATA AVAILABILITY: Our manuscript has no associated data.

CONFLICT OF INTEREST: The authors have no Conflict of interest to declare for this article.

References

- [1] G. Aronsson, *Minimization problems for the functional $\sup_x F(x, f(x), f'(x))$ I, II, III*, Arkiv fur Mat. **33** - 53 (1965); 409 - 431 (1966); 509 - 512 (1969).
- [2] A. Balci, L. Behn, L. Diening, J. Storn, *Examples of p -harmonic Maps*, SIAM Journal on Mathematical Analysis **58**(1), 260-275 (2026).
- [3] E.N. Barron, R.R. Jensen, C.Y. Wang, *The Euler equation and absolute minimizers of L^∞ functionals*, Arch. Rational Mech. Anal. **157**, 255-283 (2001).
- [4] A. Braides, *A handbook of Γ -convergence*, Handbook of Differential Equations: Stationary Partial Differential Equations, North-Holland, **3**, 101-213 (2006).
- [5] C. Brizzi, L. De Pascale, *A property of Absolute Minimizers in L^∞ Calculus of Variations and of solutions of the Aronsson-Euler equation*, Advances in Differential Equations **28** (3/4), 287-310 (2023).
- [6] S. Carano, N. Katzourakis, R. Moser, *Existence, uniqueness and characterisation of vector-valued absolute minimisers for a second order L^∞ -variational problem*, Preprint arXiv <https://arxiv.org/abs/2504.04181> (2025).
- [7] S. Carano, R. Moser, *An L^∞ -variational problem involving the Fractional Laplacian*, Preprint arXiv <https://arxiv.org/abs/2510.14476> (2025).
- [8] B. Dutton, N. Katzourakis, *On Second-Order L^∞ Variational Problems with Lower-Order Terms*, Discrete Contin. Dyn. Syst., **49**, 1-21 (2026).
- [9] D. Gilbarg, N. Trudinger, *Elliptic Partial Differential Equations of Second Order*. Classics in Mathematics, reprint of the 1998 edition, Springer.
- [10] M. Giaquinta, L. Martinazzi, *An Introduction to the Regularity Theory for Elliptic Systems, Harmonic Maps and Minimal Graphs*, Publications of the Scuola Normale Superiore (2018).
- [11] N. Katzourakis, *L^∞ Variational Problems for maps and the Aronsson PDE System*, J. Differential Equations **253**(7), 2123 - 2139 (2012).
- [12] N. Katzourakis, *An L^∞ regularisation strategy to the inverse source identification problem for elliptic equations*, SIAM Journal Math. Analysis **51**(2), 1349-1370 (2019).
- [13] N. Katzourakis, *An Introduction to Viscosity Solutions for Fully Nonlinear PDE with Applications to Calculus of Variations in L^∞* , Springer Briefs in Mathematics, DOI 10.1007/978-3-319-12829-0 (2015).
- [14] N. Katzourakis, *∞ -minimal submanifolds*, Proc. Amer. Math. Soc. **142**, 2797-2811 (2014).
- [15] N. Katzourakis, *On the structure of ∞ -harmonic maps*, Comm. Partial Differential Equations **39**, 2091–2124 (2014).
- [16] N. Katzourakis, R. Moser, *Existence, uniqueness and structure of second order absolute minimisers*, ARMA **231**(3), 1615-1634 (2019).
- [17] N. Katzourakis, R. Moser, *Minimisers of supremal functionals and mass-minimising 1-currents*, Calc. Var. **64**(26) (2025).

- [18] N. Katzourakis, R. Moser, *Variational problems in L^∞ involving semilinear second order differential operators*, ESAIM: COCV, **29**(76) (2023).
- [19] N. Katzourakis, R. Moser. *Existence, uniqueness and characterisation of local minimisers in higher order Calculus of Variations in L^∞* , to appear in Analysis & PDE (2026).
- [20] N. Katzourakis, T. Pryer, *On the numerical approximation of p -Biharmonic and ∞ -Biharmonic functions*, Numerical Methods for Partial Differential Equations, **35**(1), 155-180 (2018).
- [21] N. Katzourakis, T. Pryer, *2nd order L^∞ Variational Problems and the ∞ -Polylaplacian*, Advances in Calculus of Variations **13**, 115-140 (2020).
- [22] F. Peng, Y. R. Zhang, Y. Zhou, *Regularity from p -harmonic potentials to ∞ -harmonic potentials in convex rings*, Proc. London Math. Soc. **130**(6), e70062 (2025).
- [23] K. Uhlenbeck, *Regularity for a class of non-linear elliptic systems*, Acta Math. **138**, 219–240 (1977).

SIMONE CARANO

Department of Mathematics and Statistics, University of Reading
 Whiteknights Campus, Pepper Lane, Reading RG6 6AX, UK
 E-mail: s.carano@reading.ac.uk

NIKOS KATZOURAKIS

Department of Mathematics and Statistics, University of Reading
 Whiteknights Campus, Pepper Lane, Reading RG6 6AX, UK
 E-mail: n.katzourakis@reading.ac.uk

ROGER MOSER

Department of Mathematical Sciences, University of Bath
 Bath BA2 7AY, UK
 E-mail: r.moser@bath.ac.uk