

Linear Model Selection and Regularization Exercises

Simone Collier

1 Getting Started

Start by loading the packages that have the data we need. If you need to install the packages first then run `install.packages("PACKAGENAME")` in your console before running the code chunk.

```
library(ISLR2)
library(leaps)
library(glmnet)
```

We will be making use of the `Hitters` data set in the `ISLR` package. It contains baseball player's salaries along with other information about their performance in the previous year.

```
names(Hitters)
```

```
## [1] "AtBat"      "Hits"       "HmRun"      "Runs"       "RBI"        "Walks"
## [7] "Years"      "CAtBat"     "CHits"      "CHmRun"     "CRuns"      "CRBI"
## [13] "CWalks"     "League"     "Division"   "PutOuts"    "Assists"    "Errors"
## [19] "Salary"     "NewLeague"
```

We can use `is.na()` to see whether there is any missing data in the set. Using the `sum()` function in combination on the `Salary` column will tell us how many player's salaries we are missing.

```
sum(is.na(Hitters$Salary))
```

```
## [1] 59
```

The `na.omit()` function allows us to create a new data set from `Hitters` that removes rows with no entries.

```
Hitters.noNA <- na.omit(Hitters)
```

We want to predict a baseball player's salary based on the other predictors in the data set using linear regression.

2 Subset Selection

2.1 Best Subset Selection

We can use the `regsubsets()` function from the `leaps` package to perform best subset selection. It identifies the best model (smallest RSS) among those with the same number of predictors.

```
best.subset.fit <- regsubsets(Salary ~ ., data = Hitters.noNA)
summary(best.subset.fit)
```

```
## Subset selection object
## Call: regsubsets.formula(Salary ~ ., data = Hitters.noNA)
## 19 Variables (and intercept)
##           Forced in Forced out
## AtBat      FALSE      FALSE
## Hits       FALSE      FALSE
## HmRun       FALSE      FALSE
## Runs       FALSE      FALSE
## RBI        FALSE      FALSE
## Walks      FALSE      FALSE
## Years      FALSE      FALSE
## CAtBat     FALSE      FALSE
## CHits      FALSE      FALSE
## CHmRun     FALSE      FALSE
## CRuns      FALSE      FALSE
## CRBI       FALSE      FALSE
## CWalks     FALSE      FALSE
## LeagueN    FALSE      FALSE
## DivisionW  FALSE      FALSE
## PutOuts    FALSE      FALSE
## Assists    FALSE      FALSE
## Errors     FALSE      FALSE
## NewLeagueN FALSE      FALSE
## 1 subsets of each size up to 8
## Selection Algorithm: exhaustive
##           AtBat Hits HmRun Runs RBI Walks Years CAtBat CHits CHmRun CRuns CRBI
## 1 ( 1 ) " " " " " " " " " " " " " " " " " " " " " " " " " " " " " " " " " " "
## 2 ( 1 ) " " "*" " " " " " " " " " " " " " " " " " " " " " " " " " " "
## 3 ( 1 ) " " "*" " " " " " " " " " " " " " " " " " " " " " " " " " " "
## 4 ( 1 ) " " "*" " " " " " " " " " " " " " " " " " " " " " " " " " " "
## 5 ( 1 ) "*" "*" " " " " " " " " " " " " " " " " " " " " " " " " " " "
## 6 ( 1 ) "*" "*" " " " " " " "*" " " " " " " " " " " " " " " " " " " "
## 7 ( 1 ) " " "*" " " " " " " "*" " " "*" "*" "*" " " " " " " " " " " "
## 8 ( 1 ) "*" "*" " " " " " " "*" " " " " " " "*" "*" " " " " " " " " " "
##           CWalks LeagueN DivisionW PutOuts Assists Errors NewLeagueN
## 1 ( 1 ) " " " " " " " " " " " " " " " " " " " " " " " " " " " " "
## 2 ( 1 ) " " " " " " " " " " " " " " " " " " " " " " " " " " " " "
## 3 ( 1 ) " " " " " " "*" " " " " " " " " " " " " " " " " " " " " " "
## 4 ( 1 ) " " " " "*" "*" " " " " " " " " " " " " " " " " " " " " " "
## 5 ( 1 ) " " " " "*" "*" " " " " " " " " " " " " " " " " " " " " " "
## 6 ( 1 ) " " " " "*" "*" " " " " " " " " " " " " " " " " " " " " " "
## 7 ( 1 ) " " " " "*" "*" " " " " " " " " " " " " " " " " " " " " " "
## 8 ( 1 ) "*" " " "*" "*" " " " " " " " " " " " " " " " " " " " " " " "
```

The `summary()` function displays the best combination of predictors for models with 1-8 predictors. The asterisks indicate that the variable is included in the model. The `nvmax` argument in the function `regsubsets()` can be increased to return models with more than 8 predictors.

```
# Perform best subset selection with all predictors (all columns except Salary)
best.subset.fit <- regsubsets(Salary ~ ., data = Hitters.noNA, nvmax = ncol(Hitters) - 1)
summary.best.subset <- summary(best.subset.fit)
```

2.2 Stepwise Selection


We can use `regsubsets()` to perform forward and backward stepwise selection with the argument `method = "forward"` or `method = "backward"`.

```
fwd.fit <- regsubsets(Salary ~ ., data = Hitters.noNA,
                      nvmax = ncol(Hitters) - 1, method = "forward")
bwd.fit <- regsubsets(Salary ~ ., data = Hitters.noNA,
                      nvmax = ncol(Hitters) - 1, method = "backward")
summary.fwd <- summary(fwd.fit)
summary.bwd <- summary(bwd.fit)
```

2.3 Deciding Between Models using Indirect Error Estimation

The `summary()` function for `regsubsets()` output gives the R^2 (`rsq`), RSS (`rss`), adjusted R^2 (`adjr2`), C_p (`cp`), and BIC (`bic`). We can make a plot of the adjusted R^2 statistic of the best subset selection outputs to help us decide which model to select. We can use the `which.max()` function to find at what index the maximum adjusted R^2 value occurs.

```
# type = "l" connects the points of the plot with lines
plot(summary.best.subset$adjr2, xlab = "Number of Predictors",
      ylab = "Adjusted R^2", type = "l")
# find the index of the maximum adjusted R^2
max.adj2 <- which.max(summary.best.subset$adjr2)
# add point at maximum adjusted R^2
points(max.adj2, summary.best.subset$adjr2[max.adj2], col = "red")
```



Recall that in the case of the C_p and BIC statistics, the best model will have the smallest C_p or BIC value.

Plot the C_p and BIC statistics and include the minimum points. Based on the three plots, which model size is the best? Justify your answer and list the predictors that are included in the model.

Choose which model size you think is the best for both the forward and backward selection. How do the three models we have chosen compare?

2.4 Deciding Between Models using Direct Error Estimation

2.4.1 Validation Set Approach

We just saw how we can indirectly estimate the test error of our models. We will now look at directly estimate the test error using the validation set approach and cross-validation.

We start by splitting the observations into a training set and a test set. We can do this by creating a random sample of TRUE and FALSE that is the same size as our data set. TRUE will signify that the element is in the training set and FALSE will signify that the element is in the test set.

```
set.seed(1)
train <- sample(c(TRUE, FALSE), nrow(Hitters.noNA), replace = TRUE)
test <- (!train)
```

Now we can perform best subset selection on the training set.

```
best.subset.fit2 <- regsubsets(Salary ~ ., data = Hitters.noNA[train, ], nvmax = ncol(Hitters.noNA) - 1)
```

Now that we have our fitted models we need to estimate the test error by predicting the response of the observations in our test set. This is a little tricky since the function `regsubsets()` which we used to fit our models does not work with the `predict()` function we have been using in previous sections. This means we will have to make our own function called `predict.regsubsets()`. Before we make the function let's go through the steps to understand what our function will need to do.

We know that `best.subset.fit2` contains 19 linear regression models. What information do we need to extract from `best.subset.fit2` in order to be able to use each of the 19 models to predict the responses on the test set?... The fitted coefficients! Recall from the linear regression module that responses for linear models are found using:

$$\hat{Y} = \hat{\beta}_0 + \hat{\beta}_1 X_1 + \hat{\beta}_2 X_2 + \cdots + \hat{\beta}_p X_p$$

So we need to write a function that extracts the fitted coefficients for each model size and then multiplies the corresponding predictors for each test observations. Let's write out each step first...

To start let's get the test data on its own. Instead of using a data frame for this we will use a matrix. This will allow us to multiply our fitted coefficients and the test observations in the matrix directly. The `model.matrix()` function is commonly used for build a predictor matrix from data in a regression context.

```
test.mat <- model.matrix(Salary ~ ., data = Hitters.noNA[test, ])
```

Run `test.mat` see what it looks like.

Now we need to extract the coefficients from each of the 19 fitted models in `best.subset.fit2` and multiply them with the corresponding test predictors. Then we can compute the test error. We can write a `for` loop for this.

```
# Make empty vector to be filled with the computed test errors
val.errors <- numeric()
for (i in 1:19) {
  # Extract the coefficients of the i-th model
  coefs <- coef(best.subset.fit2, id = i)
  # Extract the predictors from the test matrix that are relevant to the i-th model
  obs <- test.mat[, names(coefs)]
  # Predict the response on the test observations
  pred <- obs %*% coefs
  # Compute the test MSE for the i-th model
  mse <- mean((Hitters.noNA$Salary[test] - pred)^2)
  # Add the test error to the list
  val.errors <- c(val.errors, mse)
}

val.errors
```

```
## [1] 164377.3 144405.5 152175.7 145198.4 137902.1 139175.7 126849.0 136191.4
## [9] 132889.6 135434.9 136963.3 140694.9 140690.9 141951.2 141508.2 142164.4
## [17] 141767.4 142339.6 142238.2
```

We can find the model with the smallest validation set error and choose this as our model.

```
which.min(val.errors)
```

```
## [1] 7
```

Now that we have the method layed out, we can write a generic function that we can use to predict responses for models from `regsubsets()`.

```
# object is the `regsubsets()` output (i.e. best.subset.fit2)
predict.regsubsets <- function(object, newdata, id, ...) {
  # get the formula used in `regsubsets()`
  form <- as.formula(object$call[[2]])
  mat <- model.matrix(form, newdata)
  coefs <- coef(object, id = id)
  vars <- names(coefs)
  mat[, vars] %*% coefs
}
```

Let's try the function out.

```
predict.regsubsets(best.subset.fit2, Hitters.noNA[test, ], 7)
```

```
##                [,1]
## -Alvin Davis      727.39222
## -Alfredo Griffin  673.80948
## -Andre Thornton   628.54831
## -Alan Trammell     952.23602
## -Buddy Biancalana  58.65845
## -Bruce Bochy      101.17418
## -Barry Bonds      456.73992
## -Bobby Bonilla    407.28753
## -Billy Hatcher    115.22671
## -Bill Madlock     730.38011
## -Bob Melvin       54.36360
## -BillyJo Robidoux 302.31423
## -Chris Bando      303.06348
## -Chili Davis      794.88821
## -Curt Ford        210.03316
## -Carney Lansford  712.22213
## -Chet Lemon       794.89166
## -Candy Maldonado  66.70965
## -Craig Reynolds   314.13616
## -Cal Ripken       985.00289
## -Cory Snyder      206.10648
## -Chris Speier     396.98031
## -Curt Wilkerson   51.96348
## -Daryl Boston     139.57096
## -Dave Concepcion  702.98418
## -Doug DeCinces    646.27730
## -Damaso Garcia    632.80611
## -Dan Gladden      393.65465
## -Dave Henderson   398.26524
## -Don Mattingly    1350.43868
```

| | |
|-----------------------|------------|
| ## -Dale Murphy | 951.93960 |
| ## -Dwayne Murphy | 531.52823 |
| ## -Dave Parker | 1126.79635 |
| ## -Dan Pasqua | 481.40267 |
| ## -Darrell Porter | 320.23183 |
| ## -Don Slaught | 251.27139 |
| ## -Darryl Strawberry | 688.91728 |
| ## -Dale Sveum | 220.36953 |
| ## -Danny Tartabull | 400.92118 |
| ## -Eddie Milner | 375.77744 |
| ## -Eddie Murray | 1464.82918 |
| ## -George Bell | 750.67771 |
| ## -Glenn Braggs | 95.95374 |
| ## -George Brett | 1322.80475 |
| ## -Gary Gaetti | 580.35839 |
| ## -George Hendrick | 809.32355 |
| ## -Garth Iorg | 336.80881 |
| ## -Gary Pettis | 582.39550 |
| ## -Gary Redus | 523.01393 |
| ## -Glenn Wilson | 562.22245 |
| ## -Harold Baines | 684.10692 |
| ## -Howard Johnson | 310.02633 |
| ## -Hal McRae | 613.96246 |
| ## -Harold Reynolds | 41.81559 |
| ## -John Cangelosi | 398.01440 |
| ## -Jim Dwyer | 312.45512 |
| ## -Julio Franco | 660.16835 |
| ## -Jim Gantner | 599.30176 |
| ## -Johnny Grubb | 549.95017 |
| ## -John Kruk | 347.76920 |
| ## -Jeffrey Leonard | 371.20901 |
| ## -Jim Morrison | 578.52033 |
| ## -John Moses | 201.13809 |
| ## -Johnny Ray | 856.90827 |
| ## -Jerry Royster | 465.45299 |
| ## -John Russell | 290.42612 |
| ## -Jim Sundberg | 480.91708 |
| ## -Joel Youngblood | 308.26960 |
| ## -Kal Daniels | 190.77503 |
| ## -Kirk Gibson | 825.60821 |
| ## -Ken Griffey | 1095.55005 |
| ## -Kent Hrbek | 917.51361 |
| ## -Kevin Mitchell | 340.34197 |
| ## -Ken Oberkfell | 671.24863 |
| ## -Ken Phelps | 547.38188 |
| ## -Kirby Puckett | 686.02908 |
| ## -Len Dykstra | 644.62235 |
| ## -Lance Parrish | 821.74381 |
| ## -Larry Parrish | 813.01797 |
| ## -Larry Sheets | 224.19213 |
| ## -Lou Whitaker | 912.59432 |
| ## -Mike Aldrete | 216.38045 |
| ## -Mariano Duncan | 106.77883 |
| ## -Mike Heath | 422.13467 |

```
## -Mike Marshall      208.79210
## -Mike Schmidt       55.02747
## -Milt Thompson     266.45271
## -Mitch Webster     642.69038
## -Mookie Wilson     757.21771
## -Marvell Wynne     187.84572
## -Ozzie Guillen     38.78130
## -Pete Incaviglia   276.33167
## -Pete Rose         1563.41556
## -Rafael Belliard   177.67184
## -Randy Bush        320.64594
## -Rick Cerone       381.64192
## -Ron Cey           747.79334
## -Rick Dempsey      527.71147
## -Reggie Jackson    1256.48726
## -Ron Kittle        164.21179
## -Rick Leach        248.33912
## -Rick Manning      584.50829
## -Rance Mulliniks   456.03845
## -Rafael Ramirez    243.50198
## -Rick Schu         246.50534
## -Rob Wilfong       157.03932
## -Robin Yount       1529.43379
## -Scott Bradley     136.45261
## -Sid Bream         790.53638
## -Steve Buechele    109.03581
## -Shawon Dunston    185.99881
## -Scott Fletcher    490.36808
## -Steve Jeltz       382.78993
## -Steve Sax         925.56961
## -Tom Foley         287.71981
## -Tony Gwynn        832.23122
## -Terry Harper      175.26040
## -Tommy Herr        758.38409
## -Tim Lulett        -50.06588
## -Terry Kennedy     120.53833
## -Tito Landrum      194.56136
## -Tom Paciorek      391.51903
## -Terry Pendleton   215.25127
## -Terry Puhl        392.53430
## -Tim Teufel        336.13853
## -Vince Coleman     472.54412
## -Will Clark        459.86191
## -Willie Randolph   1187.96388
## -Wayne Tolleson    390.03303
```

Now that we have decided to go with the 7 variable model we need to refit the model using ALL the observations. We did this previously so we can go ahead and grab the corresponding model.

```
coef(best.subset.fit, id = 7)
```

```
## (Intercept)      Hits      Walks      CAtBat      CHits      CHmRun
## 79.4509472    1.2833513    3.2274264    -0.3752350    1.4957073    1.4420538
```



```
##      DivisionW      PutOuts
## -129.9866432      0.2366813
```

Note this model is a little different than the one that was fit with only the training observations... It even has different variables.

```
coef(best.subset.fit2, id = 7)
```

```
## (Intercept)      AtBat      Hits      Walks      CRuns      CWalks
##  67.1085369 -2.1462987  7.0149547  8.0716640  1.2425113 -0.8337844
##      DivisionW      PutOuts
## -118.4364998      0.2526925
```

2.4.2 Cross-validation

Let's try out the cross-validation technique for estimating test error. We will use $k = 10$ folds.

```
k <- 10
n <- nrow(Hitters.noNA)
set.seed(1)
folds <- sample(rep(1:k, length = n))
```

We can make a matrix for storing the computed test errors, one for each fold and model size.

```
cv.errors <- matrix(nrow = k, ncol = 19)
```

Now we can go about computing the errors by first predicting the responses for each fold using each of the 19 models with our function `predict.regsubsets()` and then computing the MSE.

```
for (i in 1:k) {
  # Fit the linear regression by best subset selection using all of the observations
  # except those in the i-th fold
  fold.fit <- regsubsets(Salary ~ ., data = Hitters.noNA[folds != i, ], nvmax = 19)

  for (j in 1:19) {
    # Predict the response for the observations in the i-th fold using the fitted model of size j
    pred <- predict.regsubsets(fold.fit, Hitters.noNA[folds == i, ], id = j)
    # Compute the MSE using the predictions and add it to the matrix
    cv.errors[i, j] <- mean((Hitters.noNA$Salary[folds == i] - pred)^2)
  }
}
```

Now we have the cross-validation error for each fold and each model size. If we take the mean of each column in the matrix, we will have the cross-validation error for each model size.

```
mean.cv.errors <- apply(cv.errors, 2, mean)
mean.cv.errors
```

```
## [1] 143439.8 126817.0 134214.2 131782.9 130765.6 120382.9 121443.1 114363.7
## [9] 115163.1 109366.0 112738.5 113616.5 115557.6 115853.3 115630.6 116050.0
## [17] 116117.0 116419.3 116299.1
```

Make a plot of the cross-validation errors and identify the model with the lowest error with a point. Refit the models with best subset selection using the complete data set. Return the coefficients of the model which was identified as the one with the lowest CV error.

3 Ridge Regression

We will use the `glmnet()` function from the `glmnet` package to perform ridge regression. Instead of using a data frame, the `glmnet()` function requires that the predictors be in the form of a matrix and the response be in the form of a vector.

```
x <- model.matrix(Salary ~., Hitters.noNA)[, -1]
y <- Hitters.noNA$Salary
```

The `model.matrix()` automatically transforms qualitative variables into dummy variables.

The crucial step for ridge regression and the lasso is selecting the tuning parameter. We will start by fitting with many λ s (this is the default for the function) and then use cross-validation after the fact to choose the λ with the smallest MSE. Let's first split our data into a training and testing set so that we can compute the test error after the fact.


```
set.seed(1)
train <- sample(1:nrow(x), nrow(x)/2)
test <- (-train)
y.test <- y[test]
```

We choose $\alpha = 0$ in the `glmnet()` function perform ridge regression. The function standardizes the variables automatically so we need not worry about that.

```
ridge.mod <- glmnet(x[train, ], y[train], alpha = 0)
```

Now we will use the function `cv.glmnet()`. This function performs cross-validation with a bunch of λ values with a default of `nfolds = 10`.

```
set.seed(1)
cv.ridge <- cv.glmnet(x[train, ], y[train], alpha = 0)
plot(cv.ridge)
```



We can find the λ with the smallest cross-validation error and make predictions for the test set using the model in `ridge.mod` with the corresponding λ .

```
best.lambda <- cv.ridge$lambda.min
ridge.pred <- predict(ridge.mod, s = best.lambda, newx = x[test, ])
```

Compute the test MSE.

Now we can refit the ridge regression using the complete data set and return the coefficients.

```
ridge <- glmnet(x, y, alpha = 0)
predict(ridge, type = "coefficients", s = best.lambda)[1:20,]
```

| | | | | | | |
|----|-------------|------------|-------------|--------------|------------|------------|
| ## | (Intercept) | AtBat | Hits | HmRun | Runs | RBI |
| ## | 15.44383120 | 0.07715547 | 0.85911582 | 0.60103106 | 1.06369007 | 0.87936105 |
| ## | Walks | Years | CAtBat | CHits | CHmRun | CRuns |
| ## | 1.62444617 | 1.35254778 | 0.01134999 | 0.05746654 | 0.40680157 | 0.11456224 |
| ## | CRBI | CWalks | LeagueN | DivisionW | PutOuts | Assists |
| ## | 0.12116504 | 0.05299202 | 22.09143197 | -79.04032656 | 0.16619903 | 0.02941950 |
| ## | Errors | NewLeagueN | | | | |
| ## | -1.36092945 | 9.12487765 | | | | |


4 The Lasso

We will perform the lasso the exact same way as ridge regression except we have `alpha = 1` in the `glmnet()` function to specify the lasso method is to be used.

```
lasso.mod <- glmnet(x[train, ], y[train], alpha = 1)
```

Now we can perform cross-validation and compute the MSE.

```
set.seed(1)
cv.lasso <- cv.glmnet(x[train, ], y[train], alpha = 1)
plot(cv.lasso)
```



```
best.lambda2 <- cv.lasso$lambda.min
lasso.pred <- predict(lasso.mod, s = best.lambda2, newx = x[test, ])
mse <- mean((lasso.pred - y.test)^2)
mse
```

```
## [1] 143668.8
```

```
lasso <- glmnet(x, y, alpha = 1)
predict(lasso, type = "coefficients", s = best.lambda2)[1:20, ]
```

```
##      (Intercept)      AtBat      Hits      HmRun      Runs
##      -3.04787656      0.00000000      2.02551572      0.00000000      0.00000000
##           RBI           Walks           Years           CAtBat           CHits
##           0.00000000      2.26853781      0.00000000      0.00000000      0.00000000
##          CHmRun          CRuns          CRBI          CWalks          LeagueN
```

| | | | | | |
|----|---------------|------------|------------|-------------|-------------|
| ## | 0.01647106 | 0.21177390 | 0.41944632 | 0.00000000 | 20.48456551 |
| ## | DivisionW | PutOuts | Assists | Errors | NewLeagueN |
| ## | -116.59062083 | 0.23718459 | 0.00000000 | -0.94739923 | 0.00000000 |

The resulting model has 10 coefficients that are exactly equal to zero so it performed variable selection!

Fit a linear regression model to the data. Compute the test MSE for the linear regression model. Compare the results from the linear regression, ridge regression, and the lasso. Which is the best model and why?

These exercises were adapted from : James, Gareth, et al. An Introduction to Statistical Learning: with Applications in R, 2nd ed., Springer, 2021.