

Borromean Triads, the Ring Lemma, and Spectral Non-Dispersal

Jonathan Simons. This record (10.5281/zenodo.19842060) is a 2026 exploratory draft. It remains published and citable.¹ This page is the public-facing face of the record. It is not a retraction notice. The original draft follows the errata.

Live exploratory stack (cite these, not prize packaging)

Live exploratory stack: Φ -renormalization 10.5281/zenodo.22050974 (not a Clay Navier-Stokes proof); Route C 10.5281/zenodo.22050963 (not a proof of RH); Ring lemma 10.5281/zenodo.22050976 (Goldbach remains conditional on SND); T2 under SND 10.5281/zenodo.22050965 and Bridge* stay named derived objects; inverse-GCD note 10.5281/zenodo.22050962 is a restricted Rayleigh bound, not a full-spectrum floor.

¹ August 2026 prize-claim language walked back; see errata below / page 2 of this file and status note 10.5281/zenodo.22050978.

Errata: prize-claim language walked back

The pages after this one are the original 2026 draft, unchanged. They are on the record so the walk-back is auditable. The following prize-claim language in that draft does not stand.

Why the August 2026 action happened

In August 2026, prize-claim language on several 2026 records was walked back. The walk-back was necessary: those drafts packaged Navier-Stokes, the Riemann hypothesis, and Goldbach as closed, and that packaging does not hold. The walk-back was executed badly. Public titles were stamped on 21 August 2026 so the landing page looked like a crime scene. The files were never unpublished, never tombstoned, and never taken off open access. This pack restores the public face: a clean title, an honest first page, and the errata on page 2.

Prize-claim language that does not stand

Clay Statement B / Navier-Stokes millennium packaging. The June drafts treated a regularity claim as closed. That language is walked back. The Φ -renormalization note is the live fluids object.

Quantum Lens / Q6 Goldbach as a theorem. Goldbach stays conditional on SND. Q6 is not a proof.

Montgomery-Dyson / pair correlation as a finished RH argument. That packaging is walked back, including the May 14 "Coincidence Resolved" draft. Live RH work is exploratory Route C and the Track B Möbius-GCD attack (obstruction, not a proof).

Three-in-one / $\text{SND} \equiv \text{GNC} \equiv \text{Bridge}$ as closures. Those identities were used as if they closed NS, RH, and Goldbach. They do not. The June drafts that argued that way remain published; the prize language is walked back.

Full-spectrum $\lambda_{\min}(Q_N) > -1/2$ as a RH proof. That bound is not a proof of RH.
Inverse-GCD $1/\text{gcd}$ is not the live Route C operator.

This page is the errata. It is not a tombstone. No Millennium problem is claimed. Domain Architect is a local FRA classifier (inquiry). ChatVault is search. Neither wrote these Zenodo titles.

Borromean Triads, the Ring Lemma, and Spectral Non-Dispersal

A Conditional Regularity Framework for the 3D Incompressible Navier-Stokes Equations

Jonathan Simons

Prime Field Technologies LLC, Savannah, Georgia

simonsmedical@icloud.com

NAV-42 Patent Pending (Provisional filed March 15, 2026)

Abstract

We present a conditional regularity framework for the three-dimensional incompressible Navier-Stokes equations on the periodic torus T^3 . The framework rests on three contributions. First, a three-operator augmented system (Q_1, Q_3, Q_6) designed to regulate nonlinear energy transfer across Littlewood-Paley shells by intervening only where spectral concentration develops. Second, the *Ring Lemma*: on a single LP shell S_j^* , the vorticity direction field $\xi = \omega/\|\omega\|$ satisfies $\|\text{grad } \xi\|_{L^\infty} \leq C \cdot 2^{j*}$, directly answering Constantin-Fefferman Question 2.2 -- the rotation rate of the vorticity direction is controlled by the shell frequency, not by the vorticity amplitude. Third, under a four-part *Spectral Non-Dispersal* condition [SND], a Global Summation Lemma yields conditional global H^1 bounds. One precisely-stated open problem remains. No claim of unconditional regularity is made.

1. Introduction

1.1 The Problem and the Obstruction

The 3D incompressible Navier-Stokes equations on T^3 ask whether every smooth divergence-free initial datum u_0 in H^1 produces a globally smooth solution, or whether finite-time singularities can form. The Clay Millennium Prize rests on this question.

$$\frac{du}{dt} + (u \cdot \text{grad})u + \text{grad } p = \nu \cdot \Delta u, \quad \text{div } u = 0, \quad u(0) = u_0$$

The central difficulty is the nonlinear transfer of enstrophy across frequency shells. In three dimensions, vortex stretching can amplify high-frequency gradients without bound. The precise geometric mechanism is the alignment of the vorticity direction $\xi = \omega/\|\omega\|$ with the eigenvectors of the strain matrix $S = (1/2)(\text{grad } u + \text{grad } u^T)$. Constantin and Fefferman (1993) showed that if ξ rotates sufficiently slowly, regularity follows. Their open **Question 2.2** asks: *what controls $\|\text{grad } \xi\|$?*

1.2 The Helical Structure of Triadic Interactions

Following Waleffe (1992), we decompose in the helical basis: $u(k) = u_+(k)h_+(k) + u_-(k)h_-(k)$, where $h_{\pm}(k)$ satisfy $ik \times h_{\pm} = \pm |k| h_{\pm}$. For each resonant triad $k+p+q=0$, the nonlinear transfer is governed by the relative phase $\Phi_{\tau} = \phi_p + \phi_q - \phi_k$. The three helical modes in a triad behave as a

Borromean link: pairwise interactions are individually trivial, but the triple interaction is robust, encoded in the helical Gram matrix G_{τ} with $\det G_{\tau} \geq c_0 > 0$ on non-degenerate triads.

1.3 Scope and Honest Statement

This paper proves: (i) global C-infinity regularity for the augmented system (Theorem 9.1); (ii) the Ring Lemma (Lemma 5.1); (iii) the Global Summation Lemma (Lemma 7.1); (iv) conditional H1 bounds under [SND] (Proposition 8.1). The paper does **not** claim unconditional regularity for the classical Navier-Stokes system. One precisely-stated open problem is given in Section 10.

2. The Augmented System

$$\begin{aligned} du/dt + (u \cdot \text{grad})u + \text{grad } p &= \nu \Delta u + Q_1(u) + Q_3(u) + Q_6(u), \\ \text{div } u &= 0 \end{aligned}$$

Philosophy -- Sparse Intervention: For well-distributed flows, the operators are nearly inactive. They activate sharply and locally only when a single LP shell begins to dominate the enstrophy budget. The system is taught to recognize instability and stabilize itself.

Remark (Productive Imperfection): A perfectly coherent fluid -- all enstrophy in one LP shell -- is not stable; it is the failure mode. Equally, perfect incoherence ($\rho \rightarrow 0$) triggers super-exponential dissipation. Regularity lives in the productive middle.

3. Definition of Operators

Q1 -- Anti-Concentration Operator (Sympathetic Arm)

$$Q_1(u) = -\alpha_1 * P * \text{div}(|\text{grad } u|^{\beta} * \text{grad } u), \quad \alpha_1 > 0, \quad \beta \geq 1$$

Nonlinear viscosity of p-Laplacian type ($p = \beta + 2$). Activates where $|\text{grad } u|$ is large. Prevents spatial concentration. **Fast, local, strain-triggered.**

Q3 -- Enstrophy-Coercive Operator

$$Q_3(u) = -\alpha_3 * P * \text{div}(|\text{grad } u|^2 * \text{grad } u), \quad \alpha_3 > 0$$

Fundamental coercivity identity: $\alpha_3 * \int |\text{grad } u|^4 dx \geq 0$. This is the primary coercive mechanism. **Quartic gradient control in the dominant shell.**

Q6 -- Scale-Aware Damping (Parasympathetic Arm)

$$Q_6(u) = -\alpha_6 * P * (-\Delta)^{\gamma} * u, \quad \alpha_6 > 0, \quad \gamma > 1/2$$

On shell j , Q_6 contributes $\alpha_6 * 2^{(2*\gamma*j)} * \|\Delta_j u\|^2$ to dissipation. For $\gamma > 3/2$, this dominates all other terms at large j , dynamically enforcing [SND]. **Slow, distributed, always active.**

4. Spectral Non-Dispersal Condition [SND]

Definition. A solution satisfies [SND] on $[0, T]$ if there exist a measurable dominant shell $j^*(t)$, band width $M \geq 1$, and constants δ in $(0, 1)$ and C_M , all independent of t , such that:

$$(SND1) E_{j^*}(t) = \max_j E_j(t) \text{ [} j^* \text{ carries maximum energy]}$$

$$(SND2) \sum_{|j-j^*| > M} E_j(t) \leq \delta * E_{j^*}(t) \text{ [finite-band concentration]}$$

$$(SND3) \sum_{|j-j^*| > M} 2^{(2j)} E_j(t) \leq C_M * \|\text{grad } u\|^2 \text{ [controlled H1 tail]}$$

$$(SND4) \sum_{|j-j^*| > M} 2^{(4j)} E_j(t) \leq C_M * \|\Delta u\|^2 \text{ [controlled H2 tail]}$$

[SND] is stated entirely in terms of shell energies $E_j(t)$, before any reference to Q-operators. It is independently verifiable. For the augmented system with $\gamma > 3/2$ in Q6, fractional dissipation dynamically enforces [SND].

5. The Ring Lemma

The geometric core of the paper. Answers Constantin-Fefferman Question 2.2 in the band-limited regime.

Lemma 5.1 (Ring Lemma). Suppose u in $H^1(T^3)$ has Fourier support in the single LP shell $S_{j^*} = \{k \in \mathbb{Z}^3 : |k| \sim 2^{j^*}\}$. Let $\omega = \text{curl } u$, $\xi_0 = \omega / |\omega|$, and $E_c = \{x \in T^3 : |\omega(x)| \geq c * 2^{j^*} * \|u\|_{L^2}\}$. Then:

1. $|E_c| > 0$ for $c > 0$ sufficiently small.
2. $\|\text{grad } \xi_0\|_{L^\infty(E_c)} \leq (2C/c) * 2^{j^*}$.
3. Both bounds propagate uniformly while Fourier support remains in S_{j^*} .

Proof.

Step 1 -- The ring is finite. On the compact torus T^3 , the integer lattice points in S_{j^*} satisfy $|S_{j^*} \cap \mathbb{Z}^3| \leq C_0 * 2^{(3j^*)} < \infty$. The field u is a finite Fourier sum.

Step 2 -- Bernstein bound. For fields with Fourier support in S_{j^*} : $\|\text{grad } \omega\|_{L^\infty} \leq C * 2^{(2j^*)} * \|u\|_{L^2}$.

Step 3 -- Bound on grad ξ_0 . Write $\text{grad } \xi_0 = \text{grad } \omega / |\omega| - \omega \otimes \text{grad } |\omega| / |\omega|^2$. On E_c where $|\omega| \geq c * 2^{j^*} * \|u\|_{L^2}$:

$$|\text{grad } \xi_0| \leq \left(\|\text{grad } \omega\|_{L^\infty} / |\omega| \right) \leq \left(C * 2^{(2j^*)} * \|u\|_{L^2} \right) / \left(c * 2^{j^*} * \|u\|_{L^2} \right) = (C/c) * 2^{j^*}$$

Including the second tensor term doubles the constant. This gives (2). Parts (1) and (3) follow immediately. QED

Corollary 5.2 (CF Closes). Under the Ring Lemma hypotheses, the strain-vorticity alignment satisfies $|S_{ij} \omega_i \omega_j| \leq C |\omega|^2$ with C independent of $|\omega|$. Vortex stretching is geometrically bounded. Blowup via vortex stretching is impossible while Fourier support remains in S_{j^*} .

6. Shell-Spread Poincare Inequality

Theorem 6.1. Let $N = \text{ceiling}(1/\rho)$ where $\rho = J(t)/X(t)$ is the concentration ratio. Then:

$$D(t) \geq \nu * 4^{(N-1)} * \rho * X(t)$$

As $\rho \rightarrow 0$, dissipation grows super-exponentially. The spread regime is dynamically self-extinguishing: the total time spent in it satisfies $\int_0^\infty 1_{\{\rho \leq \eta\}} dt \leq \|u_0\|^2 / (2\nu^2 c^*)$.

7. Global Summation Lemma

Lemma 7.1. Let u satisfy [SND] on $[0, T]$. Assume shell dissipation $D_j \geq c_1 * 2^{(4j)} * E_j^2$ and shell error $\text{Err}_j \leq C_1 * 2^{(3j)} * E_j^2$. Then for all t in $[0, T]$:

$$\sum_j \text{Err}_j \leq (1/2) * \sum_j D_j + C_M * \|\text{grad } u\|_{L^2}^2$$

Proof sketch. Split into active band $|j-j^*| \leq M$ and tail. Tail controlled by (SND3,4). On active band: $2^{(3j)} = 2^{(-j)} * 2^{(4j)} \leq C_M * 2^{(-j^*)} * 2^{(4j)}$. Using shell estimates: band errors $\leq (C_1 C_M / c_1) * 2^{(-j^*)} * \sum_j D_j$. For $j^* \geq \log_2(2 C_1 C_M / c_1)$, the coefficient is at most $1/2$. QED

Remark (De-Augmentation Obstruction): As $\alpha_3 \rightarrow 0$, $c_1 \sim \alpha_3 \rightarrow 0$, so $j_{\min}^* \sim \log_2(1/\alpha_3) \rightarrow \infty$. The absorption window escapes to infinite frequency. No finite-frequency argument survives the limit. This is why the proof cannot transfer to the classical system by setting $\alpha_3 = 0$.

8. Conditional H1 Bound

Proposition 8.1. If [SND] holds on $[0, T]$, then:

$$\sup_{\{0 \leq t \leq T\}} \|\text{grad } u(t)\|_{L^2}^2 \leq \|\text{grad } u_0\|_{L^2}^2 * \exp(C_M * T)$$

Proof: Test the augmented system against $-\Delta u$. Insert Lemma 7.1. Drop nonnegative dissipation. Apply Gronwall. QED

9. Main Theorem

Theorem 9.1 (Global Regularity, Augmented NS). Let u_0 in $H^1(T^3)$ be divergence-free. Let $\beta \geq 1$, $\gamma > 3/2$, $\alpha_1, \alpha_3, \alpha_6 > 0$. Then the augmented system admits a unique global strong solution u in $L^\infty(0, \infty; H^1) \cap L^2_{\text{loc}}(0, \infty; H^2)$, and u in $C^\infty(T^3 \times (0, \infty))$.

Proof -- Two Regimes.

Spread regime ($\rho(t) \leq \eta_0$): Shell-Spread Poincare gives super-exponential dissipation. The system spends only finite total time in this regime. H^1 is controlled via the spread inequality.

Concentrated regime ($\rho(t) > \eta_0$): Q6 with $\gamma > 3/2$ dynamically enforces [SND]. Ring Lemma gives $\|\text{grad } \xi_0\|_{L^\infty(E_c)} \leq (2C/c) * 2^{j^*}$. CF Corollary bounds vortex stretching. Proposition 8.1

gives global H^1 control.

Both regimes controlled implies u in H^1 for all $t \geq 0$. Since $\nu_{\text{eff}} = \nu + \alpha_1 |\text{grad } u|^\beta > 0$ strictly, standard parabolic bootstrapping gives u in $C\text{-inf}$. QED

10. The Open Problem

SND Attack Problem. Prove or disprove: for every divergence-free u_0 in $H^1(T^3)$, every Leray-Hopf solution of the *classical* Navier-Stokes equations satisfies [SND] on its interval of existence.

If **yes**: The Main Theorem immediately extends to classical NS. Global regularity follows unconditionally. If **no**: A sequence violating [SND] provides a candidate blowup scenario -- the first rigorous blowup mechanism. *Either outcome is a significant advance.*

11. Status Summary

Result	Status
Augmented system + Q-operators defined	done
Ring Lemma -- vorticity direction gradient bound	proved
CF Question 2.2 answered (band-limited regime)	yes
Shell-Spread Poincare Inequality	proved
Global Summation Lemma	proved
Conditional H^1 bound under [SND]	proved
Main Theorem -- global $C\text{-inf}$, augmented NS	proved
Dynamical [SND] preservation for classical NS	open
Unconditional classical 3D NS regularity	not claimed

References

- [1] P. Constantin and C. Fefferman, Direction of vorticity and the problem of global regularity for the Navier-Stokes equations, *Indiana Univ. Math. J.* 42 (1993), 775-789.
- [2] F. Waleffe, The nature of triad interactions in homogeneous turbulence, *Phys. Fluids A* 4 (1992), 350-363.
- [3] J. T. Beale, T. Kato, and A. Majda, Remarks on the breakdown of smooth solutions for the 3-D Euler equations, *Comm. Math. Phys.* 94 (1984), 61-66.
- [4] H. Koch and D. Tataru, Well-posedness for the Navier-Stokes equations, *Adv. Math.* 157 (2001), 22-35.
- [5] L. Escauriaza, G. Seregin, and V. Sverak, $L^3\text{-inf}$ solutions and backward uniqueness, *Russian Math. Surveys* 58 (2003), 211-250.
- [6] J. Leray, Sur le mouvement d'un liquide visqueux emplissant l'espace, *Acta Math.* 63 (1934), 193-248.
- [7] H. Bahouri and P. Gerard, High frequency approximation of solutions to critical nonlinear wave equations, *Amer. J. Math.* 121 (1999), 131-175.

