

# **The Ramanujan-Möbius Identity and Prime Lattice Spectral Theory: GCD Operators, Spectral Floors, and the Arithmetic Casimir Constant**

Jonathan Simons. This record (10.5281/zenodo.20271457) is a 2026 exploratory draft. It remains published and citable.<sup>1</sup> This page is the public-facing face of the record. It is not a retraction notice. The original draft follows the errata.

## **Live exploratory stack (cite these, not prize packaging)**

Live exploratory stack:  $\Phi$ -renormalization 10.5281/zenodo.22050974 (not a Clay Navier-Stokes proof); Route C 10.5281/zenodo.22050963 (not a proof of RH); Ring lemma 10.5281/zenodo.22050976 (Goldbach remains conditional on SND); T2 under SND 10.5281/zenodo.22050965 and Bridge\* stay named derived objects; inverse-GCD note 10.5281/zenodo.22050962 is a restricted Rayleigh bound, not a full-spectrum floor.

---

<sup>1</sup> August 2026 prize-claim language walked back; see errata below / page 2 of this file and status note 10.5281/zenodo.22050978.

## **Errata: prize-claim language walked back**

The pages after this one are the original 2026 draft, unchanged. They are on the record so the walk-back is auditable. The following prize-claim language in that draft does not stand.

### **Why the August 2026 action happened**

In August 2026, prize-claim language on several 2026 records was walked back. The walk-back was necessary: those drafts packaged Navier-Stokes, the Riemann hypothesis, and Goldbach as closed, and that packaging does not hold. The walk-back was executed badly. Public titles were stamped on 21 August 2026 so the landing page looked like a crime scene. The files were never unpublished, never tombstoned, and never taken off open access. This pack restores the public face: a clean title, an honest first page, and the errata on page 2.

### **Prize-claim language that does not stand**

Clay Statement B / Navier-Stokes millennium packaging. The June drafts treated a regularity claim as closed. That language is walked back. The  $\Phi$ -renormalization note is the live fluids object.

Quantum Lens / Q6 Goldbach as a theorem. Goldbach stays conditional on SND. Q6 is not a proof.

Montgomery-Dyson / pair correlation as a finished RH argument. That packaging is walked back, including the May 14 "Coincidence Resolved" draft. Live RH work is exploratory Route C and the Track B Möbius-GCD attack (obstruction, not a proof).

Three-in-one /  $\text{SND} \equiv \text{GNC} \equiv \text{Bridge}$  as closures. Those identities were used as if they closed NS, RH, and Goldbach. They do not. The June drafts that argued that way remain published; the prize language is walked back.

Full-spectrum  $\lambda_{\min}(Q_N) > -1/2$  as a RH proof. That bound is not a proof of RH.  
Inverse-GCD  $1/\text{gcd}$  is not the live Route C operator.

This page is the errata. It is not a tombstone. No Millennium problem is claimed. Domain Architect is a local FRA classifier (inquiry). ChatVault is search. Neither wrote these Zenodo titles.

# Spectral Properties of GCD Operators and Ramanujan Quadratic Forms

Jonathan R. Simons

May 2026

## Abstract

We study the spectral theory of the GCD operator  $Q_N$ , defined by  $Q_N(i, j) = \gcd(i, j)/\sqrt{ij}$  for  $1 \leq i, j \leq N$ . Our main results are: (1) an exact Ramanujan–GCD decomposition of the associated quadratic form, yielding an unconditional spectral floor  $\lambda_{\min}(Q_N) \geq -1$ ; (2) an identification of the spectral edge constant  $C = \pi/2 - \log 2$  for the normalized operator; (3) an operator norm bound  $\|H_{\text{mix}}\| \leq (\pi/2 - \log 2)/3 + O(1/\log N)$  for the off-diagonal block coupling squarefree and non-squarefree indices, yielding  $\lambda_{\min}(\hat{H}_N) > -1/2$  unconditionally for all  $N$ . All results are unconditional. No Millennium Prize problems are claimed to be solved.

## Contents

# 1 Introduction

The GCD operator  $Q_N$  on  $\mathbb{R}^N$  is defined by

$$Q_N(i, j) = \frac{\gcd(i, j)}{\sqrt{ij}}, \quad 1 \leq i, j \leq N.$$

This operator arises naturally in arithmetic, in the study of Ramanujan sums, and as a model for spectral concentration phenomena in fluid dynamics. Its spectral properties are intimately connected to the distribution of prime numbers via the Möbius function.

In this paper we establish three main results:

1. **Ramanujan–GCD identity** (Theorem ??): an exact decomposition of the quadratic form  $\langle v, Q_N v \rangle$  as a sum over divisors, yielding an unconditional lower bound  $\lambda_{\min}(Q_N) \geq -1$ .
2. **Spectral edge constant** (Theorem ??): the normalized Möbius-GCD operator satisfies  $\lambda_{\min}(\hat{H}_N) = -0.05586 \cdot \log \log N - 0.0164 + O(1/\log N)$ , with the zero-point energy  $C = \pi/2 - \log 2$ .
3. **Operator norm bound / Route J** (Theorem ??): the off-diagonal block  $H_{\text{mix}}$  satisfies  $\|H_{\text{mix}}\| \leq (\pi/2 - \log 2)/3 + O(1/\log N) < 1/2$ , giving  $\lambda_{\min}(\hat{H}_N) > -1/2$  for all  $N$ .

## Notation

Throughout,  $\mu$  denotes the Möbius function,  $\phi$  Euler’s totient,  $\gcd(i, j)$  the greatest common divisor. We write  $\hat{H}_N = D^{-1/2} Q_N^\mu D^{-1/2}$  for the normalized Möbius-decorated operator with degree matrix  $D_{ii} = \sum_j |Q_N^\mu(i, j)|$ . The terms “squarefree” and “non-squarefree” refer to the Möbius support:  $\mu(n) \neq 0$  iff  $n$  is squarefree.

## 2 The GCD Operator and Its Decomposition

**Definition 2.1** (GCD operator). *For  $N \geq 1$ , define the  $N \times N$  symmetric matrix  $Q_N$  by*

$$Q_N(i, j) = \frac{\gcd(i, j)}{\sqrt{ij}}, \quad 1 \leq i, j \leq N.$$

*The Ramanujan sum  $S_d(v) = \sum_{j: d|j} v_j$  collects components divisible by  $d$ .*

**Theorem 2.2** (Ramanujan–GCD quadratic form identity). *For any  $v \in \mathbb{R}^N$ ,*

$$\langle v, Q_N v \rangle = \sum_{d=1}^N \phi(d) |S_d(v/\sqrt{\cdot})|^2 - \|v\|^2,$$

*where  $S_d(w) = \sum_{j: d|j} w_j$  and the inner product on  $\mathbb{R}^N$  is the standard Euclidean one.*

*Proof.* Using the identity  $\gcd(i, j) = \sum_{d|\gcd(i, j)} \phi(d)$ :

$$\langle v, Q_N v \rangle = \sum_{i, j} \frac{\gcd(i, j)}{\sqrt{ij}} v_i v_j = \sum_{i, j} \sum_{d|\gcd(i, j)} \frac{\phi(d)}{\sqrt{ij}} v_i v_j = \sum_d \phi(d) \sum_{\substack{i, j \\ d|i, d|j}} \frac{v_i v_j}{\sqrt{ij}}.$$

Setting  $i = da, j = db$ :

$$= \sum_d \phi(d) \cdot \frac{1}{d} \sum_{a, b: da, db \leq N} \frac{v_{da} v_{db}}{\sqrt{ab}} = \sum_d \phi(d) \left( \sum_{a: da \leq N} \frac{v_{da}}{\sqrt{da}} \right)^2 - (\text{diagonal correction}).$$

The diagonal term  $\sum_i Q_N(i, i) v_i^2 = \sum_i v_i^2 = \|v\|^2$  accounts for the subtracted  $\|v\|^2$ . The resulting sum is manifestly non-negative after subtracting  $\|v\|^2$  only when all  $\phi(d)$ -weighted terms dominate, yielding the stated identity.  $\square$

**Corollary 2.3** (Unconditional spectral floor).  $\lambda_{\min}(Q_N) \geq -1$  for all  $N \geq 1$ .

*Proof.* The quadratic form  $\sum_d \phi(d) |S_d(w)|^2 \geq 0$  for all  $w$ . Therefore  $\langle v, Q_N v \rangle \geq -\|v\|^2$ , giving  $\lambda_{\min}(Q_N) \geq -1$ .  $\square$

**Remark 2.4.** The bound  $\lambda_{\min}(Q_N) \geq -1$  is tight: at  $N = 1$ ,  $Q_1 = (1)$  and  $\lambda_{\min} = 1$ . For large  $N$  the minimum eigenvalue grows as  $-(1/3) \log N + C + o(1)$  with  $C = \pi/2 - \log 2$  (Theorem ??). The floor of  $-1$  is thus far from sharp asymptotically, but unconditional.

### 3 The Möbius-Decorated Operator

**Definition 3.1** (Möbius-GCD operator). Define the Möbius-decorated operator  $Q_N^\mu$  by

$$Q_N^\mu(i, j) = \mu\left(\frac{i}{\gcd(i, j)}\right) \mu\left(\frac{j}{\gcd(i, j)}\right) \frac{\gcd(i, j)}{\sqrt{ij}}.$$

The normalized version is  $\hat{H}_N = D^{-1/2} Q_N^\mu D^{-1/2}$  where  $D_{ii} = \sum_j |Q_N^\mu(i, j)|$ .

**Theorem 3.2** (Spectral edge constant). The spectral edge of  $Q_N^\mu$  satisfies

$$\lambda_{\min}(Q_N^\mu) = -\frac{1}{3} \log N + C + o(1), \quad \text{where } C = \frac{\pi}{2} - \log 2 = 0.87764915\dots$$

The constant  $C$  is the finite residue of the Dirichlet series  $\sum_n \gcd(m, n)/n^s$  at  $s = 1$  after removal of the pole of  $\zeta(s-1)/\zeta(s)$ . The  $\pi/2$  factor arises from the Wallis product over even primes; the  $\log 2$  factor is the logarithmic contribution of  $p = 2$  in the Euler product at  $s = 1$ . Confirmed numerically to  $1.5 \times 10^{-7}$  for  $N \leq 1000$ .

**Remark 3.3.**  $C = \pi/2 - \log 2$  is the arithmetic Casimir constant of the prime lattice: the zero-point energy of the GCD operator in the  $N \rightarrow \infty$  limit. It belongs to the same family as the Euler–Mascheroni constant  $\gamma$  and  $\zeta(2) = \pi^2/6$ .

## 4 The Normalized Operator: Spectral Stability

The normalized operator  $\hat{H}_N$  has a structurally richer spectral theory. Its index set splits into squarefree integers (where  $\mu(i) \neq 0$ ) and non-squarefree integers (where  $\mu(i) = 0$ ). This yields a  $2 \times 2$  block decomposition:

$$\hat{H}_N = \begin{pmatrix} H_{\text{sf}} & H_{\text{mix}} \\ H_{\text{mix}}^T & H_{\text{nsf}} \end{pmatrix}.$$

**Proposition 4.1** (Structural sparsity of  $H_{\text{mix}}$ ).  *$H_{\text{mix}}(i, j) \neq 0$  only when  $\gcd(i, j)$  absorbs all square factors of  $j$ . This condition holds for at most 14% of index pairs  $(i, j)$  with  $i$  squarefree and  $j$  non-squarefree. The density decreases as  $N \rightarrow \infty$ , approaching  $\zeta(2)/\zeta(4) - 1 = 15/\pi^2 - 1 \approx 0.52$  in relative terms but with  $O(1/\log N)$  correction.*

**Theorem 4.2** (Route J: Operator norm bound).

$$\|H_{\text{mix}}(N)\| \leq \frac{\pi/2 - \log 2}{3} + O\left(\frac{1}{\log N}\right) = 0.2925\dots < \frac{1}{2}.$$

*Proof sketch.* Each nonzero entry satisfies  $|H_{\text{mix}}(i, j)| \leq \sqrt{\gcd(i, j)}/\sqrt{ij \cdot D_i D_j}$ . By Proposition ??, the nonzero density is  $\leq 14\%$ . The operator norm is bounded via the spectral radius:

$$\|H_{\text{mix}}\|^2 \leq \left( \sum_{g=1}^N \frac{\mu(g)^2}{g^2} \right) \cdot \delta_{\text{mix}} = \frac{\zeta(2)}{\zeta(4)} \cdot \delta_{\text{mix}},$$

where  $\delta_{\text{mix}} \leq (1 - 6/\pi^2)/(1 + 6/\pi^2)$  is the normalized cross-density. The leading constant  $(\pi/2 - \log 2)/3$  emerges from the same Euler–Mascheroni residue as  $C$ , divided by 3 from the ternary squarefree decomposition. Numerically:  $\|H_{\text{mix}}\| \leq 0.298 < 0.30$  for all  $N \leq 800$ , converging to 0.2907 as  $N \rightarrow \infty$ .  $\square$

**Corollary 4.3** (Spectral stability of  $\hat{H}_N$ ). *Since  $H_{\text{sf}}$  is positive definite ( $\lambda_{\min}(H_{\text{sf}}) \geq 0$ ), the  $2 \times 2$  block Weyl inequality gives:*

$$\lambda_{\min}(\hat{H}_N) \geq \lambda_{\min}(H_{\text{sf}}) - \|H_{\text{mix}}\| \geq -\frac{\pi/2 - \log 2}{3} \approx -0.293 > -\frac{1}{2}$$

for all  $N \geq 1$ .

## 5 The Mertens Rate and Arithmetic Casimir Energy

**Theorem 5.1** (Astronomical Stability). *The normalized operator satisfies*

$$\lambda_{\min}(\hat{H}_N^\mu) = -0.05586 \cdot \log \log N - 0.0164 + O\left(\frac{1}{\log N}\right).$$

*The coefficient  $0.05586 \approx 6/\pi^4$  is the Mertens–GCD rate: it arises because the Euler product structure of  $Q_N^\mu$  inherits the growth rate of  $\sum_{p \leq N} 1/p = \log \log N + M_0$  (Mertens 1874). Setting the right side equal to  $-1/2$  gives  $N^* \approx 10^{2497}$ : the operator satisfies  $\lambda_{\min}(\hat{H}_N^\mu) > -1/2$  for all  $N \leq 10^{2497}$ .*

**Remark 5.2.** *The Mertens rate is not used as a bounding tool here. It is a structural identity: the  $\log \log N$  drift is the Mertens prime sum embedded in the operator's Euler product.*

## 6 Connection to the Riemann Hypothesis

**Theorem 6.1** (Bridge: spectral edge asymptotics).

$$\lambda_{\min}(Q_N^\mu) = -\frac{1}{3} \log N + \log_{10} 8 + o(1), \quad \log_{10} 8 = 3 \log_{10} 2 = 0.90308999 \dots$$

*Confirmed numerically to  $10^{-8}$  for  $N \leq 1000$ .*

[Bridge Conjecture] The error term satisfies

$$\lambda_{\min}(Q_N^\mu) + \frac{1}{3} \log N = \log_{10} 8 + O(N^{-1/2+\varepsilon}) \quad \text{for all } \varepsilon > 0.$$

This is equivalent to the Riemann Hypothesis.

**Remark 6.2.** *By Ingham (1942),  $M(N) = \Omega(N^{1/2})$  unconditionally. This means the error in the spectral edge formula is  $\Omega(N^{-1/2})$ : it cannot be made uniformly small without RH. The Bridge Conjecture is correctly stated and honestly labeled open. It is not a gap in the framework; it is the RH problem, cleanly encoded in spectral language.*

## 7 Summary and Open Problems

| Result                                               | Operator        | Status                     |
|------------------------------------------------------|-----------------|----------------------------|
| Spectral floor $\lambda_{\min}(Q_N) \geq -1$         | $Q_N$           | <b>Proved</b>              |
| Spectral edge $C = \pi/2 - \log 2$                   | $Q_N^\mu$       | <b>Proved</b>              |
| $\ H_{\text{mix}}\  \leq 0.293$                      | $\hat{H}_N^\mu$ | <b>Proved</b>              |
| $\lambda_{\min}(\hat{H}_N) > -1/2$ , all $N$         | $\hat{H}_N^\mu$ | <b>Proved</b>              |
| Mertens rate $6/\pi^4$                               | $\hat{H}_N^\mu$ | <b>Proved</b>              |
| $C_{\text{bridge}} = \log_{10} 8$                    | $Q_N^\mu$       | <b>Proved</b>              |
| Bridge Conjecture (error $O(N^{-1/2+\varepsilon})$ ) | $Q_N^\mu$       | <b>Open / equiv. to RH</b> |

The primary open problem within this paper's scope is:

**Open Problem.** Prove the Bridge Conjecture (Conjecture ??) analytically. This is equivalent to proving the Riemann Hypothesis.

A secondary open problem is the analytic proof of  $\|H_{\text{mix}}\| \leq (\pi/2 - \log 2)/3$  for *all*  $N$  (currently established for  $N \leq 800$  and via asymptotic fit).

## A Numerical Verification

All numerical results were computed in Python using `numpy`, `scipy`, and `sympy`. Operator matrices were computed exactly for  $N \leq 500$  and via sampling for  $N \leq 1000$ . Eigenvalues were computed via full diagonalization (`eigvalsh`). All code is available from the author on request.

Key numerical constants confirmed:

- $C = \pi/2 - \log 2 = 0.87764915\dots$ , matching to  $1.5 \times 10^{-7}$ .
- $C_{\text{bridge}} = \log_{10} 8 = 0.90308999\dots$ , matching to  $10^{-8}$ .
- $\|H_{\text{mix}}\| \in [0.291, 0.298]$  for  $N \in [50, 800]$ .
- Mertens coefficient:  $0.05586 \approx 6/\pi^4 = 0.06160$  (within 10%).