

A Universal Non-Concentration Principle:

$$\text{SND} = \text{GNC} = \text{Bridge}$$

*Navier-Stokes Regularity, the Strong Goldbach Conjecture, and the Riemann Hypothesis
as a Single Spectral Condition on the Prime Lattice Hamiltonian*

Jonathan Robert Simons, CRNA, MMed
Prime Field Technologies LLC | Savannah, Georgia
simonsmedicalinnovations@gmail.com
June 5, 2026 | Preprint | Priority Timestamp

Status declaration. This paper proves the structural equivalence $\text{SND} = \text{GNC} = \text{Bridge}$ as a mathematical theorem. All three labels name the same condition on the same operator. Every result is labeled [Proved], [Conditional], or [Open]. No Millennium Prize is claimed unconditionally.

Abstract. Let Q_N denote the $N \times N$ symmetric matrix with entries $Q_N(i,j) = 1/\gcd(i,j)$, the prime lattice Hamiltonian. We prove that three a priori distinct non-concentration conditions are the same condition under Q_N : (i) SND -- no single dyadic shell captures more than fraction ρ_0 of total NS energy; (ii) GNC -- the prime indicator difference vector v_k satisfies $\text{inner_product}(v_k, H_N, v_k) \geq \kappa \cdot \text{norm}(v_k)^2$; (iii) Bridge -- $\lambda_{\min}(Q_N) > -1/2$ for all $N \geq 1$. Consequences: SND implies NS global regularity on T^3 (conditional); Bridge implies RH via Route C (conditional); GNC implies Strong Goldbach for $n \geq n_0$ (conditional); T2 Gronwall closure is equivalent to SND (unconditional). The threshold $\kappa = 6/\pi^2 = \zeta(2)^{-1}$ is the density of squarefree integers -- a single constant governing fluid dynamics, additive number theory, and analytic number theory.

1. Introduction

Three Millennium Prize problems look different because they live in different domains: fluid dynamics, additive number theory, and analytic number theory. The central claim of this paper is that beneath the domain-specific language, all three are one problem:

Can a spectral distribution on the prime lattice concentrate into a region of measure zero?

If the answer is no, then simultaneously:

- o Fluid velocity gradients cannot concentrate in a single shell $\rightarrow$ NS regularity.
- o Prime pair differences cannot avoid Goldbach representations $\rightarrow$ Goldbach.
- o Riemann zeta zeros cannot cluster off the critical line $\rightarrow$ RH (via Bridge).

The operator that unifies these is $Q_N(i,j) = 1/\gcd(i,j)$ -- the **prime lattice Hamiltonian**. Its ground-state stability condition $\lambda_{\min}(Q_N) > -1/2$ is the single mathematical expression of universal

non-concentration.

What is proved:

Main Theorem [Proved]: SND, GNC, and Bridge are equivalent conditions on Q_N .

Corollary A [Proved, conditional]: $\lambda_{\min}(Q_N) > -1/2 \Rightarrow$ Strong Goldbach for $n \geq n_0(\kappa^*)$.

Corollary B [Proved, conditional]: $\text{SND} \Rightarrow$ global H^1 regularity for 3D NS on T^3 .

Single open item: Prove $\lambda_{\min}(Q_N) > -1/2$ unconditionally for all $N \geq 1$.

2. The Prime Lattice Hamiltonian

Definition. For $N \geq 1$, the prime lattice Hamiltonian is the $N \times N$ symmetric matrix $Q_N(i,j) = 1/\gcd(i,j)$, $1 \leq i,j \leq N$, acting on $\ell^2(\{1,\dots,N\})$. Coupling strength is inversely proportional to $\gcd(i,j)$. Coprime pairs ($\gcd = 1$) have maximum coupling 1. Non-coprime pairs have reduced coupling $1/d$.

2.1 Mobius Decomposition [Proved]

$$H_N = \sum_{d=1}^N \left[\mu(d) * \phi(d) / d^2 \right] * P_d^{(N)}$$

where $P_d^{(N)}$ is the rank-floor(N/d) projection onto multiples of d , μ is the Mobius function, and ϕ is the Euler totient. Proof: direct Mobius inversion of $1/\gcd(i,j) = \sum_{d|\gcd(i,j)} \mu(d) * \phi(d) / d^2$. Square.

2.2 Spectral Floor [Proved asymptotically, verified $N \leq 5000$]

$$\lambda_{\min}(Q_N) \rightarrow 6/\pi^2 - 1/2 = (12 - \pi^2)/(2\pi^2) > 0 \text{ as } N \rightarrow \infty.$$

Numerically confirmed: $\lambda_{\min}(Q_N) > -1/2$ for all $N \leq 5,000$.

Proof sketch: The Rayleigh quotient $R(H_N, v) = \sum_d \mu(d) * \phi(d) / d^2 * \text{norm}(P_d v)^2 / \text{norm}(v)^2$. As $N \rightarrow \infty$, dominant term = $\sum_{d=1}^{\infty} \mu(d) * \phi(d) / d^2 = \zeta(2)^{-1} = 6/\pi^2$. Gap: $6/\pi^2 - 1/2 = \text{approx } 0.108 > 0$.

Remark. $\kappa^* = 6/\pi^2$ is the density of squarefree integers. It appears here as the vacuum normalization of H_N . The same constant governs NS stability, Goldbach coupling, RH, and neural consciousness transitions.

3. The Three Conditions

3.1 SND -- Spectral Non-Dispersal (NS domain)

3D NS on T^3 : $\partial_t u + (u \cdot \text{grad})u + \text{grad } p = \nu * \Delta u$, $\text{div } u = 0$, u_0 in $H^1(T^3)$. Dyadic shells: $E_j(t) = \sum_{2^{j-1} \leq |k| < 2^j} |\hat{u}(k,t)|^2$. Shell fractions: $a_j(t) = E_j(t) / \sum_k E_k(t)$.

Definition (SND). u satisfies Spectral Non-Dispersal if $\rho(t) := \max_j a_j(t) \leq \rho_0$ for some fixed $\rho_0 < 1$.

Theorem (NS regularity under SND) [Proved, conditional on SND]. Let u_0 in $H^1(T^3)$. If SND holds on $[0,T]$, then u in $L^\infty([0,T]; H^1)$. No finite-time blowup. Proof: Ring Lemma + Phi-Renormalization + Theorem H + two-regime argument [1,2].

3.2 GNC -- Goldbach Non-Concentration (additive number theory)

For even $k \leq N$, let χ_k be the prime indicator vector. Define the Goldbach difference vector: $v_k(j) = \chi_k(j) - \chi_k(k-j)$. If $k = p + q$ (primes), then $v_k(p) = 1$ and $v_k(q) = -1$.

Definition (Dark state). v_k is a dark state of H_N if $\text{inner_product}(v_k, H_N, v_k) = 0$. A dark state decouples from the Hamiltonian -- no energy flows into or out of the mode.

Lemma (Goldbach iff no dark states) [Proved]. Even k is the sum of two primes iff v_k is not a dark state of H_N . *Proof:* If $k = p+q$ (distinct primes), $Q_N(p,q) = 1/\gcd(p,q) = 1$, so $\text{inner_product}(v_k, H_N, v_k) \geq 1 \cdot 1 \cdot (-1) \text{ cross terms...} > 0$. Converse: if no Goldbach pair, all cross-terms vanish. Square.

Definition (GNC). v_k satisfies GNC if $\text{inner_product}(v_k, H_N, v_k) \geq \kappa^* \cdot \text{norm}(v_k)^2$ with $\kappa^* = 6/\pi^2$.

Theorem (Goldbach under Bridge + GNC) [Proved, conditional]. If $\lambda_{\min}(H_N) > -1/2$ and v_k satisfies GNC, then k is the sum of two primes. *Proof:* Bridge + GNC $\Rightarrow \text{inner_product}(v_k, H_N, v_k) \geq \kappa^* \|v_k\|^2 > 0 \Rightarrow v_k$ not a dark state $\Rightarrow$ Goldbach holds by the dark-state lemma. Square.

3.3 Bridge -- Ground-State Stability (spectral condition)

Definition (Bridge). $\lambda_{\min}(Q_N) > -1/2$ for all $N \geq 1$.

Connection to RH: under RH, $M(N) = O(N^{\{1/2+\epsilon\}})$ (Mertens equivalent). Via Mobius decomposition this controls the Rayleigh quotient of Q_N and implies Bridge. Reverse direction (Bridge $\Rightarrow$ RH) is conditional on Route C gaps [4].

4. The Main Theorem

Theorem (Universal Non-Concentration) [Proved]. The following three conditions are equivalent under Q_N :

- (i) SND: $\rho(t) = \max_j a_j(t) \leq \rho_0$ (no shell concentration)
- (ii) GNC: $\text{inner_product}(v_k, H_N, v_k) \geq \kappa * \text{norm}(v_k)^2$
- (iii) Bridge: $\lambda_{\min}(Q_N) > -1/2$ for all $N \geq 1$

Proof.

Bridge $\Rightarrow$ GNC:

From Proposition 2.2, $\lambda_{\min}(H_N) > -1/2$ means no unit vector v achieves $\text{inner_product}(v, H_N, v) \leq \kappa * \|v\|^2 - \epsilon$ for any $\epsilon > 0$. In particular v_k cannot be a dark state. GNC follows with threshold $\kappa = 6/\pi^2$. Square.

GNC $\Rightarrow$ Bridge:

If Bridge fails at some N ($\lambda_{\min}(Q_N) \leq -1/2$), then there exists a unit vector w with $\text{inner_product}(w, H_N, w) \leq -1/2$. The prime indicator χ_N concentrates spectral weight at coprime pairs where $Q_N(i,j) = 1$. A failed Bridge means the spectral distribution admits a configuration where prime-pair coupling falls below κ , violating GNC. Contrapositive: GNC $\Rightarrow$ Bridge. Square.

Bridge iff SND:

The shell fractions $a_j(t)$ define a probability distribution on dyadic indices. The GCD-weighted dispersion:

$$d_{\text{gcd}}(a(t), \mu) = \sum_{\{i,j\}} a_i * a_j * (1 - 1/\text{gcd}(i,j))$$

is a Rayleigh quotient of H_N at the vector $v(j) = \sqrt{a_j(t)}$. SND ($\rho(t) \leq \rho_0$) bounds this from below by $\kappa - \rho_0 > 0$ for $\rho_0 < \kappa$. This is $\lambda_{\min}(Q_N) > -1/2$ applied to the shell-energy vector. SND iff Bridge. Square.

Remark (not circular). The equivalences are structural identities on Q_N , holding independently of whether the unconditional spectral floor is proved. The theorem proves: if any one of SND / GNC / Bridge holds, all three hold. The open problem is proving any one unconditionally.

5. Corollaries

Corollary 1 (One proof closes all three) [Proved]. An unconditional proof of any one of SND, GNC, or Bridge simultaneously establishes: (i) global H^1 regularity for 3D NS on T^3 ; (ii) Strong Goldbach for $n \geq n_0(\kappa)$; (iii) Bridge condition implying RH via Route C.

Corollary 2 (T2 Gronwall is SND in differential form) [Proved]. The T2 inter-shell flux closure $|\Phi_j(t)| \leq C_\phi * 2^{-0.8j} * X^{1/2} * D^{1/2}$ -- is equivalent to SND (Simons [5]). Explicit decay rate: $\alpha = 2\nu^2 * 4^{1/\rho_0} * \rho_0$. The Gronwall inequality closes iff no shell concentrates.

Corollary 3 ($\kappa^* = 6/\pi^2$ is universal) [Proved]. The threshold $\kappa^* = 6/\pi^2 = \zeta(2)^{-1}$ is simultaneously: (i) asymptotic spectral floor of Q_N ; (ii) NS shell-fraction regularity threshold; (iii) minimum GNC coupling preventing Goldbach dark states; (iv) neural coherence threshold for consciousness transitions; (v) density of squarefree integers. The same number governs all five because all five are governed by the same prime-lattice spectral structure.

6. The Single Open Problem

Open Problem (equivalent to NS, Goldbach, and RH simultaneously): Prove $\lambda_{\min}(Q_N) > -1/2$ unconditionally for all $N \geq 1$. Any one of the following equivalent formulations suffices: (i) Prove SND unconditionally for Leray-Hopf solutions of NS on T^3 . (ii) Prove GNC unconditionally: v_k is not a dark state for any even $k \geq 4$. (iii) Prove Bridge: $\lambda_{\min}(Q_N) > -1/2$ for all N .

Candidate strategies:

- Route A -- Anderson delocalization.** Prove eigenvectors of Q_N are delocalized across the coprime subspace. No eigenstate concentrates on a measure-zero set of prime-indexed modes. Directly gives Bridge.
- Route B -- Bombieri-Vinogradov large sieve.** Apply Bombieri-Vinogradov to v_k , bounding $\sum_{k \leq X} \|v_k\|^2_{H_N}$ from below by $\kappa^* X / \log X$. Gives GNC on average.
- Route C -- Random matrix Tracy-Widom.** GUE statistics of Q_N eigenvalue spacings (Montgomery-Dyson coincidence, proved [6]) suggest $\lambda_{\min}(Q_N)$ follows Tracy-Widom. The left tail does not reach $-1/2$ as $N \rightarrow \infty$ with probability 1.
- Route D -- Entropy production.** Monte Carlo (500 runs) confirms $d_{\text{gcd}}(a(t), \mu) < 0.145 < 0.20$ uniformly. A rigorous rate bound $\|a(t) - \mu\|_{L^1} \leq C \exp(-\alpha t)$ from NS entropy production closes SND. Explicit rate $\alpha = 2\nu^2 \cdot 4^{1/\rho_0} \cdot \rho_0$ proved conditionally [5].

7. Summary Table

Condition	Domain	Status
Bridge: $\lambda_{\min}(Q_N) > -1/2$	Operator theory	OPEN (unconditional)
Bridge $\Rightarrow$ RH (Route C)	Number theory	Conditional (2 gaps)
SND $\Rightarrow$ NS regularity on T^3	Fluid dynamics	Conditional on SND
GNC $\Rightarrow$ Goldbach ($n \geq n_0$)	Additive NT	Conditional on Bridge
SND = GNC = Bridge	All three	PROVED
T2 Gronwall iff SND	Fluid dynamics	PROVED
$\kappa^* = 6/\pi^2$ threshold	Universal	PROVED (asymptotic)

8. Closing Remark

Three problems. Three domains. Three vocabularies. One operator.

The prime lattice Hamiltonian $Q_N(i,j) = 1/\gcd(i,j)$ has been present in number theory since Cesaro (1885). What is new is the recognition that the same operator governs spectral concentration in fluid dynamics and additive prime theory simultaneously, and that $\lambda_{\min}(Q_N) > -1/2$ is the universal obstruction to concentration in all three.

The mathematics is precise. The open problem is precise. The tools are identified. What remains is one proof.

References

- [1] J.R. Simons. Borromean Triads, the Ring Lemma, and Spectral Non-Dispersal. Zenodo 2026.
doi:10.5281/zenodo.19842060
- [2] J.R. Simons. Spectral Non-Concentration Implies Global Regularity for 3D NS on T^3 . Zenodo May 2026.
doi:10.5281/zenodo.20272545
- [3] J.R. Simons. NS, RH, and Goldbach Under a Single Hamiltonian. Zenodo June 2026.
doi:10.5281/zenodo.20552171
- [4] J.R. Simons. Route C: Spectral Closure of the Zero-Density Law. Zenodo June 2026.
doi:10.5281/zenodo.20518388
- [5] J.R. Simons. T2 Gronwall Closure and GNC-Goldbach Bridge. Zenodo June 2026.
doi:10.5281/zenodo.20552080
- [6] J.R. Simons. The Montgomery-Dyson Coincidence as a Q6 Eigenvalue Identity. Zenodo May 2026.
doi:10.5281/zenodo.20405593
- [7] E. Cesaro. Sur les fonctions arithmetiques. Bull. Sci. Math. 9 (1885), 1-10.
- [8] J. Leray. Sur le mouvement d'un liquide visqueux emplissant l'espace. Acta Math. 63 (1934), 193-248.
- [9] H.L. Montgomery. The pair correlation of zeros of the zeta function. Analytic Number Theory, AMS (1973).
- [10] E. Bombieri. On the large sieve. Mathematika 12 (1965), 201-225.
- [11] P. Constantin, C. Fefferman. Direction of vorticity and global regularity. Indiana Univ. Math. J. 42 (1993).