


Machine Learning Support Vector Machines (SVMs), Part 1

Slides adapted from David Sontag, who adapted from Luke Zettlemoyer, Vibhav Gogate, and Carlos Guestrin

Linear Separators


- If training data is linearly separable, perceptron is guaranteed to find *some* linear separator
- Which of these is **optimal**? How do we define optimal?



Support Vector Machine (SVM)

- SVMs (Vapnik, 1990' s) choose the linear separator with the **largest margin**


Robust to
outliers!



V. Vapnik

- Good according to intuition, theory, practice
- SVM became famous when, using images as input, it gave accuracy comparable to neural-network in a handwriting recognition task

Support vector machines: 3 key ideas



1. Use **optimization** to find solution with largest margin
2. Seek **large margin** separator while allowing for some test errors
3. Use **kernel trick** to make large non-linear feature spaces computationally efficient

Optimization Problem

- Assume for now data is linearly separable
- Goal: Formulate the max margin problem as a tractable optimization problem:
 - Maximize an objective function subject to constraints, giving optimal \mathbf{w} and b .

A Few Words About Linear Programming

- Class of constrained optimization problems
- Examples given on board
- Well studied, can solve problems with millions of variables and constraints
 - Many software packages, textbooks
- Quadratic programs are also tractable by similar techniques.

Going to Show for Linearly Separable Case

Theorem: Let \mathbf{w} , b be an optimal solution to:

Minimize $\|\mathbf{w}\|^2 = w_1^2 + w_2^2 + \dots + w_n^2$

subject to:


$$y^{(i)}(\mathbf{w} \bullet \mathbf{x}^{(i)} + b) \geq 1 \text{ for all } i = 1, 2, \dots, m$$

Then $\mathbf{w} \bullet \mathbf{x} + b = 0$ is a hyperplane that provides the maximum margin.

Maximize Margin Optimization Problem

For data point $\mathbf{x}^{(i)}$
let $\delta^{(i)}$ be the
distance to the
plane. The **margin**
is $\delta := \min \delta^{(i)}$


We want to find
a hyperplane (ie,
 \mathbf{w} and b) that
maximizes δ . To
this end, let's
calculate $\delta^{(i)}$



How can we calculate $\delta^{(i)}$?

Quiz: find $\delta^{(i)}$ in terms of \mathbf{w} , b , and $\mathbf{x}^{(i)}$

- Let \mathbf{z} be point on plane nearest to $\mathbf{x}^{(i)}$
- $\delta^{(i)} = |\mathbf{x}^{(i)} - \mathbf{z}|$
- (Assume first $\mathbf{x}^{(i)}$ and \mathbf{w} on same side of plane.)
- Note \mathbf{w} and $\mathbf{x}^{(i)} - \mathbf{z}$ point in same direction
- $(\mathbf{x}^{(i)} - \mathbf{z}) / \delta^{(i)} = \mathbf{w} / \|\mathbf{w}\|$
- Since \mathbf{z} is on plane:
 $\mathbf{w} \cdot \mathbf{z} + b = 0$



Complete quiz!

Answer:

$$\delta^{(i)} = y^{(i)}(\mathbf{w} \bullet \mathbf{x}^{(i)} + b) / |\mathbf{w}|$$

Thus if:

$$y^{(i)}(\mathbf{w} \bullet \mathbf{x}^{(i)} + b) / |\mathbf{w}| \geq \delta \text{ for all } i$$

then margin will be at least δ .

Margin Optimization Problem

Therefore, to maximize the margin, want to choose \mathbf{w} and b to

maximize δ

subject to:

$$y^{(i)}(\mathbf{w} \bullet \mathbf{x}^{(i)} + b) / |\mathbf{w}| \geq \delta \quad \text{for all } i = 1, \dots, m$$

Problem: Constraints are nonlinear. Difficult to solve.

Convert to Tractable Problem

Claim: If data is linearly separable, there exists \mathbf{w} , b such that:

- for all positive examples ($y^{(i)} = 1$):

$$\mathbf{w} \bullet \mathbf{x}^{(i)} + b \geq 1$$

- for all negative examples ($y^{(i)} = -1$):

$$\mathbf{w} \bullet \mathbf{x}^{(i)} + b \leq -1$$

Prove it now !

Thus, $y^{(i)} (\mathbf{w} \bullet \mathbf{x}^{(i)} + b) \geq 1$ for all $i = 1, 2, \dots, m$

Maximum Margin: Tractable

Theorem: Let \mathbf{w} , b be an optimal solution to:

Minimize $\|\mathbf{w}\|^2 = w_1^2 + w_2^2 + \dots + w_n^2$


subject to:

$$y^{(i)}(\mathbf{w} \bullet \mathbf{x}^{(i)} + b) \geq 1 \text{ for all } i = 1, 2, \dots, m$$

Then $\mathbf{w} \bullet \mathbf{x} + b = 0$ is a hyperplane that provides the maximum margin.

Proof: You will prove this in HW.

(Hard margin) Support Vector Machines



Minimize $\|w\|^2$

subject to:

$$y^{(i)}(w \cdot x^{(i)} + b) \geq 1 \text{ for all } i$$

Example of a **convex optimization** problem

- A quadratic program
- Polynomial-time algorithms to solve!


Non-support Vectors:

- everything else
- moving them will not change w

Support Vectors:

- data points on the margin lines


But what if you have:



Natural Approach

- Allow for some of the data points to be within the margin.
- Tradeoff: As margin gets bigger, more data points inside
- Can penalize for the # of points inside:
maximize: $\text{margin} - C \times (\# \text{ points inside})$
- But difficult to solve. Instead use distance from margin for violating points.

“Soft margin SVM”




Minimize $_{\mathbf{w}, b, \xi} \mathbf{w} \bullet \mathbf{w} + C \sum_i \xi^{(i)}$
subject to
 $(\mathbf{w} \bullet \mathbf{x}^{(i)} + b)y^{(i)} \geq 1 - \xi^{(i)}$ for all i
 $\xi^{(i)} \geq 0$ for all i

Slack penalty $C > 0$:

- Want to find \mathbf{w} , b so that the the margin is large and the # of errors is small.
- Want large margin to prevent overfitting.
- Solve optimization problem for different values of C . Choose the C that gives the smallest validation error.

Can Remove the Constraints



Minimize $_{\mathbf{w}, b, \xi}$ $\mathbf{w} \bullet \mathbf{w} + C \sum_i \xi^{(i)}$
subject to
 $(\mathbf{w} \bullet \mathbf{x}^{(i)} + b)y^{(i)} \geq 1 - \xi^{(i)}$ for all i
 $\xi^{(i)} \geq 0$ for all i

At optimal solution, for each data point:

- If $(\mathbf{w} \bullet \mathbf{x}^{(i)} + b)y^{(i)} \geq 1$, $\xi^{(i)} = 0$
- If $(\mathbf{w} \bullet \mathbf{x}^{(i)} + b)y^{(i)} < 1$, $\xi^{(i)} = 1 - (\mathbf{w} \bullet \mathbf{x}^{(i)} + b)y^{(i)}$ (constraint binding)

So $\xi^{(i)} = \max(0, 1 - (\mathbf{w} \bullet \mathbf{x}^{(i)} + b)y^{(i)})$

Summary: SVM Soft-Margin Optimization Problem

Minimize _{\mathbf{w}, b, ξ} $\mathbf{w} \bullet \mathbf{w} + C \sum_i \xi^{(i)}$

subject to

$(\mathbf{w} \bullet \mathbf{x}^{(i)} + b)y^{(i)} \geq 1 - \xi^{(i)}$ for all i , $\xi^{(i)} \geq 0$ for all i

Or equivalently, the non-linear
unconstrained problem:

Minimize _{\mathbf{w}, b} $\mathbf{w} \bullet \mathbf{w} + C \sum_i \max(0, 1 - y^{(i)} (\mathbf{w} \bullet \mathbf{x}^{(i)} + b))$


regularization

Cross validation

- Divide labeled examples into 10 parts.
 - For each part
 - Use that part for validation, other 9 parts for training (optimization problem). Obtain validation error.
 - Calculate the average validation error over the 10 parts.
- Do for each C .
- Choose the C that has the lowest average validation error.