PSC 202 SYRACUSE UNIVERSITY


# INTRODUCTION TO POLITICAL ANALYSIS

MORE BIVARIATE HYPOTHESIS TESTING, HYPOTHESIS TESTING WITH A SAMPLE

#### MORE ON REGRESSION LINE

- How is linear regression useful?
- Caveats about linear regression


• Linear regression widely used in private sector



- Insurance company has to decide how much to charge you
- How much to charge you depends on how much in damages they expect to have to pay for you
- Guessing won't do
  - If they overestimate how much damage someone will cause, they charge too much (and the person might buy insurance elsewhere)
  - If they underestimate, they charge too little (and lose money)

- They use linear regression
- Have data on how much damage other customers have caused
  - Regression analysis of damages caused (Y), depending on age (X)
  - Based on your age, predict how much damage you will cause
    - Damages = a + b\*age
  - That determines your rate

Linear regression also widely used in public policy research



 Infant mortality rates (Death under 1 year of age per 1,000 live births)

- Some of these rates are appalling
  - Mali: Out of 1,000 babies born alive, 100 die before their first birthday
- If we want to lower infant mortality rates, we need to know what causes them

• Infant mortality rate = 39.9 - 0.0008889\*GDP per capita

- Infant mortality rate = 39.9 0.0008889\*GDP per capita
  - For each dollar that GDPpc is higher, infant mortality expected to decrease by 0.0008889
  - If GDPpc=0, infant mortality is expected to be 39.9


- Infant mortality rate = 39.9 0.0008889\*GDP per capita
  - GDP per capita of the U.S. is \$41,627
    - Expected rate: 39.9 0.0008889\*41,627=2.90

- Infant mortality rate = 39.9 0.0008889\*GDP per capita
  - GDP per capita of the U.S. is \$41,627
    - Expected rate: 39.9 0.0008889\*41,627=2.90
  - GDP per capita of Mexico is \$11,877
    - Expected rate: 39.9 0.0008889\*11,877=29.34


## TODAY

- How is linear regression useful?
- Caveats about linear regression


# **PLOT**



# OH NO...




## LINEARITY




- Linear regression really means linear
- Often, effect of x on y is not linear

# FROM OUR SURVEY




- Score=-6.8 + 0.8 \* Lib/Cons
- Intercept is negative!

## FROM OUR SURVEY




• Always start an analysis by getting to know your data, make plots etc.

## ANOTHER THING




- This line is the line that minimizes squared prediction error
- But: Even this line has a lot of prediction error!

## MORE GENERALLY




- Same regression equation in both situations
  - Y = 2 + 3\*X
- But: X explains Y much better in the first than in the second
- Regression equation does not tell us how much it explains

- Need: measure of how well independent variable explains dependent variable in a linear regression
- Idea: How much of the variation in Y can we predict using X?



- We take mean (25) and compute squared prediction error for each observation
- =Variance of Y (test score): 47.5



- Now: We take regression line and compute squared prediction error for each observation
- ="Residual variance"=29.6


- Squared prediction error without regression line: 47.5
- Squared prediction error with regression line (for hours spent studying): 29.6
- 29.6 is 62.3% of 47.5
  - So we were able to reduce squared prediction error by 100-62.3=37.7%
  - In other words, hours spent studying explains
    37.7% of variance in test scores

#### R-SQUARE


- Measure is called R<sup>2</sup>
- Interpretation: R<sup>2</sup> tells us how much variation of the dependent variable is explained by the independent variable

#### R-SQUARE

- Typically, not expressed as percentage (between 0 and 100), but as fraction (between 0 and 1)
  - 0: The independent variable explains none of the variation in the dependent variable
  - 1: The independent variable explains all of the variation in the dependent variable



- Hours spent studying explains 37.7% of variance in test scores
- So:  $R^2 = 0.377$




- Hours spent studying explains 100% of variance in test scores
- So:  $R^2 = 1$

# BACK TO OUR EXAMPLE

. reg therm_2	libcons_1						
Source	SS	df	MS	Numb	er of obs	=	74
				- F(1,		=	8.06
Model	3834.01698	1	3834.01698	Prob	> F	=	0 0050
Residual	34232.5776	72	475.452467		uared	=	0.1007
				- Adj	R-squared	=	U.U00Z
Total	38066.5946	73	521.4602	Root	MSE	=	21.805
therm_2	Coef.	Std. Err.	t	P> t	[95% (	Conf.	Interval]
libcons_1	347605	.1224088	-2.84	0.006	59162	24	1035876
_cons	63.79618	4.3579	14.64	0.000	55.108	887	72.4835


- DV: Rating of J. Biden (therm\_2)
- IV: Liberal-conservative scale (libcons\_1)

## JOE BIDEN



- Here: R<sup>2</sup>=0.10
- Lib/cons rating only explains 10% of variance in ratings of J.
  Biden
- So 90% remain unexplained...

# DONALD TRUMP




#### DONALD TRUMP

reg therm_1	libcons_1						
Source	SS	df	MS	Numb	per of obs	5 =	51
					49)	=	39.09
Model	16298.2825	1	16298.282	2 <b>5</b> Prob	) > F	=	2.000
Residual	20428.345	49	416.90	<b>5</b> R-sc	quared	=	0.4438
				— Adj	R-squared	= t	0.4324
Total	36726.6275	50	734.53254		MSE	=	20.418
therm_1	Coef.	Std. Err.	t	P> t	[95% (	Conf.	Interval]
libcons_1	.7643888	.1222537	6.25	0.000	.5187	L08	1.010067
_cons	-6.759461	5.118334	-1.32	0.193	-17.045	514	3.526216

- DV: Rating of D. Trump (therm\_1)
- IV: Liberal-conservative scale (libcons\_1)

#### DONALD TRUMP



- Here: R<sup>2</sup>=0.44
- Lib/cons rating explains 44% of variance in ratings of D. Trump
- So only 56% remain unexplained...

#### WHAT WE CAN DO

- We can now...
  - Estimate how much an independent variable X affects a dependent variable Y
  - Tell how much of the variance in Y is explained by X

# BIVARIATE RELATIONSHIPS

# Independent Variable

<u>e</u>		Nominal/Ordinal	Interval		
ependent Variable	Nominal/Ordinal	Cross-Tabulation	Not In This Class		
Depende	Interval	Mean Comparison	Correlation Coefficient, Linear Regression		

#### NOW

- Is the effect of lib/cons on ratings of J. Biden real?
- Or is it only something that we found in our sample, but lib/cons actually has no effect in the population?


#### REMEMBER

POLITICS SEPTEMBER 22, 2021

# Biden's Approval Rating Hits New Low of 43%; Harris' Is 49%


Results for this Gallup poll are based on telephone interviews conducted Sept. 1-17, 2021, with a random sample of 1,005 adults, aged 18 and older, living in all 50 U.S. states and the District of Columbia. For results based on the total sample of national adults, the margin of sampling error is ±4 percentage points at the 95% confidence level.

# BIVARIATE RELATIONSHIP



What explains why some people approve of J.
 Biden while others do not?

# BIVARIATE RELATIONSHIP



- Hypothesis: In a comparison of individuals, women are more likely to approve of J. Biden than men
  - "gender gap"

#### BIVARIATE RELATIONSHIP

# Biden Approval Ratings Diverge by Gender, Education, Race

Job Approval Ratings of President Biden, by Subgi	roup		
	Approve	Disapprove	N
	%	%	
All U.S. adults	56	39	2,937
Gender			
Men	49	45	1,643
Women	62	34	1,294

#### **PROBLEM**

- Is the effect of gender on approval real?
  - Does it exist in the population?
- Or is it only something that we found in our sample, but gender actually has no effect in the population?

#### **PROBLEM**

- We have a random sample
  - Men: 49% approval
  - Women: 62% approval
- Want to know: is mean approval rating of men and women in the population the same or not?

# SOLUTION

- Idea: Use relation between two variables in sample to make inference about relation between two variables in population
  - Of course, means we can make mistakes

# ALTERNATIVE HYPOTHESIS

 There is a relationship between the independent and dependent variable in the population

H<sub>A</sub> or H<sub>1</sub>

## **NULL HYPOTHESIS**

- In the population, there is no relationship between dependent and independent variable
  - If there is a difference in the sample, it is due to random sampling error

H<sub>0</sub>


# IN OUR CASE

- H<sub>0</sub>: In a comparison of individuals, there is no difference between men and women in approval of Biden
- $H_A$ : In a comparison of individuals, women are more likely to approve of Biden than men

## BACK TO MISTAKES

- Idea: Use relation between two variables in sample to make inference about relation between two variables in population
  - Of course, means we can make mistakes

#### **ERRORS**



#### **ERRORS**




# TYPE I ERROR

- We conclude there is a relationship between X and Y when in reality there is not
  - "Type I error"
  - We falsely reject H<sub>0</sub>

# TYPE I ERROR

- We conclude there is a relationship between X and Y when in reality there is not
  - Example: There is no difference between men and women in approval rating in the population, but we conclude there is

#### **ERRORS**




#### TYPE II ERROR

- We conclude there is no relationship between X and Y when in reality there is
  - "Type II error"
  - We falsely do not reject H<sub>0</sub>

## TYPE II ERROR

- We conclude there is no relationship between X and Y when in reality there is
  - Example: There is a difference between men and women in approval rating in the population, but we conclude there is none

#### **ERRORS**



## DECISION

- It's really bad if we conclude there is a relationship when in reality there is not (Type I error)
  - Type II error is also not great, but not as bad
- Thus: We privilege H<sub>0</sub>

#### DECISION

- By default: We start out with assumption that there is no relationship in population (so  $H_0$  is true)
  - No difference between men and women in Biden approval in population

## DECISION

- Ask: Is there enough evidence in the sample to reject H<sub>0</sub>?
  - Is the observed difference between mean and women in sample large enough to reject null hypothesis that no difference between them in population?