

Characteristic	Overall, N = 956 ¹	Severity 1, N = 608 ¹	Severity 2, N = 348 ¹	p-value ²
Treatment according to Guideline 2016				0.002
aim not reached	599 (63%)	404 (66%)	195 (56%)	
aim reached	357 (37%)	204 (34%)	153 (44%)	
Treatment according to Guideline 2019				<0.001
aim not reached	760 (79%)	505 (83%)	255 (73%)	
aim reached	196 (21%)	103 (17%)	93 (27%)	
Treatment according to Guideline 2016 in Patients with low intensity Statin				0.005
aim not reached	360 (66%)	256 (70%)	104 (57%)	
aim reached	189 (34%)	111 (30%)	78 (43%)	

Characteristic	Overall, N = 956 ¹	Severity 1, N = 608 ¹	Severity 2, N = 348 ¹	p-value ²
Treatment according to Guideline 2016 in Patients with high intensity Statin				0.2
aim not reached	239 (59%)	148 (61%)	91 (55%)	
aim reached	168 (41%)	93 (39%)	75 (45%)	
Treatment according to Guideline 2019 in Patients with low intensity Statin				<0.001
aim not reached	457 (83%)	320 (87%)	137 (75%)	
aim reached	92 (17%)	47 (13%)	45 (25%)	
Treatment according to Guideline 2019 in Patients with high intensity Statin				0.2
aim not reached	303 (74%)	185 (77%)	118 (71%)	
aim reached	104 (26%)	56 (23%)	48 (29%)	

Severity 1: PAD grade IIa and IIb

Severity 2: III, IV, and diabetic foot syndrom

Characteristic	Overall, N = 956¹	Severity 1, N = 608¹	Severity 2, N = 348¹	p-value²
¹ Statistics presented: n (%)				
² Statistical tests performed: chi-square test of independence				