

# Algebraic Topology Miscellaneous Notes

Simon Xiang

September 16, 2020

Miscellaneous notes for the Fall 2020 graduate section of Algebraic Topology (Math 380C) at UT Austin, taught by Dr. Allcock. The course was loaded with pictures and fancy diagrams, so I didn't  $\text{\TeX}$  any notes for the lectures themselves. However, I did take some miscellaneous supplementary notes, here they are. Source files: [https://git.simonxiang.xyz/math\\_notes/files.html](https://git.simonxiang.xyz/math_notes/files.html)

## Contents

<b>1</b>	<b>Category Theory</b>	<b>2</b>
1.1	Motivation . . . . .	2
1.2	Categories . . . . .	2
1.3	Functors . . . . .	3
1.4	Natural Transformations . . . . .	5
1.5	Homotopy Categories and Homotopy Equivalence . . . . .	7
<b>2</b>	<b>Free Groups</b>	<b>8</b>
2.1	Words and Reduced Words . . . . .	8
2.2	Free Groups . . . . .	8
2.3	Homomorphisms of Free Groups . . . . .	9
2.4	Free Products of Groups . . . . .	10
<b>3</b>	<b>Van Kampen's Theorem</b>	<b>12</b>
3.1	The van Kampen Theorem . . . . .	12

## §1 Category Theory

Today we talk about abstract nonsense! These notes will follow Evan Chen's Napkin §60 and May's "A Concise Course in Algebraic Topology" §2.

### §1.1 Motivation

Why do we talk about categories? Categories rise from objects (sets, groups, topologies) and maps between them (bijections, isomorphisms, homeomorphisms). Algebraic topology speaks of maps from topologies to groups, which makes maps between categories a suitable tool for us.

**Example 1.1.** Here are some examples of morphisms between objects:

- A bijective homomorphism between two groups  $G$  and  $H$  is an isomorphism. What also works is two group homomorphisms  $\phi: G \rightarrow H$  and  $\psi: H \rightarrow G$  which are mutual inverses, that is  $\phi \circ \psi = \text{id}_H$  and  $\psi \circ \phi = \text{id}_G$ .
- Metric (or topological) spaces  $X$  and  $Y$  are isomorphic if there exists a continuous bijection  $f: X \rightarrow Y$  such that  $f^{-1}$  is also continuous.
- Vector spaces  $V$  and  $W$  are isomorphic if there is a bijection  $T: V \rightarrow W$  that's a linear map (aka,  $T$  and  $T^{-1}$  are linear maps).
- Rings  $R$  and  $S$  are isomorphic if there is a bijective ring homomorphism  $\phi$  (or two mutually inverse ring homomorphisms).

### §1.2 Categories

**Definition 1.1** (Category). A *category*  $\mathcal{A}$  consists of

- A class of *objects*, denoted  $\text{obj}(\mathcal{A})$ .
- For any two objects  $A_1, A_2 \in \text{obj}(\mathcal{A})$ , a class of *arrows* (also called *morphisms* or *maps* between them). Let's denote the set of arrows by  $\text{Hom}_{\mathcal{A}}(A_1, A_2)$ .
- For any  $A_1, A_2, A_3 \in \text{obj}(\mathcal{A})$ , if  $f: A_1 \rightarrow A_2$  is an arrow and  $g: A_2 \rightarrow A_3$  is an arrow, we can compose the two arrows to get  $h = g \circ f: A_1 \rightarrow A_3$  an arrow, represented in the *commutative diagram* below:

$$\begin{array}{ccc} A_1 & \xrightarrow{f} & A_2 \\ & \searrow h & \downarrow g \\ & & A_3 \end{array}$$

The composition operation can be denoted as a function

$$\circ: \text{Hom}_{\mathcal{A}}(A_2, A_3) \times \text{Hom}_{\mathcal{A}}(A_1, A_2) \rightarrow \text{Hom}_{\mathcal{A}}(A_1, A_3)$$

for any three objects  $A_1, A_2, A_3$ . Composition must be associative, that is,  $h \circ (g \circ f) = (h \circ g) \circ f$ . In the diagram above, we say  $h$  *factors* through  $A_2$ .

- Every object  $A \in \text{obj}_{\mathcal{A}}$  has a special *identity arrow*  $\text{id}_A$ . The identity arrow has the expected properties  $\text{id}_A \circ f = f$  and  $f \circ \text{id}_A = f$ .

**Note.** We can't use the word "set" to describe the class of objects because of some weird logic thing (there is no set of all sets). But you can think of a class as a set.

From now on,  $A \in \mathcal{A}$  is the same as  $A \in \text{obj}(\mathcal{A})$ . A category is *small* if it has a set of objects, and *locally small* if  $\text{Hom}_{\mathcal{A}}(A_1, A_2)$  is a set for any  $A_1, A_2 \in \mathcal{A}$ .

**Example 1.2** (Basic Categories). Here are some basic examples of categories:

- We have the category of groups  $\text{Grp}$ .
  - The objects of  $\text{Grp}$  are groups.
  - The arrows of  $\text{Grp}$  are group homomorphisms.
  - The composition of  $\text{Grp}$  is function composition.
- Describe the category  $\text{CRing}$  (of commutative rings) in a similar way.
- Consider the category  $\text{Top}$  of topological spaces, whose arrows are continuous maps between spaces.
- Also consider the category  $\text{Top}_*$  of topological spaces with a distinguished base-point, that is, a pair  $(X, x_0)$ ,  $x_0 \in X$ . Arrows are continuous maps  $f: X \rightarrow Y$  with  $f(x_0) = y_0$ .
- Similarly, the category of (possibly infinite-dimensional) vector spaces over a field  $k$   $\text{Vect}_k$  has linear maps for arrows. There is even a category  $\text{FDVect}_k$  of finite-dimensional vector spaces.
- Finally, we have a category  $\text{Set}$  of sets, arrows denote any map between sets.

### §1.3 Functors

Motivation: maps between categories, objects rising from other objects.

**Example 1.3** (Basic Functors). Here are some basic examples of functors:

- Given an algebraic structure (group, field, vector space) we can take its underlying set  $S$ : this is a functor from  $\text{Grp} \rightarrow \text{Set}$  (or whatever you want to start with).
- If we have a set  $S$ , if we consider the vector space with basis  $S$  we get a functor  $\text{Set} \rightarrow \text{Vect}$ .
- Taking the power set of a set  $S$  gives a functor  $\text{Set} \rightarrow \text{Set}$ .

- Given a locally small category  $\mathcal{A}$ , we can take a pair of objects  $(A_1, A_2)$  and obtain a set  $\text{Hom}_{\mathcal{A}}(A_1, A_2)$ . This turns out to be a functor  $\mathcal{A}^{\text{op}} \times \mathcal{A} \rightarrow \text{Set}$ .

Finally, the most important example (WRT this course):

- In algebraic topology, we build groups like  $H_1(X)$ ,  $\pi_1(X)$  associated to topological spaces. All these group constructions are functors  $\text{Top} \rightarrow \text{Grp}$ .

**Definition 1.2 (Functors).** Let  $\mathcal{A}$  and  $\mathcal{B}$  be categories. A *functor*  $F$  takes every object of  $\mathcal{A}$  to an object of  $\mathcal{B}$ . In addition, it must take every arrow  $A_1 \xrightarrow{f} A_2$  to an arrow  $F(A_1) \xrightarrow{F(f)} F(A_2)$ . Refer to the commutative diagram:

$$\begin{array}{ccccc}
 & A_1 & & B_1 = F(A_1) & \\
 & \downarrow f & \xrightarrow{\quad F \quad} & \downarrow F(f) & \\
 \mathcal{A} \ni & & & & \in \mathcal{B} \\
 & A_2 & & B_2 = F(A_2) & 
 \end{array}$$

Functors also satisfy the following requirements:

- Identity arrows get sent to identity arrows, that is, for each identity arrow  $\text{id}_A$ , we have  $F(\text{id}_A) = \text{id}_{F(A)}$ .
- Functors respect composition: if  $A_1 \xrightarrow{f} A_2 \xrightarrow{g} A_3$  are arrows in  $\mathcal{A}$ , then  $F(g \circ f) = F(g) \circ F(f)$ .

More precisely, these are covariant functors. A contravariant functor  $F$  reverses the direction of arrows, so that  $F$  sends  $f: A_1 \rightarrow A_2$  to  $F(f): F(A_2) \rightarrow F(A_1)$ , and satisfies  $F(g \circ f) = F(f) \circ F(g)$  instead. A category  $\mathcal{A}$  has an opposite category  $\mathcal{A}^{\text{op}}$  with the same objects and with  $\mathcal{A}^{\text{op}}(A_1, A_2) = \mathcal{A}(A_2, A_1)$ . A contravariant functor  $F: \mathcal{A} \rightarrow \mathcal{B}$  is just a covariant functor  $\mathcal{A}^{\text{op}} \rightarrow \mathcal{B}$ .

**Example 1.4.** We have already talked about *free* and *forgetful* functors in Example 1.3: the forgetful functors are functors from spaces to sets (the underlying set of a group) and free functors are from sets to spaces (the basis set forming a vector space).

- Another example of a forgetful functor is a functor  $\text{CRing} \rightarrow \text{Grp}$  by sending a ring  $R$  to its abelian group  $(R, +)$ .
- Another example of a free functor is a functor  $\text{Set} \rightarrow \text{Grp}$  by taking the free group generated by a set  $S$  (who would have known this is free?)

Here is a cool example: functors preserve isomorphism. If two groups are isomorphic, then they must have the same cardinality. In the language of category theory, this can be expressed as such: if  $G \cong H$  in  $\text{Grp}$  and  $U: \text{Grp} \rightarrow \text{Set}$  is the forgetful functor, then  $U(G) \cong U(H)$ . We can generalize this to *any* functor and category!

**Theorem 1.1.** If  $A_1 \cong A_2$  are isomorphic objects in  $\mathcal{A}$  and  $F: \mathcal{A} \rightarrow \mathcal{B}$  is a functor then

$$F(A_1) \cong F(A_2).$$

*Proof.* Let's go diagram chasing!

$$\mathcal{A} \ni \begin{array}{c} A_1 \\ \uparrow f \\ \downarrow g \\ A_2 \end{array} \xrightarrow{\quad F \quad} \begin{array}{c} B_1 = F(A_1) \\ \uparrow F(f) \\ \downarrow F(g) \\ B_2 = F(A_2) \end{array} \in \mathcal{B}$$

The main idea of the proof follows from the fact that functors preserve composition and the identity map.  $\square$

This is very very useful for us (people who are doing algebraic topology) because functors will preserve isomorphism between spaces (we get that homotopic spaces have isomorphic fundamental groups).


**Note.** As a meme (or not really, but it's still funny), we can construct the category  $\text{Cat}$  whose objects are categories and arrows are functors.

## §1.4 Natural Transformations


We talked about maps between objects which led to categories, and then maps between categories which lead to functors. Now let's talk about maps between functors, the natural transformation: this is actually not too strange (recall the homotopy, a "deformation" from a map to another map).

In this case, we also want to pull a map (functor)  $G$  to another map  $G$  by composing a bunch of arrows in the target space  $\mathcal{B}$ .

**Definition 1.3** (Natural Transformations). Let  $F, G: \mathcal{A} \rightarrow \mathcal{B}$  be two functors. A *natural transformation*  $\alpha: F \rightarrow G$  denoted



consists of, for each  $A \in \mathcal{A}$  an arrow  $\alpha_A \in \text{Hom}_{\mathcal{B}}(F(A), G(A))$ , which is called the component of  $\alpha$  at  $A$ . Pictorially, it looks like this:




The  $\alpha_A$  are subject to the “naturality” requirement such that for any  $A_1 \xrightarrow{f} A_2$ , the following diagram commutes:

$$\begin{array}{ccc} F(A_1) & \xrightarrow{F(f)} & F(A_2) \\ \alpha_{A_1} \downarrow & & \downarrow \alpha_{A_2} \\ G(A_1) & \xrightarrow{G(f)} & G(A_2) \end{array}$$

The arrow  $\alpha_A$  represents the path that  $F(A)$  takes to get to  $G(A)$  (like in a homotopy from  $f$  to  $g$  the point  $f(t)$  gets deformed to the point  $g(t)$  continuously). Think of  $f$  representing the homotopy and the basepoints being  $F(A_1), G(A_1)$  to  $F(A_2), G(A_2)$ .

Natural transformations can be composed. Take two natural transformations  $\alpha: F \rightarrow G$  and  $\beta: G \rightarrow H$ . Consider the following commutative diagram:



We can also construct inverses: suppose  $\alpha$  is a natural transformation such that  $\alpha_A$  is an isomorphism for each  $A$ . Then we construct an inverse arrow  $\beta_A$  in the following way:

$$\begin{array}{ccc}
 & & F(A) \in \mathcal{B} \\
 & \nearrow F & \uparrow \alpha_A \\
 \mathcal{A} \ni A & & \\
 & \searrow G & \downarrow \beta_A \\
 & & G(A) \in \mathcal{B}
 \end{array}$$

We say  $\alpha$  is a *natural isomorphism*. Then  $F(A) \cong G(A)$  *naturally* in  $A$  (and  $\beta$  is an isomorphism too!) We write  $F \cong G$  to show that the functors are naturally isomorphic.

**Example 1.5.** If  $F: \text{Set} \rightarrow \text{Grp}$  is the free functor that sends a set to the free group on such set and  $U: \text{Grp} \rightarrow \text{Set}$  is the forgetful functor sending a free group to its generating set, then we have a natural inclusion of  $S \hookrightarrow UF(S)$ . The functors  $F$  and  $U$  are left and right adjoint to each other, in the sense that we have a natural isomorphism

$$\text{Grp}(F(S), A) \cong \text{Set}(S, U(A))$$

for a set  $S$  and an abelian group  $A$ . This expresses the “universal property” of free objects: a map of sets  $S \rightarrow U(A)$  extends uniquely to a homomorphism of groups  $F(S) \rightarrow A$ .

**Definition 1.4.** Two categories  $\mathcal{A}$  and  $\mathcal{B}$  are equivalent if there are functors  $F: \mathcal{A} \rightarrow \mathcal{B}$  and  $G: \mathcal{B} \rightarrow \mathcal{A}$  and natural isomorphisms  $FG \rightarrow \text{Id}$  and  $GF \rightarrow \text{Id}$ , where the  $\text{Id}$  are the respective identity functors.

## §1.5 Homotopy Categories and Homotopy Equivalence

Todo

## §2 Free Groups

Not to be confused with free *abelian* groups. Whether or not we can count is uncertain, but can we even spell? These notes will follow Fraleigh §39 and Hatcher §1.2.

### §2.1 Words and Reduced Words

Let  $A$  be a set of elements (not necessarily finite). We say  $A$  is an *alphabet* and think of the  $a_i \in A$  as *letters*. Symbols of the form  $a_i^n$  are *syllables* and *words* are a finite string of syllables. We denote the *empty word* 1 as the word with no syllables.

**Example 2.1.** Let  $A = \{a_1, a_2, a_3\}$ . Then

$$a_1 a_3^{-4} a_2^2 a_3, a_2^3 a_2^{-1} a_3 a_1, \text{ and } a_3^2$$

are all words (given that  $a_i^1 = a_i$ ).

We can reduce  $a_i^m a_i^n$  to  $a_i^{m+n}$  (*elementary contractions*) or replacing  $a_i^0$  by 1 (dropping something out of the word). Using a finite number of elementary contractions, we get something called a *reduced word*.

**Example 2.2.** The reduced word of  $a_2^3 a_2^{-1} a_3 a_1^2 a_1^{-7}$  is  $a_2^2 a_3 a_1^{-5}$ .

Is it obvious or not that the reduced form of a word is unique? Does it stay the same rel elementary contractions? Apparently you have to be a great mathematician to know.

### §2.2 Free Groups

Denote the set of all reduced words from our alphabet  $A$  as  $F[A]$ . We give  $F[A]$  a group structure in the natural way: for two words  $w_1$  and  $w_2$  in  $F[A]$ , let  $w_1 \cdot w_2$  be the result by string concatenation of  $w_2$  onto  $w_1$ .

**Example 2.3.** If  $w_1 = a_2^3 a_1^{-5} a_3^2$  and  $w_2 = a_3^{-2} a_1^2 a_3 a_2^{-2}$ , then  $w_1 \cdot w_2 = a_2^3 a_1^{-3} a_3 a_2^{-2}$ .

“It would seem obvious” that this indeed forms a group on the alphabet  $A$ . Man, the weather outside today is nice.

**Definition 2.1** (Free Group). The group  $F[A]$  described above is the *free group generated by  $A$* .

Sometimes we have a group  $G$  and a generating set  $A = \{a_i \mid i \in I\}$ , and we want to know whether or not  $G$  is *free* on  $\{a_i\}$ , that is,  $G$  is the free group generated by  $\{a_i\}$ .

**Definition 2.2** (Free Generators). If  $G$  is a group with a set  $A = \{a_i\}$  of generators and is isomorphic to  $F[A]$  under a map  $\phi: G \rightarrow F[A]$  such that  $\phi(a_i) = a_i$ , then  $G$  is *free on  $A$* , and the  $a_i$  are *free generators of  $G$* . A group is *free* if it is free on some nonempty set  $A$ .

Oh you’ll be free... free indeed...



**Example 2.4.**  $\mathbb{Z}$  is the free group on one generator.

I wish we would call it the “free group on  $n$  letters” as opposed to the “free group on  $n$  generators”, which is lame, to be consistent with the whole “mathematicians don’t know how to spell” theme.

**Example 2.5.**  $\mathbb{Z}$  is the free group on one letter.

Much better. Time for theorem spam.

**Theorem 2.1.** *If  $G$  is free on  $A$  and  $B$ , then  $A$  and  $B$  have the same order, that is, any two sets of free generators of a free group have the same cardinality.*

*Proof.* Refer “to the literature”. ⊠

**Definition 2.3 (Rank).** If  $G$  is free on  $A$ , then the number of letters in  $A$  is the *rank* of the free group  $G$ .

**Theorem 2.2.** *Two free groups are isomorphic if and only if they have the same rank.*

*Proof.* Immediate. ⊠

**Theorem 2.3.** *A nontrivial proper subgroup of a free group is free.*

*Proof.* Back “to the literature”. ⊠

**Example 2.6.** Let  $F[\{x, y\}]$  be the free group on  $\{x, y\}$ . Let

$$y_k = x^k y x^{-k}$$

for  $k \geq 0$ . The  $y_k$  for  $k \geq 0$  are free generators for the subgroup of  $F[\{x, y\}]$  that they generate. So the rank of the free subgroup of a free group can be much greater than the rank of the whole group.

## §2.3 Homomorphisms of Free Groups

**Theorem 2.4.** *Let  $G$  be generated by  $A = \{a_i \mid i \in I\}$  and let  $G'$  be any group. If  $a_i'$  for  $i \in I$  are any elements in  $G'$  not necessarily distinct, then there is at most one homomorphism  $\phi: G \rightarrow G'$  such that  $\phi(a_i) = a_i'$ . If  $G$  is free on  $A$ , then there is exactly one such homomorphism.*

*Proof.* Let  $\phi$  be a homomorphism from  $G$  into  $G'$  such that  $\phi(a_i) = a_i'$ . Then any  $x \in G$  can be written as a finite product of the generators  $a_i$ , denoted

$$x = \prod_j a_{i_j}^{n_j},$$

the  $a_i$  not necessarily distinct. Since  $\phi$  is a homomorphism, we have

$$\phi(x) = \prod_j \phi(a_{i_j}^{n_j}) = \prod_j (a'_{i_j})^{n_j},$$

so a homomorphism is completely determined by its values on elements of a generating set. This shows that there is at most one homomorphism such that  $\phi(a_i) = a_i'$ .

Now suppose that  $G$  is free on  $A$ , that is,  $G = F[A]$ . For

$$x = \prod_j a_{i_j} \in G,$$

define  $\psi: G \rightarrow G'$  by

$$\psi(x) = \prod_j (a'_{i_j})^{n_j}.$$

The map is well defined, since  $F[A]$  consists precisely of reduced words. Since the rules for computation involving exponents are formally the same as those involving exponents in  $G$ , it can be seen that  $\psi(xy) = \psi(x)\psi(y)$  for any elements  $x$  and  $y$  in  $G$ , so  $\psi$  is indeed a homomorphism.  $\square$

Note that this theorem states that a group homomorphism is completely determined by its value on each element of a generating set: eg, a homomorphism of a cyclic group is completely determined by its value on any single generator.

**Corollary 2.1.** *Every group  $G'$  is a homomorphic image of a free group  $G$ .*

*Proof.* Let  $G' = \{a'_i \mid i \in I\}$ , and let  $A = \{a_i \mid i \in I\}$  be a set with the same number of elements as  $G'$ . Let  $G = F[A]$ . Then by Theorem 2.4 there exists a homomorphism  $\psi$  mapping  $G$  into  $G'$  such that  $\psi(a_i) = a'_i$ . Clearly the image of  $G$  under  $\psi$  is all of  $G'$ .  $\square$

Only the free group on one letter is abelian.

## §2.4 Free Products of Groups

**Definition 2.4** (Free Products). As a set, the free product  $*_{\alpha} G_{\alpha}$  consists of all words  $g_1 g_2 \cdots g_m$  of arbitrary finite length  $m \geq 0$ , where each letter  $g_i$  belongs to a group  $G_{\alpha_i}$  and is not the identity element of  $G_{\alpha_i}$ , and adjacent letters  $g_i$  and  $g_{i+1}$  belong to different groups  $G_{\alpha}$ , that is,  $\alpha_i \neq \alpha_{i+1}$ .

Basically, reduced words with alternating letters from different groups. The group operation is concatenation: what if the end of  $w_1$  and the beginning of  $w_2$  belong to the same  $G_{\alpha}$ ? Merge them into a syllable: what if they merge into the identity, and so the next two letters are from the same alphabet? Merge again, and repeat forever. Eventually we'll get a reduced word.

How to prove this is associative? Relate it to a subgroup of the symmetric group, it takes care of a lot of work. So we have the free product  $\mathbb{Z} * \mathbb{Z}$ , which is also free. Note that  $\mathbb{Z}_2 * \mathbb{Z}_2$  is *not* a free group: since  $a^2 = e = b^2$ , powers of  $a$  and  $b$  are not needed. So  $\mathbb{Z}_2 * \mathbb{Z}_2$  consists of the alternating words  $a, b, ab, ba, aba, bab, abab, \dots$  together with the empty word.

A basic property of the free product  $*_{\alpha} G_{\alpha}$  is that any collection of homomorphisms  $\varphi_{\alpha}: G_{\alpha} \rightarrow H$  extends uniquely to a homomorphism  $\varphi: *_{\alpha} G_{\alpha} \rightarrow H$ . Namely, the value of  $\varphi$  on a word  $g_1 \cdots g_n$  with  $g_i \in G_{\alpha_i}$  must be  $\varphi_{\alpha_1}(g_1) \cdots \varphi_{\alpha_n}(g_n)$ , and using this formula to define  $\varphi$  gives a well-defined homomorphism since the process of reducing an unreduced product in  $*_{\alpha} G_{\alpha}$  goes not affect its image under  $\varphi$ .

**Example 2.7.** For a free product  $G * H$ , the inclusions  $G \hookrightarrow G \times H$  and  $H \hookrightarrow G \times H$  induce a surjective homomorphism  $G * H \rightarrow G \times H$ .

## §3 Van Kampen's Theorem

OK guys, let's decompose big spaces into smaller ones and compute their fundamental groups. These notes follow Hatcher §1.2.

### §3.1 The van Kampen Theorem

Let's take a space  $X$  and say it's the union of path-connected open subsets  $A_\alpha$ , each of which contains the basepoint  $x_0 \in X$ . Then the homomorphisms  $j_\alpha: \pi_1(A_\alpha) \rightarrow \pi_1(X)$  induced by the inclusions  $A_\alpha \hookrightarrow X$  extend to a homomorphism  $\Phi: \ast_\alpha \pi_1(A_\alpha) \rightarrow \pi_1(X)$ . The van Kampen theorem will say that  $\Phi$  is often onto but in general, we can expect  $\Phi$  to have a nontrivial kernel.

For if  $i_{\alpha\beta}: \pi_1(A_\alpha \cap A_\beta) \rightarrow \pi_1(A_\alpha)$  is the homomorphism induced by the inclusion  $A_\alpha \cap A_\beta \hookrightarrow A_\alpha$  then  $j_\alpha i_{\alpha\beta} = j_\beta i_{\beta\alpha}$ , both of these compositions being induced by the inclusion  $A_\alpha \cap A_\beta \hookrightarrow X$ , so the kernel of  $\Phi$  contains all the elements of the form  $i_{\alpha\beta}(\omega) i_{\beta\alpha}(\omega)^{-1}$  for  $\omega \in \pi_1(A_\alpha \cap A_\beta)$ .

Van Kampen says under fairly broad hypotheses that this determines all of  $\Phi$ .

**Theorem 3.1.** *If  $X$  is the union of path-connected open sets  $A_\alpha$  each containing the basepoint  $x_0 \in X$  and if each intersection  $A_\alpha \cap A_\beta$  is path-connected, then the homomorphism*

$$\Phi: \ast_\alpha \pi_1(A_\alpha) \rightarrow \pi_1(X)$$

*is onto. Furthermore, if each intersection  $A_\alpha \cap A_\beta \cap A_\gamma$  is path-connected, then the kernel of  $\Phi$  is the normal subgroup  $N$  generated by all elements of the form  $i_{\alpha\beta}(\omega) i_{\beta\alpha}(\omega)^{-1}$  for  $\omega \in \pi_1(A_\alpha \cap A_\beta)$ , and hence  $\Phi$  induces an isomorphism*

$$\pi_1(X) = \ast_\alpha \pi_1(A_\alpha) / N.$$

**Example 3.1** (Wedge Sums). I like the visual of the wedge sum but the terminology of the smash product. We define the wedge sum  $\bigvee_\alpha X_\alpha$  with basepoints  $x_\alpha \in X_\alpha$  as the disjoint union  $\coprod_\alpha X_\alpha$  with all the basepoints  $x_\alpha$  identified to a single point.