

# Complex Analysis Lecture Notes

Simon Xiang

These are my lecture notes for the Fall 2020 section of Complex Analysis (Math 361) at UT Austin with Dr. Radin. These were taken live in class, usually only formatting or typo related things were corrected after class. You can view the source code here: [https://git.simonxiang.xyz/math\\_notes/file/freshman\\_year/complex\\_analysis/master\\_notes.tex.html](https://git.simonxiang.xyz/math_notes/file/freshman_year/complex_analysis/master_notes.tex.html). I was also unhappy with the textbook, so some supplementary notes from different texts are found at the bottom of the document.

## Contents

1	August 27, 2020	3
1.1	Basic Properties of Complex Numbers	3
1.2	Real and Imaginary Parts	3
1.3	Complex Numbers in the Plane	3
2	September 1, 2020	4
2.1	Units and Zero Divisors in the Complex Numbers	4
2.2	Polar Coordinate Notation	4
2.3	On the Norm (Modulus) of a Complex Number	4
2.4	Euler's Formula	5
3	September 3, 2020	6
3.1	Fractional Powers	6
3.2	Point Set Topology	6
3.3	Interior, Closure, Boundary	6
3.4	Open and Closed Sets	7
3.5	Path Connectedness	7
4	September 8, 2020	8
4.1	Accumulation Points	8
4.2	Limits	8
4.3	Continuity	8
5	September 10, 2020	10
5.1	More on Continuity	10
5.2	Limits near Infinity	10
5.3	Derivates	11
5.4	Product, Quotient, and Chain Rules	12
6	September 15, 2020	13
6.1	Cauchy-Riemann Equations	13
6.2	Weak Converse of CR Equations	14
6.3	CR Equations in Polar Coordinates	14
7	September 17, 2020	15
7.1	CR Equations (cont)	15
7.2	Analytic Functions	15
7.3	Harmonic Equations	16
8	September 22, 2020	17
8.1	Trig functions	17
8.2	Hyperbolic trig functions	18
9	September 24, 2020	19
9.1	Logarithmic functions	19
10	October 1, 2020	19

10.1	Catching up: logarithms and branches . . . . .	19
10.2	Branches of the logarithm . . . . .	20
10.3	Logarithmic identities . . . . .	20
10.4	Complex exponents . . . . .	21
11	October 6, 2020 . . . . .	22
11.1	Parametrized curves . . . . .	22
12	Holomorphic Functions . . . . .	23
12.1	Continuous functions . . . . .	23
12.2	Holomorphic functions . . . . .	23
12.3	Complex-valued functions as mappings . . . . .	23

## §1 August 27, 2020

### §1.1 Basic Properties of Complex Numbers

We talk about functions  $f: \mathbb{C} \rightarrow \mathbb{C}$  that map variables  $z \mapsto f(z)$ . This course is “not a very hard course” (it’s a fun course!). Holomorphic functions have very nice properties automatically that real valued differentiable functions simply don’t have.

**Definition 1.1** (Complex Addition). We define complex numbers as ordered pairs  $z = (x, y)$  where  $x, y \in \mathbb{R}$ , with the binary operation of complex addition being defined as

$$(x_1, y_1) + (x_2, y_2) = (x_1 + x_2, y_1 + y_2),$$

where  $+$  denotes addition on the reals.

Once we define multiplication and additive/multiplicative inverses, we will have (almost) formed the field  $\mathbb{C}$ .

**Definition 1.2** (Complex Multiplication). For  $x, y \in \mathbb{C}$ , we have

$$(x_1, y_1)(x_2, y_2) = (x_1x_2 - y_1y_2, x_1y_2 + y_1x_2).$$

Note: for  $a \in \mathbb{R}$ , we define

$$a(x, y) = (ax, ay).$$

Recall  $(a, 0)(x, y) = (ax, ay)$ . So one can understand that  $a \in \mathbb{R}$  is simply the real analog of  $(a, 0)$  (or simply,  $\text{Re}(a, 0) = a \in \mathbb{R}$ ).

How do we define multiplication of a complex number by a real number? We can think of the reals acting (in a group sense) on the complex numbers, with the operation being the standard multiplication.

**Example 1.1.** Take  $(1, 0)(x, y) = (x, y)$ . So  $1(x, y) = (x, y)$  (where  $1 \in \mathbb{R}$ ).

**Example 1.2** (Complex Addition is Commutative). We have already defined the sum of two complex numbers  $z_1 + z_2$  as  $z_3 = z_1 + z_2 = (x_1 + x_2, y_1 + y_2)$ . Since addition is commutative on the real numbers, we have

$$z_1 + z_2 = (x_1 + x_2, y_1 + y_2) = (x_2 + x_1, y_2 + y_1) = z_2 + z_1,$$

so complex addition is commutative.

Claim: multiplication of complex numbers is commutative. You can verify this at home.

**Theorem 1.1** (Distributive Law). We have

$$z_1(z_2 + z_3) = z_1z_2 + z_1z_3,$$

for  $z_1, z_2, z_3 \in \mathbb{C}$ .

*Proof.* This follows from the fact that  $\mathbb{C}$  has a ring structure. □

### §1.2 Real and Imaginary Parts

**Definition 1.3.** If  $z = (x, y)$ , then  $x = \text{Re } z$  and  $y = \text{Im } z$ . Furthermore, we can associate a complex number with a point in the plane in many ways:

(insert figure 1 later)

### §1.3 Complex Numbers in the Plane

Point: the plane is just a plane. The plane doesn’t have to have a coordinate system (coordinate axes don’t have to be perpendicular). Any coordinate system is “useful” for adding complex numbers. For example, you can interpret complex addition as simply vector addition in the plane (no need for orthogonal axes!).

**Definition 1.4** (Additive Inverse). We have

$$-(x, y) = (-1)(x, y) = (-x, -y).$$

So  $(x, y) + [-(x, y)] = (0, 0)$ .

Note:  $(x, y)(0, 1) = (-y, x)$ , a *rotation* of  $(x, y)$  by  $90^\circ$ . Another note: We have  $(x, y) \in \mathbb{C} \cong x + iy$  and  $i = (0, 1)$ . So

$$(x, y) \cong x + iy \cong (x, 0) + (0, 1)(y, 0).$$

## §2 September 1, 2020

### §2.1 Units and Zero Divisors in the Complex Numbers

Recall from last time: A complex number can be defined as  $(x, y) = x + iy$ , where  $x, y \in \mathbb{R}$ . Addition is easy:  $(x_1 + iy_1) + (x_2 + iy_2) = (x_1 + y_1) + i(y_1 + y_2)$ . In particular,  $(0, 0) = 0 + i \cdot 0 = 0$ . For multiplication, assume  $i^2 = -1$ . Then

$$\begin{aligned}(x_1 + iy_1)(x_2 + iy_2) &= (x_1x_2 + iy_1x_2 + iy_2x_1 + i^2y_1y_2) \\ &= x_1x_2 - y_1y_2 + i(y_1x_2 + y_2x_1).\end{aligned}$$

On division: what does it mean to divide complex numbers? We say the multiplicative unit of a complex number (wrt the ring  $\mathbb{C}$ ) as the unique  $\frac{1}{z} = z^{-1}$  s.t.  $z \cdot z^{-1} = z^{-1} \cdot z = (1, 0) \in \mathbb{C}$  (the unity of  $\mathbb{C}$ ). Assume  $(x, y)(x, y)^{-1} = (1, 0)$ . Then do  $u$  and  $v$  exist such that the system of equations

$$\begin{cases} xu - yv = 1 \\ xv + yu = 0 \end{cases}$$

holds? Yes, iff the determinant  $\begin{vmatrix} x & -y \\ y & x \end{vmatrix} = x^2 + y^2$  is non zero.

**Definition 2.1** (Complex Conjugate). We have  $(x, -y)$  the complex conjugate of the complex number  $z = (x, y)$ , denoted  $\bar{z}$ .

We show that  $\mathbb{C}$  has no zero divisors and is therefore an integral domain. WLOG, assume there exists  $z_1, z_2$  such that  $z_1 \neq 0, z_1z_2 = 0$ : then we have  $z_1^{-1}$  exists. So  $z_1^{-1}z_1z_2 = 1z_2 = 0$ , therefore  $z_2 = 0$ . For example: the group  $GL_n(\mathbb{R})$  is not an integral domain, since we have zero divisors (two matrices that when multiplied equal zero).

### §2.2 Polar Coordinate Notation

**Definition 2.2** (Polar Coordinates). Think of  $(x, y)$  as rectangular coordinates in the  $xy$ -plane, and consider the *polar coordinate* notation  $z = [r, \theta]$ , where  $r = \sqrt{x^2 + y^2} = |z|$  (modulus of  $z$ ), and  $\theta = \arctan(\frac{y}{x})$ . So  $[r, \theta] = (r \cos \theta, r \sin \theta)$ .

**Example 2.1** (Multiplication with Polar Coordinates). We have

$$[r_1, \theta_1][r_2, \theta_2] = (r_1 \cos \theta_1, r_1 \sin \theta_1)(r_2 \cos \theta_2, r_2 \sin \theta_2).$$

Then

$$\begin{aligned}(r_1 \cos \theta_1 + ir_1 \sin \theta_1)(r_2 \cos \theta_2 + ir_2 \sin \theta_2) &= \\ r_1r_2 [\cos \theta_1 \cos \theta_2 - \sin \theta_1 \sin \theta_2] + ir_1r_2 [\sin \theta_1 \cos \theta_2 + \sin \theta_2 \cos \theta_1] &= \\ r_1r_2 \cos(\theta_1 + \theta_2) + r_1r_2i \sin(\theta_1 + \theta_2) &= \\ [r_1r_2, \theta_1 + \theta_2].\end{aligned}$$

**Example 2.2.** Assume that a complex number  $z = (x, y)$  is nonzero. Then

$$\frac{1}{(x, y)} = \frac{1(x, -y)}{(x, y)(x, -y)} = \frac{(x, -y)}{x^2 + y^2}.$$

### §2.3 On the Norm (Modulus) of a Complex Number

**Example 2.3.** Some properties of the modulus (norm)  $|z|$ :

1.  $|z_1z_2| = |z_1||z_2|$ ,
2.  $\left|\frac{z_1}{z_2}\right| = \left|z_1 \cdot \frac{1}{z_2}\right| = \left|z_1 \cdot \frac{\bar{z}_2}{|z_2|^2}\right| = |z_1| \frac{|z_2|}{|z_2|^2} = \frac{|z_1|}{|z_2|}$  (clearly  $|\bar{z}_2| = |z_2|$ ),
3.  $|z_1 + z_2| \leq |z_1| + |z_2|$  ( $\mathbb{C}$  is a metric space, so the triangle inequality holds),
4.  $|z_1 + z_2| \geq ||z_1| - |z_2||$  (reverse triangle inequality).

We prove the Reverse Triangle Inequality.

*Proof.* We have  $|z_1| = |z_1 + z_2 - z_2| \leq |z_1 + z_2| + |z_2|$ , so  $|z_1 + z_2| \geq |z_1| - |z_2|$ . A similar argument holds for  $z_2$ .  $\square$

Think of the polar angle as only well defined for multiples of  $2\pi$ . Define the argument (angle) as  $\text{Arg} = -\pi < \theta \leq \pi$  (what??). So  $\text{Arg}(1, 1) = \frac{\pi}{4}$ ,  $\text{Arg}(-1, 0) = \pi$ . OTOH, we would have  $\arg(1, 1) = \frac{\pi}{4} + 2\pi n$ .

## §2.4 Euler's Formula

**Theorem 2.1** (Euler's Formula). *We claim*

$$e^{i\theta} = \cos \theta + i \sin \theta.$$

*Proof.* Try using Maclaurin series. □

This suggests  $e^{i\theta_1} e^{i\theta_2} = e^{i(\theta_1 + \theta_2)}$ . We proved this when we showed  $[r_1, \theta_1][r_2, \theta_2] = [r_1 r_2, \theta_1 + \theta_2]$ .

The reason why Dr. Radin says to "forget about Euler" is because he's trying to make a semi-rigorous (or self-contained) construction of the complex numbers. I think it's fine to rely on intuition from other courses, this isn't Real Analysis (nowhere near as rigorous). If we truly were to construct the field  $\mathbb{C}$ , we would have to cover polynomial rings and the fields generated by PID's quotient irreducible polynomials, then show that  $\mathbb{C} \simeq \mathbb{R}[x]/\langle x^2 + 1 \rangle$  (and show that this new field is algebraically closed too!). Of course this isn't feasible. So let's just think of this as Euler's Formula, and not some weird definition!

Back to math: using our newfound formula, we can simply say  $\arg z = \theta$  such that  $z = re^{i\theta}$  for any  $z \in \mathbb{C}$ . Similarly,  $\text{Arg } z$  is just  $\theta$  restricted to the interval  $(-\pi, \pi]$ .

**Example 2.4.** If  $z = re^{i\theta}$  nonzero, then what is the polar form of  $\frac{1}{z}$ ? It must be

$$\frac{1}{r} e^{-i\theta}.$$

**Example 2.5.** We've seen that  $e^{i\theta_1} e^{i\theta_2} = e^{i(\theta_1 + \theta_2)}$ . Then

$$e^{i\theta_1} (e^{i\theta_2} e^{i\theta_3}) = e^{i\theta_1} e^{i(\theta_2 + \theta_3)} = e^{i(\theta_1 + \theta_2 + \theta_3)}.$$

So  $(\cos \theta + i \sin \theta)^m = \cos(m\theta) + i \sin(m\theta)$ . This is known as *de Moivre's formula*.

## §3 September 3, 2020

### §3.1 Fractional Powers

Let  $z_0 \in \mathbb{C}$ , and define the fractional power  $(z_0)^{\frac{1}{m}}$  for  $m \geq 2$ . This is a complex number such that

$$\left[(z_0)^{\frac{1}{m}}\right]^m = z_0.$$

This may not be unique. To determine the value of the fractional power  $(z_0)^{\frac{1}{m}}$ , let  $z_0 = r_0 e^{i\theta_0}$ ,  $r_0 = |z_0|$ ,  $\theta_0 \in \text{Arg } z_0$ . Then

$$(z_0)^{\frac{1}{m}} = (r_0)^{\frac{1}{m}} e^{i\frac{\theta_0}{m}}.$$

**Example 3.1.** In polar form,  $z_0 = i = e^{i\frac{\pi}{2}}$ . We want  $i^{\frac{1}{6}}$ , one value is  $e^{i\frac{\pi}{12}}$ . Also,

$$e^{i\frac{[\frac{\pi}{2}+2\pi]}{6}} = e^{i[\frac{\pi}{12}+\frac{\pi}{3}]} = e^{i\frac{5\pi}{12}}.$$

In general,  $i = e^{i[\frac{\pi}{2}+2\pi m]}$ , so  $e^{i[\frac{\pi}{12}+\frac{m\pi}{3}]}$  is a value of  $i^{\frac{1}{6}}$  for any  $m$ . In particular, consider the choices  $m = 0, 1, \dots, 5$ . Then

(insert figure later- it has to do with roots of unity on the circle group tho)

This method gives all possible  $n$ -th roots. In particular, in the circle group  $U_1$ , each “walk” is equal to a multiplication of  $\zeta$ .

We will eventually generalize the fractional power  $z_0^{p/q}$  to  $z_0^w$ . Yada yada no exponentials allowed reeee. If you’re going to formalize do it right or don’t do it at all. Half baked rigor is about as useful as a potato (at least a potato can feed your family).

### §3.2 Point Set Topology

Why are we studying abstract nonsense? We need topology to define limits of complex numbers. We will eventually define a derivative as a quotient of deltas, eg

$$\frac{\Delta f}{\Delta z} \rightarrow \frac{df}{dz} \quad \text{as } \Delta z \rightarrow 0.$$

We’ll talk about open and closed sets and accumulation points and such (basic things needed for limits). Consider

$$\tilde{S} = \{z \mid |z| \leq 1 \text{ and } |z| \neq 1 \text{ if } \text{Re } z < 0\}.$$

**Definition 3.1** (Open Ball). We define an open ball

$$B(z_0, \epsilon) = \{z \mid |z - z_0| < \epsilon\}.$$

### §3.3 Interior, Closure, Boundary

**Definition 3.2** (Interior Point). We have an *interior point* a point in a set such that there exists an open ball centered at the point entirely contained in the set. We define the set of all interior points of a set  $X$  as  $\text{Int}(X)$ .

Note that  $\text{Int}(\tilde{S}) = \{z \mid |z| < 1\}$ .

**Definition 3.3** (Exterior Point). A point  $z_0$  is an exterior point of  $S$  if there exists a ball

$$B(z_0, \epsilon) \subseteq S^c,$$

ie,  $z_0 \in \text{Int}(S^c)$ .

**Definition 3.4** (Boundary Point). A point  $z_0$  is a boundary point of  $S$  if for ball  $B(z_0, \epsilon)$  centered at  $z_0$ ,  $B(z_0, \epsilon) \cap S \neq \emptyset$  and  $B(z_0, \epsilon) \cap S^c \neq \emptyset$ . We define the *boundary* of a set  $S$  as the set of all boundary points, denoted  $\partial S$ .

Basic things: points can’t be both in the interior and exterior, boundary and interior, etc etc.

**Theorem 3.1.** For any set  $S$ ,  $\text{Int}(S)$ ,  $\text{Ext}(S)$ , and  $\partial S$  form a partition of  $S$ .

We will use  $S^\circ$  to denote the interior and  $(S^c)^\circ$  to denote the exterior of a set from now on.

**Example 3.2.**  $\partial \tilde{S} = \{z \mid |z| = 1\}$ .

**Example 3.3.** We have the unit circle  $S = \{z \mid |z| = 1\} \cup zi$  (where  $zi$  is a point).  $S^\circ = \emptyset$ ,  $zi \in \partial S$ , any point on the rim  $\in \partial S$ , so  $\partial S = S$ . By our previous theorem,  $(S^c)^\circ = S^c$ . (Who even studies the exterior of a set??)

### §3.4 Open and Closed Sets

From now on a set refers to a subset of  $\mathbb{C}$ .

**Definition 3.5** (Open Sets). A set is open if it contains none of its boundary. Alternatively, a set is open iff  $S = S^\circ$ .

**Example 3.4.**  $\mathbb{C}$  is open (and closed)! Furthermore,  $\partial\mathbb{C} = \emptyset$  (which is an alternate condition for a set to be clopen). Note that  $\emptyset$  is also both open and closed, since  $\partial\emptyset = \emptyset$ . This also makes sense if we look at it from the interior perspective (no interior points in  $\emptyset$ , every point has an open ball in  $\mathbb{C}$ ).

**Definition 3.6** (Closed Sets). A set is closed if it contains all of its boundary. (What do you mean not the complement of open???)

**Theorem 3.2.**  $S$  is closed  $\iff S^c$  is open.

*Proof.* Immediate. In general topology, we define open sets this way. □

**Example 3.5.** Like I said earlier, both  $\mathbb{C}$  and  $\emptyset$  are closed. In general topology, we define both  $S, \emptyset \in \tau$ , since they're complements of course they're both open and closed. Exercise: prove that no other sets are both open and closed.

**Definition 3.7** (Closure). The closure  $\bar{S}$  of  $S$  is the union

$$S \cup \partial S.$$

Clearly  $\bar{S}$  is always closed (by our definition).

**Theorem 3.3.**  $S^\circ$  is open for any  $S$ .

Doesn't this follow from the definition too??

### §3.5 Jank Connectedness

**Definition 3.8** (Path-connectedness). A set  $S$  is path-connected if every pair of points  $z_1, z_2 \in S$  is connected by a continuous path in  $S$ .

Every path-connected set is connected (can be written as the union of two disjoint sets). Something about polygonal paths?? Dr. Radin is right, this is most definitely not standard. Is this what physicists do to topology?

Now he's talking about the Topologist's sine curve (the classic counterexample). This is a counterexample to the (false) idea that connected implies path-connected by exhibiting a set that is connected but not path-connected (but we haven't even talked about the standard definition of connectedness yet!).

## §4 September 8, 2020

### §4.1 Accumulation Points

**Definition 4.1.** A connected open set is a *domain*.

**Definition 4.2.** A *region* is a domain that contains none, some, or all of its boundary.

**Definition 4.3** (Bounded Set). A set  $S$  is bounded if

$$S \subseteq B(x_0, \epsilon).$$

for some  $x_0 \in \mathbb{C}$ ,  $\epsilon > 0$ .

**Definition 4.4** (Accumulation Points).  $z_0$  is an accumulation point of  $S$  if for all balls  $B(z_0, \frac{1}{m})$  centered at  $z_0$ , we have

$$B(z_0, \frac{1}{m}) \setminus \{z_0\} \cap S \neq \emptyset.$$

**Example 4.1.** Let  $S = \mathbb{Q}$ . Then  $\frac{1}{2}, \sqrt{2}$  etc are accumulation points of  $S$  (this relies on the fact that  $\mathbb{Q}$  is *dense* in  $\mathbb{R}$ ). This example shows that accumulation points don't have to be in the set themselves.

**Theorem 4.1.** We have  $S$  is closed if and only if  $S$  contains all of its accumulation points, the set of which is denoted  $S'$ . Furthermore, the closure of  $S$  denoted  $\bar{S}$  is equal to  $S \cup S'$ .

*Proof.*  $\implies$  Accumulation points are either in the boundary of  $S$  or in  $S$  itself. Since  $S$  is closed, we have  $S' \subseteq S$ .  
 $\Leftarrow$  If  $z_0 \in \partial S \cap S^c$  it would be an accumulation point of  $S$ , a contradiction. So  $\partial S \subseteq S \implies S$  is closed. (I'll try to write a better proof later).  $\square$

A quick summary of basic p-set topology:

1.  $S$  is open  $\iff S = S^\circ$ ,
2.  $S$  is closed  $\iff S^c$  is open,
3.  $S$  is open  $\iff S$  contains none of  $\partial S$ ,
4.  $S$  is closed  $\iff S$  contains all of  $\partial S$ ,
5.  $S$  is closed  $\iff S$  contains all of  $S'$ .

### §4.2 Limits

Consider a map  $f: \text{Dom}(f) \rightarrow \mathbb{C}$ ,  $\text{Ran}(f) \subseteq \mathbb{C}$  (I prefer the notation  $f: X \rightarrow \mathbb{C}$  where  $X \subseteq \mathbb{C}$ , and  $\text{Ran}(f) = f[X]$ ). The fact that  $f$  is well defined on  $X$  holds because define  $X$  to be a set on which  $f$  is well defined, duh).

We want to talk about whether a function is continuous or not. Intuitively, a function is continuous if points in the image being "close" together imply that points in the preimage are also "close" together (the preimage of an open set is open).

**Definition 4.5** (Epsilon Delta Limits). For  $z_0$  an accumulation point of some subset  $X$  of  $\mathbb{C}$  (a region),  $\lim_{z \rightarrow z_0} f(z)$  exists and has a value of  $L \iff$  for all  $\epsilon > 0$ , there exists a  $\delta > 0$  such that

$$0 < |z - z_0| < \delta \implies |f(z) - L| < \epsilon,$$

where  $z \in X$ . The modulus is just a distance metric: so the epsilon delta definition is the same as what I said earlier, if points are close to each other in the codomain ( $|f(z) - L| < \epsilon$ ), then such points are close to each other in the domain ( $0 < |z - z_0| < \delta$ ).

Some notes: the limit is only defined when  $z_0$  is an accumulation point. This why accumulation points are also sometimes referred to as *limit points*.

### §4.3 Continuity

**Definition 4.6** (Continuity).  $f$  is continuous at  $z_0$  if  $\lim_{z \rightarrow z_0} f(z) = f(z_0)$ .  $f$  is said to be continuous on a set  $X$  if for all  $x \in X$ ,  $f$  is continuous at  $x$ .

We want to *analyze* a function  $f(z)$ , let  $z = (x, y)$  and  $f(z) = f(x, y) = u(x, y) + iv(x, y)$ ,  $u(x, y) = \text{Re } f$  and  $v(x, y) = \text{Im } f$ .

**Theorem 4.2.** We have

$$\lim_{z \rightarrow z_0} f(z) = L \iff \begin{cases} \lim_{z \rightarrow z_0} \text{Re } f(z) \rightarrow \text{Re } L \\ \lim_{z \rightarrow z_0} \text{Im } f(z) \rightarrow \text{Im } L. \end{cases}$$



*Proof.* Homework. ⊠

**Theorem 4.3.** Let  $f: X \rightarrow \mathbb{C}, g: Y \rightarrow \mathbb{C}$ . For an accumulation point  $z_0$  of  $X \cap Y$ , if  $\lim_{z \rightarrow z_0} f(z) = L$  and  $\lim_{z \rightarrow z_0} g(z) = M$ , then (excuse the abuse of notation)

1.  $\lim(f + g) = L + M$ ,
2.  $\lim fg = LM$ ,
3.  $\lim \frac{f}{g} = \frac{L}{M}$  if  $M \neq 0$ .

*Proof.* Same as the ones you'd find in any analysis course. ⊠

Continuity of sums, products, and quotients of functions follow from the above theorem. Now we turn our attention to the composition of functions.

**Theorem 4.4.** Suppose  $f: \mathbb{C} \rightarrow \mathbb{C}$  and  $g: X \rightarrow \mathbb{C}$ . Let  $z_0$  be an accumulation point of  $X$ . Then if  $f$  is continuous at  $z_0$  and  $g$  is continuous at  $f(z_0)$ , we have  $f \circ g$  continuous at  $z_0$ .

**Example 4.2.**  $f(z) = |z^m|$  for a fixed  $m$  is equal to  $(g \circ h)(z)$  where  $h(z) = z^m$  and  $g(w) = |w|$ . Both  $h$  and  $g$  are continuous on  $\mathbb{C}$ , so  $|z^m|$  is also continuous everywhere.

**Example 4.3.** The identity map is continuous. This is trivial (let  $\delta = \epsilon$ ). It follows that maps of the form  $z^n$  is continuous for some positive integer  $n$ .

**Corollary 4.1.** Functions of the form

$$f(z) = \frac{p(z)}{q(z)}$$

where  $p(z)$  and  $q(z)$  are polynomials are continuous given  $q(z) \neq 0$ .

**Example 4.4.** Let  $f(z) = \frac{z}{\bar{z}}, z \neq 0$ . Consider  $z = x + iy$  near 0 with  $x \neq 0, y = 0$ , then  $f(z) = 1$ . If  $x = 0, y \neq 0$  then  $f(z) = -1$ . Therefore  $\lim_{z \rightarrow z_0} \frac{z}{\bar{z}}$  does not exist (standard technique for proving multivariate limits don't exist).

## §5 September 10, 2020

### §5.1 More on Continuity

Last time we talked about the function  $\frac{z}{\bar{z}}$ . What if we define the domain as  $\mathbb{C} \setminus \{0\}$ ? Does  $\lim_{z \rightarrow z_0} \frac{z}{\bar{z}}$  exist? (AKA: is  $\frac{z}{\bar{z}}$  continuous on its domain?)

**Theorem 5.1.** Let  $f: \mathbb{C} \rightarrow \mathbb{C}$  be defined as  $f = u + iv$ . If  $f$  is continuous at  $z_0$ , then

1.  $\bar{f} = u - iv$  is continuous at  $z_0$ . We can also write  $\bar{f}$  as  $g \circ f$  where  $g(w) = \bar{w}$ .
2.  $\frac{f+\bar{f}}{2} = \operatorname{Re}(f)$  is continuous at  $z_0$ .
3.  $\frac{f-\bar{f}}{2i} = \operatorname{Im}(f)$  is continuous at  $z_0$ .

*Proof.* We prove that  $f(z) = \bar{z}$  is continuous at any  $z_0$ . Given  $\varepsilon > 0$ , consider

$$|f(z) - f(z_0)| = |\bar{z} - \bar{z}_0|.$$

We need a  $\delta > 0$  such that

$$0 < |z - z_0| < \delta \implies |\bar{z} - \bar{z}_0| < \varepsilon.$$

Claim: If  $\delta = \varepsilon$ ,  $|\bar{z} - \bar{z}_0| = |\overline{(z - z_0)}| = |z - z_0| = \delta = \varepsilon$ . This is easy to see, so we are done.  $\square$

**Note.** To show that

$$\lim_{z \rightarrow z_0} f(z) = L,$$

we consider neighborhoods (open sets around  $L$ ), or the set of  $z$  such that  $|f(z) - L| < \varepsilon$  (equivalently, the  $z$  such that  $f(z) \in B(L, \varepsilon)$ ). Also,  $\lim_{z \rightarrow z_0} f(z) - L \iff \lim_{z \rightarrow z_0} (f(z) - L) = 0 \iff \lim_{z \rightarrow z_0} (f(z) - L) = 0$ .

### §5.2 Limits near Infinity

Infinity is not a complex number!! Consider the limits

$$\lim_{z \rightarrow \infty} f(z)$$

and

$$\lim_{z \rightarrow z_0} f(z) = \infty.$$

To define these, we use neighborhoods of “ $\infty$ ”. There is no notion of “ $\pm\infty$ ” in the complex numbers. The definition is similar to the one you encountered in Real Analysis:  $z$  is “large” if  $|z| > R$  for all  $R \in \mathbb{R}$ .

**Definition 5.1** (Limits at Infinity). For  $z_0 \in \mathbb{C}$  we say

$$\lim_{z \rightarrow z_0} f(z) = \infty$$

if given some  $R > 0$ ,  $R \in \mathbb{R}$ , there exists some  $\delta > 0$  such that

$$0 < |z - z_0| < \delta \implies |f(z)| > R.$$

**Example 5.1.** We have  $\lim_{z \rightarrow 0} (\frac{1}{z}) = \infty$  since given  $R > 0$ , there exists a  $\delta > 0$  such that  $0 < |z - 0| < \delta$  implies  $|\frac{1}{z}| > R$ , namely,  $\delta = \frac{1}{R}$ , because

$$|z| < \frac{1}{R} \implies \frac{1}{|z|} > R \iff \left| \frac{1}{z} \right| > R.$$

**Definition 5.2** (Limits to Infinity). We say  $\lim_{z \rightarrow \infty} f(z) = L$ ,  $L \in \mathbb{C}$  if and only if for all  $\varepsilon > 0$ , there exists some  $R > 0$  such that

$$|z| > R \implies |f(z) - L| < \varepsilon.$$

**Example 5.2.** We have  $\lim_{z \rightarrow \infty} \frac{1}{z} = 0$ , let  $\varepsilon > 0$ ,  $R = \frac{1}{\varepsilon}$ . Then  $|f(z) - L| = \left| \frac{1}{z} \right|$ , so

$$|z| > R \implies |z| > \frac{1}{\varepsilon} \implies \varepsilon > \frac{1}{|z|} = \left| \frac{1}{z} \right|,$$

and we are done.

**Definition 5.3.** Finally, we say

$$\lim_{z \rightarrow \infty} f(z) = \infty$$

if (for  $R_1, R_2 \in \mathbb{C}$ ) given some  $R_1 > 0$ , there exists an  $R_2 > 0$  such that

$$|z| > R_2 \implies |f(z)| > R_1.$$

**Example 5.3.** We have  $\lim_{z \rightarrow \infty} z^2 = \infty$  since  $|z^2| > R$  whenever  $|z| > \sqrt{R}$ .

### §5.3 Derivates

We are finally ready to define the derivative of a function (the good stuff). Given a function  $f: X \rightarrow \mathbb{C}$ , we will only define the derivative of  $f$  at a point  $z \in X^\circ$ . Recall that  $X^\circ = \{z \in X \mid B(z, \gamma) \subseteq X\}$  for some  $\gamma > 0$ .

**Definition 5.4** (Complex Derivative). A function  $f: X \rightarrow \mathbb{C}$  is said to be *differentiable* at  $z_0 \in X^\circ$  if

$$\lim_{z \rightarrow z_0} \frac{f(z) - f(z_0)}{z - z_0}$$

exists in  $\mathbb{C}$  (so limits to infinity are not allowed. We will examine these “poles” later in the course). If the limit exists, we denote this limit as  $f'(z_0)$ .

**Example 5.4.** Let  $f: \mathbb{C} \rightarrow \mathbb{C}, z \mapsto 7$ . We claim that  $f'(z) = 0$  for all  $z$ , since

$$\frac{f(z) - f(z_0)}{z - z_0} = \frac{7 - 7}{z - z_0} = 0.$$

We only look at the points  $z$  “near” (accumulation points)  $z_0$ , so we don’t have to worry about the case where  $z = z_0$ . So given  $\varepsilon > 0$ ,

$$|z - z_0| < \delta \implies \left| \frac{f(z) - f(z_0)}{z - z_0} \right| < \varepsilon$$

for any  $\delta > 0$ .

**Example 5.5.** Let  $f: \mathbb{C} \rightarrow \mathbb{C}, z \mapsto z$ . We claim  $f'(z) = 1$  since

$$\frac{f(z) - f(z_0)}{z - z_0} = \frac{z - z_0}{z - z_0} = 1$$

for any  $z \neq 0$ . This limit is one since

$$\left| \frac{f(z) - f(z_0)}{z - z_0} - 1 \right| = \left| \frac{z - z_0}{z - z_0} - 1 \right| = 0.$$

**Example 5.6.** Let  $f: \mathbb{C} \rightarrow \mathbb{C}, f(z) = z^2$ . We will show  $f'(z_0) = 2z_0$ . We want to find a  $\delta > 0$  such that

$$0 < |z - z_0| < \delta \implies \left| \frac{f(z) - f(z_0)}{z - z_0} - 2z_0 \right| < \varepsilon.$$

So

$$\left| \frac{z^2 - z_0^2}{z - z_0} - 2z_0 \right| = |(z + z_0) - 2z_0| = |z - z_0| < \varepsilon$$

if  $|z - z_0| < \delta$  with  $\delta = \varepsilon$ . There aren’t any limit signs because we directly invoked the epsilon-delta definition.

**Example 5.7.** Consider  $f(z) = |z|$  (maps will map  $\mathbb{C} \rightarrow \mathbb{C}$  unless otherwise stated from now on). We have showed  $f$  is continuous for all  $z$ , but  $f$  isn’t differentiable at 0. Use the technique at the end of the last example (write out the piecewise definition of the absolute value and show that the limits don’t agree).

What about  $z_0 \neq 0$ ? Is  $f: \mathbb{C} \setminus \{0\} \rightarrow \mathbb{C}$  differentiable? Let  $z_0 \in \mathbb{C} \setminus \{0\}$ , then

$$\frac{f(z) - f(z_0)}{z - z_0} = \frac{|z| - |z_0|}{z - z_0} = \frac{r - r_0}{re^{i\theta} - r_0e^{i\theta_0}}.$$

We let  $z$  get close to  $z_0$  in two different ways. First, assume  $r = r_0$  but  $\theta \neq \theta_0$  (vary the angle, but all having length  $r$ ). Then

$$\frac{r - r_0}{re^{i\theta} - r_0e^{i\theta_0}} = \frac{0}{r(e^{i\theta} - e^{i\theta_0})} = 0.$$

Next, assume  $r \neq r_0$  but  $\theta = \theta_0$  (points on a line with angle  $\theta$ , vary the length). Then

$$\frac{r - r_0}{re^{i\theta} - r_0e^{i\theta_0}} = \frac{r - r_0}{e^{i\theta}(r - r_0)} = e^{-i\theta} \neq 0.$$

So  $f$  is nowhere differentiable.

## §5.4 Product, Quotient, and Chain Rules

To get  $f'(z)$  for  $f(z) = z^m$ , we want a formula. Time for induction!

**Theorem 5.2.** *If  $f'(z_0)$  and  $g'(z_0)$  exist for two functions  $f$  and  $g$ , then so do the derivatives*

1.  $(f + g)'(z_0) = f'(z_0) + g'(z_0),$
2.  $(fg)'(z_0) = f'(z_0)g(z_0) + f(z_0)g'(z_0),$
3.  $\left(\frac{f}{g}\right)'(z_0) = \frac{f'(z_0)g(z_0) - f(z_0)g'(z_0)}{[g(z_0)]^2}$  provided  $g(z_0) \neq 0$ .

**Theorem 5.3.** *If  $g$  is differentiable at  $z_0$  and  $f$  is differentiable at  $g(z_0)$  then  $f \circ g$  is differentiable at  $z_0$  and*

$$(f \circ g)'(z_0) = f'[g(z_0)]g'(z_0).$$

**Note (Leibniz Rule).** Suppose we have  $f_1, f_2, \dots, f_n$  functions all differentiable at  $z_0$ . Then

$$(f_1 f_2 f_3 \cdots f_n)'(z_0) = f_1' f_2 f_3 \cdots f_n + f_1 f_2' f_3 \cdots f_n + f_1 f_2 f_3' \cdots f_n + \cdots.$$

In particular,  $(z^n)' = n(z'z^{n-1}) = nz^{n-1}$  (just take  $f_i = f$  and it becomes clear that this is true).

## §6 September 15, 2020

I could be studying fundamental groups right now, but instead I'm sitting here verifying limits and derivatives by hand. Why?? OK so Gradescope is a meme. Anything new?

Everything so far has been awfully boring. But now it gets interesting. Finally, I've been waiting for this.

### §6.1 Cauchy-Riemann Equations

Suppose we have a function  $f: \mathbb{C} \rightarrow \mathbb{C}$ , write it as  $f(z) = u(x, y) + iv(x, y)$ . Assume  $f'(z_0)$  exists, and is equal to

$$\lim_{(x,y) \rightarrow (x_0,y_0)} \frac{(u(x, y) + iv(x, y)) - (u(x_0, y_0) + iv(x_0, y_0))}{(x + iy) - (x_0 + iy_0)}.$$

Then we rewrite this to get

$$f'(z_0) = \lim_{(x,y) \rightarrow (x_0,y_0)} \frac{u(x, y) - u(x_0, y_0) + i[v(x, y) - v(x_0, y_0)]}{(x - x_0) + i(y - y_0)}.$$

Consider two special ways  $(x, y)$  can be "near"  $(x_0, y_0)$ . First, let  $x = x_0$  but  $y \neq y_0$ . Then the quotient becomes

$$\begin{aligned} \frac{u(x_0, y) - u(x_0, y_0) + i[v(x_0, y) - v(x_0, y_0)]}{i(y - y_0)} &= \\ \frac{u(x_0, y) - u(x_0, y_0)}{i(y - y_0)} + \frac{v(x_0, y) - v(x_0, y_0)}{y - y_0}. \end{aligned}$$

Then the limit is equal to

$$\frac{1}{i} \frac{\partial u}{\partial y}(x_0, y_0) + \frac{\partial v}{\partial y}(x_0, y_0).$$

Now let  $y = y_0$  but  $x \neq x_0$ . Then the quotient becomes

$$\frac{u(x, y_0) - u(x_0, y_0) + i[v(x, y_0) - v(x_0, y_0)]}{x - x_0} = \frac{u(x, y_0) - u(x_0, y_0)}{x - x_0} + \frac{i[v(x, y_0) - v(x_0, y_0)]}{x - x_0},$$

so the limit  $f'(z_0)$  is equal to

$$\frac{\partial u}{\partial x}(x_0, y_0) + i \frac{\partial v}{\partial x}(x_0, y_0).$$

Why are we doing this? It's because if the limit exists, it should be the same whichever direction you approach it from, so you can derive some cool equalities.

The two equations must agree, so

$$\frac{1}{i} \frac{\partial u}{\partial y}(x_0, y_0) + \frac{\partial v}{\partial y}(x_0, y_0) = \frac{\partial u}{\partial x}(x_0, y_0) + i \frac{\partial v}{\partial x}(x_0, y_0).$$

Examine the real and imaginary parts, so we have

$$\frac{\partial v}{\partial y}(x_0, y_0) = \frac{\partial u}{\partial x}(x_0, y_0) \quad \text{and} \quad \frac{\partial v}{\partial x}(x_0, y_0) = -\frac{\partial u}{\partial y}(x_0, y_0). \quad (1)$$

These are known as the *Cauchy-Riemann Equations*. Furthermore,

$$f'(z_0) = \frac{\partial u}{\partial x}(x_0, y_0) + i \frac{\partial v}{\partial x}(x_0, y_0) = \frac{\partial v}{\partial y}(x_0, y_0) - i \frac{\partial u}{\partial y}(x_0, y_0).$$

What does this tell us? If your function is differentiable at a point, then we have a way to compute the derivative at that point. The converse does not hold, that is, if the Cauchy-Riemann equations hold this doesn't necessarily guarantee the existence of a derivative at that point.

**Example 6.1.** Recall that the function  $f(z) = |z|$  is nowhere differentiable. However, consider  $g(z) = |z|^2 = x^2 + y^2 = u(x, y) + iv(x, y)$ , where  $v(x, y) = 0$  and  $u(x, y) = x^2 + y^2$ . Let's check to see if this function satisfies the Cauchy-Riemann equations.  $\frac{\partial u}{\partial x} = 2x$ ,  $\frac{\partial u}{\partial y} = 2y$ ,  $\frac{\partial v}{\partial x} = 0$ ,  $\frac{\partial v}{\partial y} = 0$ . Does  $\frac{\partial u}{\partial x} = \frac{\partial v}{\partial y}$ ? Only if  $2x = 0 \implies x = 0$ . Does  $\frac{\partial u}{\partial y} = -\frac{\partial v}{\partial x}$ ? Only if  $2y = 0 \implies y = 0$ . So this function only satisfies the Cauchy-Riemann equations at the origin, which implies that the function is nowhere differentiable at any other point ( $g'(z_0)$  does not exist if  $z_0 \neq 0$ ). Now it does satisfy the CR equations at 0, but we don't have the existence of the derivative guaranteed.

Let's check the case if  $z_0 = 0$ .

$$\frac{g(z) - g(0)}{z - 0} = \frac{|z|^2 - 0}{z - 0} = \frac{\bar{z}z}{z} = \bar{z}.$$

Does  $\lim_{z \rightarrow 0} \bar{z}$  exist? Yes, and it's equal to 0. So  $g'(0)$  exists and is equal to 0.

## §6.2 Weak Converse of CR Equations

Let's talk about the opposite of the CR equations.

**Theorem 6.1** (Weak Converse of Cauchy-Riemann Equations). *Let  $f = u + iv$  be defined on a neighborhood of  $z_0 = x_0 + iy_0$ . Suppose the partial derivatives of  $u$  and  $v$  exist in that neighborhood, and are continuous at  $z_0$ . Furthermore, suppose the functions  $u$  and  $v$  satisfy the CR-equations at  $z_0$ . Then  $f'(z_0)$  exists.*

**Note.** We claim the hypotheses hold for  $|z|^2 : u(x, y) = u^2 + y^2, v(x, y) = 0$ . Oops, I went to the restroom here. I don't think I missed anything interesting though.

Now the next topic is very important.

**Example 6.2.** Let  $f(x, y) = e^x(\cos(y) + i \sin(y)) = e^x \cos(y) + ie^x \sin(y)$ . Note:  $u$  and  $v$  are nice. Let's compute the CR equations:  $\frac{\partial u}{\partial x} = e^x \cos(y) = \frac{\partial v}{\partial y} = e^x \cos(y)$ . We also have  $\frac{\partial u}{\partial y} = -e^x \sin(y) = -\frac{\partial v}{\partial x} = -e^x \sin(y)$ . Then  $f$  is differentiable everywhere, furthermore,

$$f' = \frac{\partial u}{\partial x} + i \frac{\partial v}{\partial x} = e^x \cos(y) + ie^x \sin(y) = e^x(\cos(y) + i \sin(y)) = f.$$

So  $f$  is equal to its derivative everywhere. This is probably the single most important function in the entire course.

We will eventually denote this function as  $\exp(z)$ . Note: if we use Euler's formula  $e^{i\theta} = \cos \theta + i \sin \theta$ , then

$$\exp(z) = e^x e^{iy} = e^{x+iy} = e^z.$$

But we have to make sure we can add the exponents first.

## §6.3 CR Equations in Polar Coordinates

Before discussing this further, consider polar coordinates for  $z$ . For any function  $g$ , write  $f(z) = u(r, \theta) + iv(r, \theta)$ . Then after the change of coordinates we have

$$\frac{\partial u}{\partial r} = \frac{\partial u}{\partial x} \frac{\partial x}{\partial r} + \frac{\partial u}{\partial y} \frac{\partial y}{\partial r}$$

and

$$\frac{\partial u}{\partial \theta} = \frac{\partial u}{\partial x} \frac{\partial x}{\partial \theta} + \frac{\partial u}{\partial y} \frac{\partial y}{\partial \theta}.$$

where  $x = r \cos \theta, y = r \sin \theta$ . We have  $\frac{\partial x}{\partial r} = \cos \theta, \frac{\partial x}{\partial \theta} = -r \sin \theta, \frac{\partial y}{\partial r} = \sin \theta, \frac{\partial y}{\partial \theta} = r \cos \theta$ . Use these to get

$$\frac{\partial u}{\partial r} = \frac{\partial u}{\partial x} \cos \theta + \frac{\partial u}{\partial y} \sin \theta$$

and

$$\frac{\partial u}{\partial \theta} = -\frac{\partial u}{\partial x} r \sin \theta + \frac{\partial u}{\partial y} r \cos \theta.$$

If  $f$  is differentiable,  $\frac{\partial u}{\partial x} = \frac{\partial v}{\partial y}, \frac{\partial u}{\partial y} = -\frac{\partial v}{\partial x}$ . Plugging these into the CR equations, we get

$$\frac{\partial u}{\partial r} = \frac{1}{r} \frac{\partial v}{\partial \theta}, \quad \frac{\partial v}{\partial r} = -\frac{1}{r} \frac{\partial u}{\partial \theta}, \quad \text{stop moving the page}$$

Next time: no. We'll do it next time. We have a test in 2 weeks BTW.

## §7 September 17, 2020

### §7.1 CR Equations (cont)

Last time: Cauchy Riemann equations for  $f = u + iv = u(x, y) + iv(x, y)$ . They are

$$\frac{\partial u}{\partial x} = \frac{\partial v}{\partial y}, \quad \frac{\partial u}{\partial y} = -\frac{\partial v}{\partial x}.$$

Also discussed this for polar coordinates, yada yada.

**Example 7.1.** Let  $f(z) = \frac{1}{z^4}$ .

$$\frac{1}{z^4} = \frac{1}{r^4 e^{i4\theta}} = \frac{1}{r^4} e^{-i4\theta} = \frac{1}{r^4} [\cos(4\theta) - i \sin(4\theta)].$$

So

$$\begin{aligned} f' &= e^{-i\theta} \left( -\frac{4}{r^5} \cos(4\theta) + i \frac{4}{r^5} \sin(4\theta) \right) \\ &= -\frac{4}{r^5} e^{-i\theta} (e^{-i4\theta}) \\ &= -\frac{4}{r^5} e^{-5i\theta} = -\frac{4}{r^5 e^{5i\theta}} \\ &= -\frac{4}{z^5}, \end{aligned}$$

which is a known formula for the derivative.

### §7.2 Analytic Functions

**Definition 7.1** (Analytic). A function  $f$  is said to be analytic at  $z_0$  if  $f$  is differentiable at all  $z$  in some ball centered at  $z_0$ .  $f$  is said to be analytic on a set  $S$  if for all  $z_0 \in S$ ,  $f$  is analytic at  $z_0$ .

**Example 7.2.** Let  $f(z) = z^m$ ,  $m \in \mathbb{N}$ . Then  $f'(z_0) = mz_0^{m-1}$  for all  $z_0$ . So such  $f$  are analytic in  $\mathbb{C}$ . If  $m \in \mathbb{Z} \setminus \mathbb{N}$ , this formula still holds for  $z_0$  nonzero. So  $f$  is analytic on the punctured plane  $\mathbb{C} \setminus \{0\}$ .

**Definition 7.2** (Entire function). We say a function  $f$  is *entire* if  $f$  is analytic on  $\mathbb{C}$ . For example,  $f(z) = z^3$  is entire. More generally, all polynomials are entire.

**Theorem 7.1.** If  $f'(z) = 0$  for all  $z \in D$  a domain, then  $f$  is constant in  $D$ . (Is this weak Liouville's Theorem?)

*Proof.* We will show that for any pair  $z_1, z_2 \in D$ ,  $f(z_1) = f(z_2)$ . Let  $z_1, z_2 \in D$ , then there is some finite set of straight lines connecting  $z_1$  and  $z_2$  (what is this definition reeee). Consider  $f$  on a segment  $z = z(t)$ ,  $0 \leq t \leq 1$ . Then  $F(t) = f[z(t)]$ ,  $0 \leq t \leq 1$  which is equal to  $u[x(t), y(t)] + iv[x(t), y(t)]$ . So

$$\frac{dF}{dt} = \frac{\partial u}{\partial x} \frac{\partial y}{\partial t} + \frac{\partial u}{\partial y} \frac{\partial y}{\partial t} + i \left[ \frac{\partial v}{\partial x} \frac{\partial x}{\partial t} + \frac{\partial v}{\partial y} \frac{\partial y}{\partial t} \right].$$

By assumption,  $\frac{dF}{dz} = 0$  in  $D$ . We can write this in two ways:  $\frac{\partial u}{\partial x} + i \frac{\partial v}{\partial x} = \frac{\partial v}{\partial y} - i \frac{\partial u}{\partial y} = 0$ . So  $\frac{\partial u}{\partial x} = 0 = \frac{\partial v}{\partial x} = \frac{\partial u}{\partial y} = \frac{\partial v}{\partial y}$  in  $D$ , and so  $\frac{dF}{dt} = 0$ ,  $0 \leq t \leq 1$ .

Consider  $\operatorname{Re}[F(t)] = u[x(t), y(t)]$ . It follows that  $\frac{d}{dt} u(x(t), y(t)) = 0$ ,  $0 \leq t \leq 1$ . It follows that  $u(x(t), y(t))$  is constant, since

$$\int_0^1 \frac{d}{dt} u(x(t), y(t)) dt = 0.$$

Similarly,  $V(x(1), y(1)) - V(x(0), y(0)) = 0$ . So  $F(t) = f(z(t))$  has the same values at  $z_a$  and  $z_b$ . (What?? Why??) Once we get Liouville's theorem we will get a less bad proof. This proof was big bad.  $\square$

We will show later that if  $f = u + iv$  is analytic at  $z_0$ , then  $u(x, y)$  and  $v(x, y)$  have partial derivatives of all orders in a neighborhood of  $z_0$ . If a function is analytic, then it is infinitely differentiable: what?? Complex analysis is crazy.

### §7.3 Harmonic Equations

**Definition 7.3** (Harmonic). If  $u_{xx} + u_{yy} = 0$ ,  $u$  is *harmonic* in that nbd of  $z_0$ , similarly for  $v$ . Laplace equation.

**Definition 7.4.** We say  $v$  is a *harmonic conjugate* of  $u$  in some region  $D$  if  $u$  and  $v$  are harmonic in  $D$ , and  $u, v$  satisfy the CR equations.

**Note.** For some analytic function  $f$  its true that  $\overline{f(z)} = f(\bar{z})$ .

**Theorem 7.2** (The Reflection Principle). *“Apparently this is famous, but I’ve never used it” -Dr. Radin*  
Suppose  $f$  is analytic in a domain  $D$  which is symmetric WRT the  $x$ -axis. Then for all  $z \in D$ ,

$$\overline{f(z)} = f(\bar{z}) \iff f \text{ is real on the segment of the } x\text{-axis in } D.$$

**Note.** I’ve been taking less notes because I’m simultaneously doing my weekly Differential Equations quiz while T<sub>E</sub>Xing notes. Just wanted to say that



## §8 September 22, 2020

Last time: for any  $z = (x, y) \in \mathbb{C}$ ,  $\exp(z) = e^x[\cos y + i \sin y] = e^x e^{iy}$ .  $e^z = \text{Exp}(z) = \exp(z)$ . We showed this function is differentiable on  $\mathbb{C}$  and that its derivative is itself.



The product of complex numbers  $e^{z_1} e^{z_2} = (e^{x_1} e^{iy_1})(e^{x_2} e^{iy_2})$ . Since multiplication is commutative, we have  $(e^{x_1} e^{x_2})(e^{iy_1} e^{iy_2}) = e^{x_1+x_2} e^{i(y_1+y_2)} = e^{z_1+z_2}$ . This follows from our definitions, it's not an assumption.

**Corollary 8.1.**  $e^z e^{-z} = e^0 = 1$ . So  $e^{-z} = \frac{1}{e^z}$ . Also, for  $m = 1, 2, \dots$   $(e^z)^m = e^{mz}$ . This also holds for negative integers. Finally, by our differentiation rules,

$$\frac{d}{dz} e^{az^n} = naz^{n-1} e^{az^n}.$$

So far we've covered how to differentiate polynomials (or more generally, rational functions), and now we've added  $e^z$  to our arsenal. Let's introduce some more basic functions to our list. Why do we differentiate? This is a course in functions of a complex variable, differentiating them, integrating them, etc. (I wish we covered analytic continuity). The next set of functions are trig functions.

### §8.1 Trig functions

Recall that  $e^{ix} = \cos x + i \sin x$ ,  $e^{-ix} = \cos x - i \sin x$ , so  $\frac{e^{ix} + e^{-ix}}{2} = \cos x$ ,  $\frac{e^{ix} - e^{-ix}}{2i} = \sin x$ . (Not sure if I got the definitions right). We can extend these to the complex plane, we define  $\cos z = (e^{iz} + e^{-iz})/2$ ,  $\sin z = (e^{iz} - e^{-iz})/2i$  for all  $z$ <sup>1</sup>. So  $\frac{d}{dz} \cos z = (ie^{iz} - ie^{-iz})/2 = i^2 \sin z = -\sin z$ . Similarly,  $\frac{d}{dz} \sin z = (ie^{iz} + ie^{-iz})/2i = \cos z$ . So these formulas agree with their real analog. We write the definitions again for clarity:

**Definition 8.1.** We define the trigonometric functions  $\sin z$  and  $\cos z$  on  $\mathbb{C}$  as

$$\sin z = \frac{e^{iz} - e^{-iz}}{2i}, \quad \cos z = \frac{e^{iz} + e^{-iz}}{2}.$$

Now by our new definitions of trig functions,

$$\cos z + i \sin z = \frac{e^{iz} + e^{-iz}}{2} + i \frac{e^{iz} - e^{-iz}}{2i} = e^{iz}.$$

From our definition,

$$\sin(z_1 + z_2) = \frac{e^{i(z_1+z_2)} - e^{-i(z_1+z_2)}}{2i}.$$

We claim that this is equal to  $\sin z_1 \cos z_2 + \cos z_1 \sin z_2$ . This is just a bunch of tedious manual labor. I don't really want to type this out, but here I am. We have this equal to

$$\begin{aligned} & \left( \frac{e^{iz_1} - e^{-iz_1}}{2i} \right) \left( \frac{e^{iz_2} + e^{-iz_2}}{2} \right) + \left( \frac{e^{iz_1} + e^{-iz_1}}{2} \right) \left( \frac{e^{iz_2} - e^{-iz_2}}{2i} \right) \\ &= \frac{1}{4i} \left[ e^{i(z_1+z_2)} + e^{i(z_1-z_2)} - e^{i(-z_1+z_2)} - e^{i(-z_1-z_2)} + e^{i(z_1+z_2)} - e^{i(z_1-z_2)} + e^{i(z_1+z_2)} - e^{i(-z_1-z_2)} \right] \\ &= \frac{1}{4i} \left[ 2e^{i(z_1+z_2)} - 2e^{i(-z_1-z_2)} \right] \\ &= \frac{1}{2i} \left[ e^{i(z_1+z_2)} - e^{i(-z_1-z_2)} \right] \\ &= \sin(z_1 + z_2). \end{aligned}$$

I think I may have typed the second (long) equation incorrectly, but I am not in the mood for going back and double checking this. Manual labor should be reserved for homework (and even then, I am still unwilling to do it).

We have a special case:  $\sin(z + 2\pi) = \sin(z) \cos(2\pi) + \cos(z) \sin(2\pi) = \sin(z)$ . Clearly this generalizes to  $\sin(z + 2\pi n)$  for  $n \in \mathbb{Z}$ . So  $\sin$  is periodic.

**Definition 8.2** (Tangent). Let

$$\tan z = \frac{\sin z}{\cos z}.$$

Note that this isn't defined at  $\cos z = 0$ : when does this happen? I stopped taking notes here for a little bit.

<sup>1</sup>Note that  $e^{iz}$  is differentiable, and  $\frac{d}{dz} e^{iz} = ie^{iz}$ .

## §8.2 Hyperbolic trig functions

Let's look at another class of functions: the cool dudes, hyperbolic trig functions (sinh is pronounced "sinch", cosh is pronounced "coush", etc). This gives me good memories of my first Calculus class with Dr. Neal Brand at UNT.

**Definition 8.3.** We define the hyperbolic trig functions  $\cosh z$  and  $\sinh z$  as

$$\cosh z = \frac{e^z + e^{-z}}{2}, \quad \sinh z = \frac{e^z - e^{-z}}{2}.$$

Similarly,  $\tanh z$  is defined as

$$\tanh z = \frac{e^z - e^{-z}}{e^z + e^{-z}}.$$

## §9 September 24, 2020

Last time: hyperbolic trig functions. We have some functions and theorems to deal with them, like rational functions  $p/q$ , trig functions  $\cos z, \sin z, \tan z$ , and hyperbolic trig functions  $\cosh z, \sinh z$ , the exponential function  $\exp(z)$ , etc. Where these functions are defined they are analytic.

### §9.1 Logarithmic functions

Let  $f$  be a function. Then  $f(z) = w \iff z = f^{-1}(w)$ . This is only a function if  $f$  is onto. Consider the case where  $f(z) = e^z$  and  $e^z = w$ .

**Definition 9.1** (Logarithm). We define the functional inverse of the exponential function as the *logarithm*, that is,

$$\log w = \{z \mid e^z = w\},$$

that is,  $\log = \exp^{-1}$ . Suppose  $z = x + iy$ , so  $e^z = e^{x+iy} = e^x(\cos y + i \sin y) = w$ . Write  $w$  in polar form, so  $w = |w|e^{i \arg w}$ . What values of  $z$  give rise to this? We want  $e^x = |w|, e^{iy} = e^{i \arg w}$ .  $e^x = |w| \iff x = \ln |w|$ , so  $y \in \arg w$ . Therefore we have

$$\log w = \ln |w| + i \arg w$$

for  $w \neq 0$ . So this function is multivariate.

We want to do calculus with this function. We are not terribly interested in multivalued things (actual functions lmao). One way to get an actual function is to define

$$\text{Log } w = \ln |w| + i \text{Arg}(w)$$

for  $w \neq 0$ . But that's not enough: we also want functions to be differentiable, etc. Note that this function  $\text{Log}$  wouldn't be continuous on the negative real axis. If  $w \in S^1$ , then ... I stopped paying attention here. Because of the discontinuity (which I was not paying attention for), we define  $\text{Log}(w) = \ln |w| + i \arg w$  only for  $w$  not a negative real number.

**Claim.**  $\text{Log}$  satisfies the CR equations. Verify this in your free time.

Note that  $\frac{d}{dz} \text{Log } z = e^{-i\theta}(u_r + iv_r) = e^{-i\theta}(\frac{1}{r} + 0) = \frac{1}{z}$  on its domain. This is very cool, thank you logarithm. This is useful, but we need other ways to get "honest" functions from  $\log w = \ln |w| + i \arg w$ .

aa hayaku ouchi ni kaeritai

Recall the identity that  $\arg(z_1 z_2) = \arg z_1 + \arg z_2$ , with addition being componentwise for the infinite sets. So  $\log(z_1 z_2) = \ln |z_1 z_2| + i \arg(z_1 z_2) = \ln |z_1| + \ln |z_2| + i \arg z_1 + i \arg z_2 = \log z_1 + \log z_2$ .



Where are we headed? To the next class of functions  $z^\alpha$  and beyond.

## §10 October 1, 2020

Last time: we had a test. Today we'll talk about branches and next time move onto integration, finally. So I had a test until 11 today, and I probably won't take many (any) notes because I'm doing my homework (due at 2) during the lecture. (If anybody wants to send me notes, please do, my email is [simonxiang@utexas.edu](mailto:simonxiang@utexas.edu)).

### §10.1 Catching up: logarithms and branches

So I didn't pay attention for the last week and now I'm paying for it, because I have to take extra notes to stay on track (and finish the homework). A side note: if we think of branches as sheets and the logarithm only being defined on simply connected domains, this is surprisingly similar to the theory of covering spaces in algebraic topology. Maybe I can find a way to connect a fundamental group to the codomain of the logarithm... or am I just spouting nonsense?



Motivation: solve equations of the form  $e^w = z$ ,  $z$  is nonzero (or else the earth collapses). Write this as  $e^u e^{iv} = r e^{i\Theta}$  where  $w = u + iv$ ,  $\Theta = \text{Arg } \theta$ . Since  $r_1 e^{i\theta_1} = r_2 e^{i\theta_2} \iff r_1 = r_2, \theta_1 = \theta_2 + 2\pi n$  for some  $n \in \mathbb{Z}$ , we have

$$e^u e^{iv} = r e^{i\Theta} \iff e^u = r \text{ and } v = \Theta + 2\pi n$$

for some  $n \in \mathbb{Z}$ . Now  $e^u = r \implies u = \ln r$  in the traditional real valued sense. So  $e^w = z$  if and only if

$$w = \ln r + i(\Theta + 2\pi n)$$

for some  $n \in \mathbb{Z}$ . If we write  $\log z = \ln r + i(\Theta + 2\pi n)$ , then  $e^{\log z} = z$  for  $z \neq 0$ .

**Definition 10.1** (Multivalued logarithm). We have  $\log z$  for  $z \in \mathbb{C}, z \neq 0$  defined by

$$\log z = \ln r + i(\Theta + 2\pi n),$$

where  $n \in \mathbb{Z}, z = re^{i\Theta}, \Theta = \text{Arg } \theta$ .

**Example 10.1.** Let  $z = -1 - \sqrt{3}i$ , then  $r = 2$  and  $\Theta = -\frac{2\pi}{3}$ . So

$$\log(z) = \ln 2 + i - \left(\frac{2\pi}{3} + 2\pi n\right) = \ln 2 + 2\pi i \left(n - \frac{1}{3}\right)$$

for  $n \in \mathbb{Z}$ .

Note that  $\log(e^z) = z$  does not necessarily hold. We can write  $\log z = \ln |z| + i \arg z$ , which implies

$$\log(e^z) = \ln |e^z| + i \arg(e^z) = \ln(e^x) + i(y + 2\pi n) = (x + iy) + 2i\pi n,$$

since  $|e^z| = e^x$  and  $\arg(e^z) = y + 2\pi n$  for some  $n \in \mathbb{Z}$  (this can be seen by writing  $\exp z$  as  $e^x e^{iy}$ ). So

$$\log(e^z) = z + 2\pi in$$

for  $n$  an integer. I'm sick of this. From now on,  $n$  denotes an integer, that is, some  $n \in \mathbb{Z}$ . You can figure out when this abuse of quantifiers ends by context. We can define the principle value of  $\log z$  at  $n = 0$  by

$$\text{Log } z = \ln r + i\Theta.$$

Note that  $\text{Log } z$  is well-defined and single-valued when  $z \neq 0$ , furthermore,  $\log z = \text{Log } z + 2\pi in$ . If  $z \in \mathbb{R}$ , this is just the standard logarithm from calculus.

**Example 10.2.** Here's a cool trick: we can define the logarithm of negative numbers now (something we couldn't do in calculus), since  $\log(-1) = \ln 1 + i(\pi + 2\pi n) = i\pi(2n + 1)$ ,  $\text{Log}(-1) = i\pi$ .

## §10.2 Branches of the logarithm

If we restrict  $\theta$  such that for some  $\alpha \in \mathbb{R}, \alpha < \theta < \alpha + 2\pi$ , then  $\log z = \ln r + i\theta$  with components  $u(r, \theta) = \ln r$  and  $v(r, \theta) = \theta$  is single-valued and continuous in the stated domain (?). It's defined from the  $x$ -axis to the angle it makes with  $\alpha$ —note that it isn't defined on the ray  $\theta = \alpha$ , because some neighborhood of  $z$  (on such ray) will contain points near  $\alpha$  and  $\alpha + 2\pi$ . Not only is this restricted logarithm continuous, but it's also analytic on its domain, because of CR. Then by the derivative of polar stuff, we have

$$\frac{d}{dz} \log z = e^{-i\theta} (u_r + iv_r) = e^{-i\theta} \left( \frac{1}{r} + i\theta \right) = \frac{1}{re^{i\theta}}.$$

So  $\frac{d}{dz} \log z = \frac{1}{z}$  when  $|z| > 0, \alpha < \arg z < \alpha + 2\pi$ . In particular,  $\frac{d}{dz} \text{Log } z = \frac{1}{z}$  for  $|z| > 0, -\pi < \text{Arg } z < \pi$ .

A *branch* of a multivalued function  $f$  is any single-valued function  $F$  that's analytic in some domain  $\Omega$ , where for any  $z \in \Omega$  we have  $F(z)$  one of the values of  $f$ . Whoever wrote this textbook needs to stop overusing references, please. Note that for  $\alpha \in \mathbb{R}$ ,  $\log$  restricted to  $\alpha$  is a branch of the general logarithm. We say  $\text{Log } z = \ln r + i\Theta$  for  $r > 0, -\pi < \Theta < \pi$  is the principal branch. A *branch cut* is a portion a line or curve that is introduced in order to define a branch  $F$  of a multivalued function  $f$ . Point on the branch cut are singular (have no well-defined nbd) and any point common to every branch cut of  $f$  is a branch point. For example, the branch cut for the principal branch is the origin plus the ray  $\Theta = \pi$ , and the origin is a branch point for  $\log$ .

**Example 10.3.** Take the principle branch  $\text{Log } z = \ln r + i\Theta$ . Then  $\text{Log}(i^3) = \text{Log}(-i) = \ln 1 - i\frac{\pi}{2} = -i\frac{\pi}{2}$ , but  $3 \text{Log } i = 3(\ln 1 + i\frac{\pi}{2}) = i\frac{3\pi}{2}$ . So  $\text{Log}(i^3) \neq 3 \text{Log } i$ .

## §10.3 Logarithmic identities

These derivations aren't interesting IMO. If  $z_1, z_2 \in \mathbb{C}$ , we have

$$\log(z_1 z_2) = \log z_1 + \log z_2, \tag{2}$$

$$\log\left(\frac{z_1}{z_2}\right) = \log z_1 - \log z_2. \tag{3}$$

Let  $z \in \mathbb{C}$ . Then if you write  $z = re^{i\theta}$ , it can be seen that

$$z^n = e^{n \log z}. \tag{4}$$

Furthermore, for  $z \neq 0, k \in \mathbb{N}$ , we have

$$z^{1/k} = \exp\left(\frac{1}{k} \log z\right), \tag{5}$$

$$\exp\left(\frac{1}{k} \log z\right) = \sqrt[k]{r} \exp\left[i\left(\frac{\Theta}{k} + \frac{2\pi n}{k}\right)\right]. \tag{6}$$

## §10.4 Complex exponents

**Definition 10.2** (Complex exponential function). For  $z \neq 0$ ,  $c \in \mathbb{C}$ , we have the function  $z^c$  defined as

$$z^c = e^{c \log z}. \quad (7)$$

We already know this holds for  $c = n$  or  $c = \frac{1}{n}$ . Usually powers of  $z$  are multivalued.

**Example 10.4.** We have  $i^{-2i} = \exp(-2i \log i)$ , where  $\log i = \ln 1 + i\left(\frac{\pi}{2} + 2\pi n\right) = i\pi\left(2n + \frac{1}{2}\right)$ . Then  $i^{-2i} = \exp[\pi(4n + 1)]$ . Note that every value of  $i^{-2i}$  lies in  $\mathbb{R}$ .

Since  $1/e^z = e^{-z}$ , then  $\frac{1}{z^c} = \frac{1}{\exp(c \log z)} = \exp(-c \log z) = z^{-c}$ . So  $1/i^{2i} = i^{-2i}$ , and we have  $\frac{1}{2i} = \exp[\pi(4n + 1)]$ . For  $z = re^{i\theta}$  and  $\alpha \in \mathbb{R}$ , the branch  $\log z$  of  $\alpha$  is single-valued and analytic on its domain. Using that branch, it follows that  $z^c = \exp(c \log z)$  is also single-valued and analytic on such domain. Then the derivative of that branch of  $z^c$  is given by

$$\frac{d}{dz} z^c = \frac{d}{dz} \exp(c \log z) = \frac{c}{z} \exp(c \log z) = c \frac{\exp(c \log z)}{\exp(\log z)} = c \exp[(c-1) \log z] = cz^{c-1},$$

for  $|z| > 0$ ,  $\alpha < \arg z < \alpha + 2\pi$ . The principle value is what you think it is: P.V.  $z^c = e^{c \text{Log } z}$ , and so is the principal branch of  $z^c$ , when  $|z| > 0$ ,  $-\pi < \text{Arg } z < \pi$ .

**Example 10.5.** The principal value of  $(-i)^i$  is

$$\exp[i \text{Log}(-i)] = \exp\left[i\left(\ln 1 - i\frac{\pi}{2}\right)\right] = \exp \frac{\pi}{2},$$

that is, P.V.  $(-i)^i = \exp \frac{\pi}{2}$ .

**Example 10.6.** The principle branch of  $z^{2/3}$  can be written as

$$\exp\left(\frac{2}{3} \text{Log } z\right) = \exp\left(\frac{2}{3} \ln r + \frac{2}{3} i\Theta\right) = \sqrt[3]{r^2} \exp\left(i\frac{2\Theta}{3}\right).$$

So P.V.  $z^{2/3} = \sqrt[3]{r^2} \cos \frac{2\Theta}{3} + i\sqrt[3]{r^2} \sin \frac{2\Theta}{3}$ .

**Example 10.7.** Let  $z_1 = 1 + i$ ,  $z_2 = 1 - i$ ,  $z_3 = -1 - i$ . Then  $(z_1 z_2)^i = 2^i = e^{i \ln 2}$ , and

$$\begin{aligned} z_1^i &= e^{i \text{Log}(1+i)} = e^{i(\ln \sqrt{2} + i\pi/4)} = e^{-\pi/4} e^{i(\ln 2)/2}, \\ z_2^i &= e^{i \text{Log}(1-i)} = e^{i(\ln \sqrt{2} - i\pi/4)} = e^{\pi/4} e^{i(\ln 2)/2}. \end{aligned}$$

So  $(z_1 z_2)^i = z_1^i z_2^i$  as expected. But do some similar stuff with  $(z_2 z_3)^i = e^{-\pi} e^{i \ln 2}$ ,  $z_3^i = e^{3\pi/4} e^{i(\ln 2)/2}$ , and we find that  $z_2^i z_3^i = e^{2\pi} (z_2 z_3)^i$ .

We have the exponential with base  $c$  defined as

$$c^z = e^{z \log c}$$

for  $c$  a nonzero constant in  $\mathbb{C}$ . When we specify a value for  $\log c$ , this function is entire, and

$$\frac{d}{dz} c^z = \frac{d}{dz} e^{z \log c} = e^{z \log c} \log c = c^z \log c.$$

## §11 October 6, 2020

### §11.1 Parametrized curves

Let's talk about integration! Define  $I = \int f(z) dz$  for parametrized curves  $\Gamma$ , given by  $w: [a, b] \rightarrow \mathbb{C}$ . We say the set of points

$$\{w(t) \mid a \leq t \leq b\}$$

is the "trace" of the curve  $w$ , denoted  $\text{tr } w$ <sup>2</sup>.

**Example 11.1.** Let  $w(t) = e^{i2\pi t}$  for  $t \in [0, \frac{1}{2}]$ . Unfortunately, I'm not cool enough to live-TeX figures in class, so try to use your imagination to see what this curve would look like (a semicircle). Also consider  $w(t) = (1+i)t$  for  $t \in [1, 3]$ . This one looks like a straight line.

**Example 11.2.** Consider  $w(t) = e^{it}$  for  $t \in [0, 4\pi]$ . What is  $\text{tr } w$ ? It's simply the unit circle.

Given a parametrized  $\Gamma$  and a function  $f(z)$  defined on at least  $\text{tr } \Gamma$ , we define  $\int_{\Gamma} f(z) dz$  by a limit of Riemann sums (inb4 not as powerful as integration by Lebesgue measure). I know we never defined what  $t$  is, but in each interval  $[t_j, t_{j+1}]$  pick some  $\hat{t}_{j+i}$  and compute  $f(\hat{t}_1)[w(t_1) - w(t_0)] + f(\hat{t}_2)[w(t_2) - w(t_1)] + \dots$ . Using summation notation, we have the sum written as

$$\sum_{j=1}^m f(w(\hat{t}_j)) [w(t_j) - w(t_{j-1})].$$

**Example 11.3.** Consider  $w(t) = e^{i2\pi t}$  for  $t \in [0, 1]$ , and  $f(z) = z^3$ . Choose  $t_j = \frac{j}{m}$ , so  $t_0 = 0$  and  $t_m = 1$ . Let  $\hat{t}_j = t_j$ . Consider the approximations of  $\int_{\Gamma} z^3 dz$ , given by

$$\begin{aligned} \sum_{j=1}^m f[w(t_j)] [w(t_j) - w(t_{j-1})] &= \\ \sum_{j=1}^m f[e^{i2\pi t_j}] [e^{i2\pi t_j} - e^{i2\pi t_{j-1}}] &= \\ \sum_{j=1}^m f[e^{i2\pi \frac{j}{m}}] [e^{\frac{i2\pi j}{m}} - e^{\frac{i2\pi(j-1)}{m}}] &= \\ \sum_{j=1}^m f[e^{i2\pi \frac{j}{m}}] [e^{\frac{i2\pi j}{m}} - e^{\frac{i2\pi(j-1)}{m}}] &= \dots \end{aligned}$$

Unfortunately I was too slow to finish the work. This simplifies to

$$\left(1 - e^{\frac{2\pi i}{m}}\right) \sum_{j=1}^m \left(e^{i\frac{8\pi}{m}}\right)^j,$$

which is a geometric series of the form  $S = \sum_{j=K}^L a^j$  and are easy to compute. We have  $aS = S - a^K + a^{L+1}$ , so  $(1-a)S = a^K - a^{L+1} \implies S = \frac{a^K - a^{L+1}}{1-a}$ . This proof is pretty much the same as any one in a calculus course. For  $S = \sum_{j=1}^m e^{\left(\frac{i8\pi}{m}\right)^j}$ , we have

$$S = \frac{e^{\frac{i8\pi}{m}} - \left(e^{i\frac{8\pi}{m}}\right)^{m+1}}{1 - e^{i\frac{8\pi}{m}}} \quad (8)$$

I missed something else big, gotta get faster. So  $\int_{\Gamma} z^3 dz = 0$ .

<sup>2</sup>Wait, I'm not sure all of a sudden. If this notation is non-standard call me a fool, but this is how it works in linear algebra.

## §12 Holomorphic Functions

I want to do some real math! These notes will follow Stein and Shakarchi §1.2.

### §12.1 Continuous functions

We've already seen the standard epsilon-delta definition of continuity. An equivalent definition is the sequential definition, that is, for every sequence  $\{z_1, z_2, \dots\} \subseteq \Omega \subseteq \mathbb{C}$  such that  $\lim z_n = z_0$ , then  $f$  is continuous at  $z_0$  if  $\lim f(z_n) = f(z_0)$ . Since the notions for convergence of complex numbers and  $\mathbb{R}^2$  is the same,  $f$  of  $z = x + iy$  is continuous iff it's continuously viewed as a function of two real variables  $x$  and  $y$ . If  $f$  is continuous, then the real valued function defined by  $z \mapsto |f(z)|$  is clearly continuous (by the triangle inequality).

We say  $f$  attains a *maximum* at the point  $z_0 \in \Omega$  if

$$|f(z)| \leq |f(z_0)| \text{ for all } z \in \Omega.$$

The definition of a minimum is what you think it is.

**Theorem 12.1.** A continuous function on a compact set  $\Omega$  is bounded and attains a maximum and minimum on  $\Omega$ .

*Proof.* Same as the any one you'd find in a Real Analysis course. □

### §12.2 Holomorphic functions

Let's talk about the good stuff. Let  $\Omega \subseteq \mathbb{C}$  be open and  $f: \Omega \rightarrow \mathbb{C}$ . Then  $f$  is *holomorphic at the point*  $z_0 \in \Omega$  if the quotient

$$\frac{f(z_0 + h) - f(z_0)}{h}$$

converges to a limit when  $h \rightarrow 0$ . Here  $h \in \mathbb{C}$  and  $h \neq 0$  with  $z_0 + h \in \Omega$ , such that the quotient is well-defined. This limit is called the *derivative of  $f$*  and  $z_0$ , and is denoted by

$$f'(z_0) = \lim_{h \rightarrow 0} \frac{f(z_0 + h) - f(z_0)}{h}.$$

Note that  $h$  approaches 0 from any direction. The function  $f$  is said to be *holomorphic on  $\Omega$*  if  $f$  is holomorphic at every point of  $\Omega$ . If  $C \subseteq \mathbb{C}$  is closed, then  $f$  is *holomorphic on  $C$*  if  $f$  is holomorphic on some open set containing  $C$ . Finally, if  $f$  is holomorphic on  $\mathbb{C}$  then  $f$  is said to be *entire*. Sometimes the terms *regular* or *complex differentiable* are used in place of holomorphic, but holomorphic functions are much much nicer than real values differentiable functions. Furthermore, every holomorphic function is analytic (that is, it has a power series expansion near every point), and so we also use the term *analytic* to refer to holomorphic functions. Once again, things are not as nice in Real Analysis, with infinitely differentiable real valued functions not having power series expansions.

**Example 12.1.** We have any polynomial  $p(z) = a_0 + a_1z + \dots + a_nz^n$  entire, and  $f(z) = \frac{1}{z}$  holomorphic on the punctured plane  $\mathbb{C} \setminus \{0\}$ . However,  $f(z) = \frac{\bar{z}}{z}$  is not entire, as  $\frac{f(z_0+h)-f(z_0)}{h} = \frac{\bar{h}}{h}$ , which has no limit as  $h \rightarrow 0$ .

If we write the definition of a holomorphic function as  $f$  being holomorphic at  $z_0 \in \mathbb{C}$  iff there exists an  $a \in \mathbb{C}$  such that

$$f(z_0 + h) - f(z_0) - ah = h\psi(h),$$

where  $\psi$  is a function defined for all "small"  $h$ , and  $\lim_{h \rightarrow 0} \psi(h) = 0$ , we can see that  $f$  is holomorphic implies  $f$  is continuous (clearly  $a = f'(z_0)$ ). The basic stuff, distribution over addition, product rule, quotient rule, chain rule, yada yada all apply.

### §12.3 Complex-valued functions as mappings

OK, here's the difference between real and complex valued functions again. In terms of real variables,  $f(z) = \bar{z}$  corresponds to  $F: (x, y) \rightarrow (x, -y)$ , which is differentiable in the real sense, its derivative at a point being the map corresponding to its Jacobian. In fact,  $F$  is linear and is equal to its derivative at any point, and is therefore infinitely differentiable. So a complex valued function having a real derivative need not imply the complex valued function is holomorphic.