AQ7.3: Activity Questions 3 - Not Graded

This assignment will not be graded and is only for practice.

Level 1

If $||a||$ = 3||a|| = 3, $||b||$ = 2√2||b|| = 2 2 √ and angle between aa and bb is 45° 45∘, then the value of $a.b$a.b is

No, the answer is incorrect.

Score: 0

Accepted Answers:

(Type: Numeric) 6

1 point

1 point

Given that $\| a \|$ = 3‖a‖ = 3 and $\| b \|$ = 5‖b‖ = 5 and $a \cdot b$ = 7.5a · b = 7.5, for two vectors aa and bb in R^n Rn. Find the angle (in degrees) between the two vectors a, ba, b. Note: cos60° = 1/2, cos30° = √3/2cos 60∘ = 1/2, cos 30∘ = 3 √ /2.

- 30° 30∘
- 60° 60∘
- 120° 120∘
- 150° 150∘

No, the answer is incorrect.

Score: 0

Accepted Answers:

60° 60∘

1 point

Consider two vectors a = (3, 4, − 1), b = (2, − 1, 2)a = (3, 4, −1), b = (2, −1, 2) in R^3 R3. Choose the correct options.

- [ ] Length/Norm of aa is √26 26 √ and that of bb is 33.
- [ ] Length/Norm of aa is 55 and that of bb is 33.
- [ ] Length/Norm of aa is √24 24 √ and that of bb is √7 7 √ .
- [ ] Angle between the two vectors is 9090 degrees.

☐
Angle between the two vectors is approximately 00 degrees.

No, the answer is incorrect.

Score: 0

Accepted Answers:

Length/Norm of aa is $\sqrt{26}$ 26

√

and that of bb is 33.

Angle between the two vectors is 9090 degrees.

1 point

If the dot product of two vectors is zero, then choose the correct option.

○ The two vectors are perpendicular to each other.
○ The two vectors are parallel to each other.
○ The two vectors are exactly the same.
○ The length of the two vectors must be the same.

No, the answer is incorrect.

Score: 0

Accepted Answers:

The two vectors are perpendicular to each other.

Level 2

If $a = (6, -1, 3), b = (4, c, 2)$a = (6, −1, 3), b = (4, c, 2) where $a, b \in R^3$ a, b ∈ R3 and aa and bb are perpendicular to each other, then find the value of cc?

No, the answer is incorrect.

Score: 0

Accepted Answers:

(Type: Numeric) 30

1 point

1 point

Consider two vectors $a = (1, 2), b = (2, 2)$a = (1, 2), b = (2, 2) and θθ is the angle between them. The sum of the two vectors is given by $c = a + b$c = a + b. Choose the correct options.

☐
Length/Norm of cc is 55.

☐
Length/Norm of cc is 2525.

☐
Length/Norm of cc is 44.

☐
$\cos\theta = \frac{3}{\sqrt{10}}$cos θ = 10

√

3.

☐
$\cos\theta = \frac{3}{\sqrt{5}}$cos θ = 5

√

3.

No, the answer is incorrect.

Score: 0

Accepted Answers:

Length/Norm of cc is 55.
$\cos\theta = \frac{3}{\sqrt{10}}$cos θ = 10

√

3.
1 point
Consider 33 vectors a, b, ca, b, c in R^3 R3 and a scalar λλ in RR. Choose the set of correct options.
Note: (.)(.) represents the dot product.

- [ ] $\lambda(a \cdot b) = (\lambda a) \cdot b$λ(a · b) = (λa) · b
- [ ] $\lambda(a \cdot b) = (\lambda b) \cdot a$λ(a · b) = (λb) · a
- [ ] $\lambda(a \cdot b) = (\lambda a) \cdot (\lambda b)$λ(a · b) = (λa) · (λb)
- [ ] $(a + c) \cdot b = a \cdot b + c \cdot b$(a + c) · b = a · b + c · b
- [ ] $(a + c) \cdot b = a \cdot c + c \cdot b$(a + c) · b = a · c + c · b
- [ ] $a \cdot a = 0, b \cdot b = 0$a · a = 0, b · b = 0 if and only if a, ba, b are null vectors.

No, the answer is incorrect.
Score: 0
Accepted Answers:
$\lambda(a \cdot b) = (\lambda a) \cdot b$λ(a · b) = (λa) · b
$\lambda(a \cdot b) = (\lambda b) \cdot a$λ(a · b) = (λb) · a
$(a + c) \cdot b = a \cdot b + c \cdot b$(a + c) · b = a · b + c · b
$a \cdot a = 0, b \cdot b = 0$a · a = 0, b · b = 0 if and only if a, ba, b are null vectors.

1 point
Consider two vectors $a = (1, 3, 5, 7, 9)$a = (1, 3, 5, 7, 9) and $b = (2, 4, 6, 8, 10)$b = (2, 4, 6, 8, 10). Choose the set of correct options.

- [ ] The length of bb is more than aa.
- [ ] The length of aa is more than bb.
- [ ] The length of $(a - b)$(a − b) is $\sqrt{5}$ 5

√

.

- [ ] The length of $(a - b)$(a − b) is $\sqrt{50}$ 50

√

.

No, the answer is incorrect.
Score: 0
Accepted Answers:
The length of bb is more than aa.
The length of $(a - b)$(a − b) is $\sqrt{5}$ 5

√

.

Check Answers

Your score is: 0/8