This assignment will not be graded and is only for practice.

4.1 Level 1:

1) Let A be a 3×3 matrix with non-zero determinant. If $det(2A) = k\ det(A)$, then what will be the value of k?
[Hint: If a scalar (c) is multiplied with one row of a matrix A, then the determinant of the new matrix will be c times the determinant of A.]

No, the answer is incorrect.

Score: 0

Accepted Answers:

(Type: Numeric) 8

1 point

2) If $A = \begin{bmatrix} 2 & -1 & 1 \\ 1 & -3 & 4 \\ -1 & 0 & 2 \end{bmatrix}$ and $B = \begin{bmatrix} 1 & 3 & -2 \\ 0 & 1 & -1 \\ 3 & 4 & 2 \end{bmatrix}$, then choose the set of correct options.

1 point

- [ ] $det(A) = -9$ and $det(B) = -3$.
- [ ] $det(A) = 9$ and $det(B) = 3$.
- [ ] $det(A) = -9$ and $det(B) = 3$.
- [ ] $det(AB) = -27$ and $det(BA) = 27$.
- [ ] $det(AB) = det(BA) = -27$.

No, the answer is incorrect.

Score: 0

Accepted Answers:

$det(A) = -9$ and $det(B) = 3$.
$det(AB) = det(BA) = -27$.

3)

1 point

Let A be a 2×2 matrix, which is given as $\begin{bmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{bmatrix}$. Define the following matrices :

$$B = \begin{bmatrix} a_{11} - a_{21} & a_{12} - a_{22} \\ a_{21} & a_{22} \end{bmatrix}, C = \begin{bmatrix} a_{11} - a_{12} & a_{12} \\ a_{21} - a_{22} & a_{22} \end{bmatrix},$$

$$D = \begin{bmatrix} a_{11} + a_{21} & a_{12} - a_{22} \\ a_{21} & a_{22} \end{bmatrix}, E = \begin{bmatrix} a_{11} - a_{21} & a_{12} + a_{22} \\ a_{21} & a_{22} \end{bmatrix}$$

Which of the matrices among $B, C, D,$ and E have the same determinant as that of the matrix A, for any real numbers $a_{11}, a_{12}, a_{21}, a_{22}$?

- B and D
- B and E
- B and C
- D and E
- C and E

No, the answer is incorrect.

Score: 0

Accepted Answers:

B and C

4) Let $A = \begin{bmatrix} a_{11} & a_{12} & a_{13} \\ ta_{11} - sa_{31} & ta_{12} - sa_{32} & ta_{13} - sa_{33} \\ ra_{31} & ra_{32} & ra_{33} \end{bmatrix}$ be a matrix and $r, s, t = 0$. Find $det(A)$

No, the answer is incorrect.

Score: 0

Accepted Answers:

(Type: Numeric) 0

1 point

5) Suppose $A = \begin{bmatrix} -1 & 4 & -2 \\ 0 & 1 & 3 \\ 1 & 0 & -2 \end{bmatrix}$ and $B = \begin{bmatrix} 2 & 0 & -1 \\ 4 & \alpha & 0 \\ 0 & -3 & -5 \end{bmatrix}$. If $det(AB) = 32$, then find the value of α.

No, the answer is incorrect.

Score: 0

Accepted Answers:

(Type: Numeric) 1

1 point

4.2 Level 2:

(Use this information to answer questions 6 and 7)

Let $A(x_1, y_1), B(x_2, y_2)$ and $C(x_3, y_3)$ be the vertices of a triangle. The area of the triangle ABC is given by $\frac{1}{2}\left|det(D)\right|$ square units, where $D = \begin{bmatrix} x_1 & y_1 & 1 \\ x_2 & y_2 & 1 \\ x_3 & y_3 & 1 \end{bmatrix}$

6) If $P(-1, 1), Q(1, 1)$ and $R(1, 0)$ are the vertices of a triangle, then find the area (in square units) of the triangle PQR.

No, the answer is incorrect.

Score: 0

Accepted Answers:

(Type: Numeric) 1

1 point

7) Let $P(x-1, 2x), Q(0, x-2)$ and $R(-x+1, 1)$ be the vertices of a triangle. If the area of the triangle is 5 square units and $x > 0$, then choose the set of correct options ***1 point***

- [ ] length of side $PQ = \sqrt{29}$ units.
- [ ] length of side $PR = \sqrt{40}$ units.
- [ ] length of side $QR = 2$ units.
- [ ] length of side $QR = 1$ units.

No, the answer is incorrect.

Score: 0

Accepted Answers:

length of side $PQ = \sqrt{29}$ units.
length of side $QR = 2$ units.

8) Find out the correct value of $det(A)$, where $A = \begin{bmatrix} 1 & 2010 & 2020 \times 2030 \\ 1 & 2020 & 2030 \times 2010 \\ 1 & 2030 & 2010 \times 2020 \end{bmatrix}$ ***1 point***

[Hint: Observe that the given matrix is of the form: $\begin{bmatrix} 1 & a & bc \\ 1 & b & ca \\ 1 & c & ab \end{bmatrix}$]

- 0
- 200
- -2000
- 2000

No, the answer is incorrect.

Score: 0

Accepted Answers:

2000

9) Let A be a square matrix of order 3 and B be a matrix that is obtained by adding 2 times the first row of A to the third row of A and adding 3 times the second row of A to the first row of A. What is the value of $det(6A^2B^{-1})$? ***1 point***

- $6\ det(A)$
- $6\ det(A)det(B)$
- $6^3\ det(A)$
- $6^3\ det(A)det(B)$

No, the answer is incorrect.

Score: 0

Accepted Answers:

$6^3\ det(A)$

10) Suppose $A = \begin{bmatrix} 19 & 20 & 21 \\ 20 & 21 & 19 \\ 21 & 19 & 20 \end{bmatrix}$. Find the value of $\frac{1}{60} det(A^2)$.

[You can try to compute the determinant of a matrix of the form: $\begin{bmatrix} a & b & c \\ b & c & a \\ c & a & b \end{bmatrix}$]

No, the answer is incorrect.

Score: 0

Accepted Answers:

(Type: Numeric) 540

1 point

Your score is: 0/10